

Two-stage Stochastic Variational Inequalities

Xiaojun Chen

Department of Applied Mathematics
Hong Kong Polytechnic University

The Greater Bay Area Optimization Day

24 February, 2018

2017

1. X. Chen and R.J-B. Wets, Stochastic Equilibrium and Variational Inequalities, a special issue of Math. Program., 165(2017).
2. X. Chen, T.K. Pong and R. J-B Wets, Two-stage stochastic variational inequalities: an ERM-solution procedure, Math. Program., 165(2017), 71-111.
3. R.T. Rockafellar and R. J.-B. Wets, Stochastic variational inequalities: single-stage to multistage, Math. Program., 165(2017), 331-360.
4. R.T. Rockafellar and J. Sun, Solving monotone stochastic variational inequalities and complementarity problems by progressive hedging, (2017), to appear in Math. Program.
5. X. Chen, H. Sun and H. Xu, Discrete approximation of two-stage stochastic and distributionally robust linear complementarity problems, (2017), revised version for Math. Program.
6. X. Chen, A. Shapiro and H. Sun, Convergence analysis of sample average approximation of two-stage stochastic generalized equations, submitted (2017).

1.1 Variational Inequalities (VI)

Given a nonempty closed-convex set $X \subseteq \mathbb{R}^n$ and a continuous function $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$, the variational inequality problem is to find $x \in X$ such that

$$-F(x) \in \mathcal{N}_X(x)$$

$$\text{i.e.} \quad (y - x)^T F(x) \geq 0, \quad \forall y \in X.$$

$\mathcal{N}_X(x)$ is the normal cone to the set X at x .

Complementarity problem as a special case: $X = \mathbb{R}_+^n$

$$-F(x) \in \mathcal{N}_{\mathbb{R}_+^n}(x), \quad 0 \leq x \perp F(x) \geq 0$$

System of equations as a special case: $X = \mathbb{R}^n$

$$F(x) = 0$$

Stochastic variational inequalities

Single stage stochastic variational inequalities

A random variable ξ affects the function F and the set X .

$\xi \in \Xi \subseteq \mathbb{R}^L$, a set representing future states of knowledge.

Given $F : \Xi \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ and $X_\xi \subset \mathbb{R}^n$, find $x \in X_\xi$ such that

$$-F(\xi, x) \in \mathcal{N}_{X_\xi}(x), \quad \text{i.e.,} \quad (y - x)^T F(\xi, x) \geq 0, \quad \forall y \in X_\xi.$$

This problem is well defined if ξ is known. “Wait-and-see”

Example

Consider $f : \Xi \times \mathbb{R}^n \rightarrow \mathbb{R}$ and $g : \Xi \times \mathbb{R}^n \rightarrow \mathbb{R}^m$

$$\text{minimize } f(\xi, x) \quad \text{subject to } x \in X_\xi$$

where

$$X_\xi = \{x \in \mathbb{R}_+^n \mid g(\xi, x) \geq 0\},$$

$-\nabla f(\xi, x) \in \mathcal{N}_{X_\xi}(x)$ — First order optimality condition

Wait-and-see and Here-and-now

Wait-and-see solution Given $F : \Xi \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ and $X_\xi \subset \mathbb{R}^n$, find $x_\xi \in X_\xi$ such that

$$-F(\xi, x_\xi) \in \mathcal{N}_{X_\xi}(x_\xi), \quad \text{i.e.,} \quad (y - x_\xi)^T F(\xi, x_\xi) \geq 0, \quad \forall y \in X_\xi.$$

Here-and-now solution One wants to make a decision x before knowing ξ . Let $X \equiv \mathbb{E}[X_\xi] = \{\mathbb{E}[x_\xi] \mid x_\xi \in X_\xi, \mathbb{E}[x_\xi] < \infty\}$.

• **Expected Residual minimization (ERM) solution**

$$\min_{x \in X} \mathbb{E}[\|r(\xi, x)\|^2],$$

where $r(\xi, \cdot)$ is a residual function.

• **Expected value (EV) solution**

$$-\mathbb{E}[F(\xi, x)] \in \mathcal{N}_X(x)$$

Stochastic complementarity problems

A random variable ξ affects $F : \Xi \times \mathbb{R}^n \rightarrow \mathbb{R}^n$,

$$x \geq 0, \quad F(\xi, x) \geq 0, \quad x^T F(\xi, x) = 0, \quad \text{for } \xi \in \Xi.$$

i.e. $\Phi(x, F(\xi, x)) = 0$, where $\Phi_i(x, F(\xi, x)) = \min(x_i, F_i(\xi, x))$, $1 \leq i \leq n$

- Expected residual minimization (ERM) formulation

Chen-Fukushima(2005)

$$\min_{x \in \mathbb{R}_+^n} \mathbb{E}[r(\xi, x)], \quad r(\xi, x) = \|\Phi(x, F(\xi, x))\|^2$$

- Expected value (EV) formulation

Gürkan-Özge-Robinson(1999), Ruszczynski-Shapiro(2003),
Jiang-Xu(2008)

$$x \geq 0, \quad \mathbb{E}[F(\xi, x)] \geq 0, \quad x^T \mathbb{E}[F(\xi, x)] = 0$$

$$\Leftrightarrow \min_{x \in \mathbb{R}^n} r(x) := \|\Phi(x, \mathbb{E}[F(\xi, x)])\|^2$$

Two-stage stochastic variational inequalities

Given the (induced) probability space $(\Xi \subset \mathbb{R}^L, \mathcal{A}, P)$, find a pair $(x \in \mathbb{R}^{n_1}, u : \Xi \rightarrow \mathbb{R}^{n_2} \text{ } \mathcal{A}\text{-measurable})$, such that the following collection of variational inequalities is satisfied:

$$\begin{aligned} -\mathbb{E}[G(\xi, x, u_\xi)] &\in \mathcal{N}_D(x) \\ -F(\xi, x, u_\xi) &\in \mathcal{N}_{C_\xi}(u_\xi) \quad \text{for a.e. } \xi \in \Xi. \end{aligned}$$

- $G : (\Xi, \mathbb{R}^{n_1}, \mathbb{R}^{n_2}) \rightarrow \mathbb{R}^{n_1}$ a vector-valued function, continuous with respect to (x, u) for all $\xi \in \Xi$, \mathcal{A} -measurable and integrable with respect to ξ .
- $\mathcal{N}_D(x)$ the normal cone to the nonempty closed-convex set $D \subset \mathbb{R}^{n_1}$ at $x \in \mathbb{R}^{n_1}$.
- $F : (\Xi, \mathbb{R}^{n_1}, \mathbb{R}^{n_2}) \rightarrow \mathbb{R}^{n_2}$ a vector-valued function, continuous with respect to (x, u) for all $\xi \in \Xi$ and \mathcal{A} -measurable with respect to ξ .
- $\mathcal{N}_{C_\xi}(v)$ the normal cone to the nonempty closed-convex set $C_\xi \subset \mathbb{R}^{n_2}$ at $v \in \mathbb{R}^{n_2}$, the random set C_ξ is \mathcal{A} -measurable.

Two-stage stochastic variational inequalities

The definition of the normal cone yields the following equivalent formulation:

$$\begin{aligned} \text{find } \bar{x} \in D \text{ and } \bar{u} : \Xi \rightarrow \mathbb{R}^{n_2}, \mathcal{A}\text{-measurable, such that } \bar{u}_\xi \in_{\text{as}} C_\xi \text{ and} \\ \langle \mathbb{E}[G(\xi, \bar{x}, \bar{u}_\xi)], x - \bar{x} \rangle \geq 0, \quad \forall x \in D, \\ \langle F(\xi, \bar{x}, \bar{u}_\xi), v - \bar{u}_\xi \rangle \geq 0, \quad \forall v \in C_\xi, \quad \text{for a.e. } \xi \in \Xi \end{aligned}$$

Two-stage stochastic linear variational inequalities (SLVI)

$$\begin{aligned} 0 &\in Ax + \mathbb{E}[B(\xi)u(\xi)] + q_1 + \mathcal{N}_D(x), \\ 0 &\in N(\xi)x + M(\xi)u(\xi) + q_2(\xi) + \mathcal{N}_{C_\xi}(u(\xi)), \quad \text{for a.e. } \xi \in \Xi. \end{aligned}$$

Two-stage stochastic linear complementarity problems (SLCP)

$$D = \mathbb{R}_+^{n_1}, C_\xi = \mathbb{R}_+^{n_2}$$

$$\begin{aligned} 0 \leq x \quad \perp \quad Ax + \mathbb{E}[B(\xi)u(\xi)] + q_1 \geq 0, \\ 0 \leq u(\xi) \quad \perp \quad N(\xi)x + M(\xi)u(\xi) + q_2(\xi) \geq 0, \quad \text{for a.e. } \xi \in \Xi. \end{aligned}$$

Two-stage SLCP

X. Chen, H. Sun and H. Xu, Discrete approximation of two-stage stochastic and distributionally robust linear complementarity problems, (2017).

Two-stage stochastic linear complementarity problems (SLCP)

$$\begin{aligned} 0 \leq x \quad \perp \quad Ax + \mathbb{E}[B(\xi)u(\xi)] + q_1 \geq 0, \\ 0 \leq u(\xi) \quad \perp \quad N(\xi)x + M(\xi)u(\xi) + q_2(\xi) \geq 0, \quad \text{for a.e. } \xi \in \Xi. \end{aligned}$$

Assumption There exists a positive continuous function $\kappa(\xi)$ such that $\mathbb{E}[\kappa(\xi)] < +\infty$ and for almost every ξ ,

$$(x^T, u^T) \begin{pmatrix} A & B(\xi) \\ N(\xi) & M(\xi) \end{pmatrix} \begin{pmatrix} x \\ u \end{pmatrix} \geq \kappa(\xi)(\|x\|^2 + \|u\|^2), \quad \forall x \in R^{n_1}, u \in R^{n_2}.$$

The two-stage SLCP has a unique solution $(x, u(\cdot)) \in R^{n_1} \times \mathcal{U}$.

\mathcal{U} is the space of measurable functions defined on Ξ .

Convergence of the sample average approximation

Two-stage stochastic generalized equations

X. Chen, A. Shapiro and H. Sun, Convergence analysis of sample average approximation of two-stage stochastic generalized equations, (2017). **without assuming relatively complete recourse**

$$\begin{aligned}0 &\in \mathbb{E}[\Phi(x, u(\xi), \xi)] + \Gamma_1(x), \quad x \in D, \\0 &\in \Psi(x, u(\xi), \xi) + \Gamma_2(u(\xi), \xi), \quad \text{for a.e. } \xi \in \Xi.\end{aligned}$$

Here $D \subset \mathbb{R}^n$ is a nonempty closed convex set,

$$\Phi : \mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \times \mathbb{R}^L \rightarrow \mathbb{R}^{n_1}, \quad \Psi : \mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \times \mathbb{R}^L \rightarrow \mathbb{R}^{n_2},$$

$\Gamma_1 : \mathbb{R}^{n_1} \rightrightarrows \mathbb{R}^{n_1}$ and $\Gamma_2 : \mathbb{R}^{n_2} \rightrightarrows \mathbb{R}^{n_2}$ are multifunctions (point-to-set mappings).

If for almost all $\xi \in \Xi$, $\Theta(x, u(\xi), \xi) := \begin{pmatrix} \Phi(x, u(\xi), \xi) \\ \Psi(x, u(\xi), \xi) \end{pmatrix}$ is strongly

monotone **at** $(x, u(\cdot))$,

then the two-stage SGE with $\Gamma_1(x) = \mathcal{N}_D(x)$ and $\Gamma_2(u(\xi), \xi) = \mathbb{R}_+^{n_2}$

has a unique solution $(x, u(\cdot)) \in \mathbb{R}^{n_1} \times \mathcal{U}$.

Algorithms for two-stage SVI

R.T. Rockafellar and J. Sun, Solving monotone stochastic variational inequalities and complementarity problems by progressive hedging, (2017). $\Xi = \{\xi^1, \dots, \xi^\nu\}$.

Extend to non-monotone stochastic VI (joint work with D. Sun and J. Yang)

Example: two-stage stochastic linear VI

$$0 \in Ax + \sum_{j=1}^{\nu} p_j B(\xi^j) u(\xi^j) + q_1 + \mathcal{N}_D(x),$$

$$0 \in N(\xi^j)x + M(\xi^j)u(\xi^j) + q_2(\xi^j) + \mathcal{N}_{C_{\xi^j}}(u(\xi^j)), \quad \text{for } j = 1, \dots, \nu,$$

where $p_j > 0$ and $\sum_{j=1}^{\nu} p_j = 1$.

Let

$C_j = C_{\xi^j}$, $B_j = B(\xi^j)$, $N_j = N(\xi^j)$, $M_j = M(\xi^j)$, $q_{2j} = q(\xi^j)$
 $D \subset \mathbb{R}^{n_1}$ and $C_j \subset \mathbb{R}^{n_2}$ are boxes. Let $\Omega = D \times C_1 \times \dots \times C_\nu$.

Example: two-stage linear SVI

$$0 \in Mz + q + \mathcal{N}_\Omega(z), \quad (1)$$

$$M = \begin{pmatrix} A & p_1 B_1 & \dots & p_\nu B_\nu \\ N_1 & M_1 & & \\ \vdots & & \ddots & \\ N_\nu & & & M_\nu \end{pmatrix}, \quad q = \begin{pmatrix} q_1 \\ q_{21} \\ \vdots \\ h_{2\nu} \end{pmatrix} \quad \text{and} \quad z = \begin{pmatrix} x \\ u_1 \\ \vdots \\ u_\nu \end{pmatrix}.$$

When $\Omega = \mathbb{R}^{n_1 + \nu n_2}$, (1) reduces to $Mz + q = 0$

When $\Omega = \mathbb{R}_+^{n_1 + \nu n_2}$, (1) reduces to $0 \leq z \perp Mz + q \geq 0$

PH Algorithm From x^k , u_j^k and w_j^k with $\sum_{j=1}^\nu p_j w_j^k = 0$, $j = 1, \dots, \nu$

Step 1 Determine \hat{x}_j^k , \hat{u}_j^k for each j by solving

$$0 \in \begin{pmatrix} A & B_j \\ N_j & M_j \end{pmatrix} \begin{pmatrix} x \\ u \end{pmatrix} + \begin{pmatrix} q_1 + w_j^k + r(x - x^k) \\ q_{2j} + r(u - u_j^k) \end{pmatrix} + \begin{pmatrix} \mathcal{N}_D(x) \\ \mathcal{N}_{C_j}(u) \end{pmatrix}$$

Step 2 Update by $x^{k+1} = \sum_{j=1}^\nu p_j \hat{x}_j^k$, $u_j^{k+1} = \hat{u}_j^k$

$$w_j^{k+1} = w_j^k + r(\hat{x}_j^k - x^{k+1}), \quad j = 1, \dots, \nu.$$

Progressive Hedging Algorithm

Give initial points $x^0 \in \mathbb{R}^{n_1}$, $u_j^0 \in \mathbb{R}^{n_2}$ and $w_j^0 \in \mathbb{R}^{n_1}$, $j = 1, \dots, \nu$ such that $\sum_{j=1}^{\nu} p_j w_j^0 = 0$. Choose $r > 0$. Let $k = 0$.

Step 1. For $j = 1, \dots, \nu$, solve the VI

$$\begin{aligned} -Ax - B_j u - q_1 - w_j^k - r(x - x^k) &\in \mathcal{N}_D(x), \\ -N_j x - M_j u - q_{2j} - r(u - u_j^k) &\in \mathcal{N}_{C_j}(u), \end{aligned}$$

and obtain a solution $(\hat{x}_j^k, \hat{u}_j^k)$, $j = 1, \dots, \nu$.

Step 2. Let $x^{k+1} = \sum_{j=1}^{\nu} p_j \hat{x}_j^k$.

$$u_j^{k+1} = \hat{u}_j^k, \quad w_j^{k+1} = w_j^k + r(\hat{x}_j^k - x^{k+1}), \quad j = 1, \dots, \nu.$$

PHA for monotone SVI is an application of Douglas-Rachford splitting method; convergence analysis for non-monotone SVI. (joint work with D. Sun and J. Yang)

Algorithms for two-stage SVI

X. Chen, T.K. Pong and R. J-B Wets, Two-stage stochastic variational inequalities: an ERM-solution procedure, Math. Program., 165(2017).

Using suitable residual functions, the two-stage stochastic VI can be formulated as the two-stage stochastic optimization problem

$$\begin{aligned} \min \quad & \theta(x) + \lambda \mathbb{E}[r(\xi, u(\xi, x)) + Q(\xi, x)] \\ \text{s.t.} \quad & x \in D \\ & u(\xi, x) = x + W y_\xi^*, \quad Q(\xi, x) = \frac{1}{2} (y_\xi^*)^T H y_\xi^*, \quad \xi \in \Xi, \quad (2) \end{aligned}$$

where

$$y_\xi^* = \operatorname{argmin}\left\{ \frac{1}{2} y_\xi^T H y_\xi \mid x + W y_\xi \in C_\xi \right\}.$$

$\lambda > 0$, $H \in \mathbb{R}^{n_2 \times n_2}$ is positive definite, $y_\xi \in \mathbb{R}^{n_2}$ is the recourse variable, $W \in \mathbb{R}^{n_1 \times n_2}$ is the recourse matrix and θ, r are residual functions.

Douglas-Rachford splitting method

Applications & Future research

1. Optimality conditions for a stochastic program
2. A Walras equilibrium problem
3. Prevailing network flow analysis (traffic, data transmission, high-speed rail, airline, power system)
4. Stochastic convex game

I. Distributionally robust two-stage variational inequalities

$$\begin{aligned} 0 &\in Ax + \mathbb{E}_P[B(\xi)u(\xi)] + q_1 + \mathcal{N}_D(x), \quad P \in \mathcal{P} \\ 0 &\in N(\xi)x + M(\xi)u(\xi) + q_2(\xi) + \mathcal{N}_{C_\xi}(u(\xi)), \quad \text{for a.e. } \xi \in \Xi. \end{aligned}$$

II. Stochastic dynamic variational inequalities

$$\begin{aligned} \dot{x}(t) &= A(t)x(t) + \mathbb{E}_{P_t}[B(t, \xi)u(t, \xi)] + q_1(t) + \mathcal{N}_D(x(t)), \quad t \in [0, T] \\ 0 &\in N(t, \xi)x(t) + M(t, \xi)u(t, \xi) + q_2(t, \xi) + \mathcal{N}_{C_\xi}(u(t, \xi)), \quad \text{for a.e. } \xi \in \Xi \end{aligned}$$

III. Multistage stochastic variational inequalities

$$x(\xi) = (x_1, x_2(\xi_1), x_3(\xi_1, \xi_2), \dots, x_N(\xi_1, \xi_2, \dots, \xi_{N-1}))$$

References

1. X. Chen and M. Fukushima, Expected residual minimization method for stochastic linear complementarity problems, *Math. Oper. Res.*, 2005.
2. H. Fang, X. Chen and M. Fukushima, Stochastic R0 matrix linear complementarity problems, *SIAM J. Optim.*, 2007.
3. X. Chen, C. Zhang and M. Fukushima, Robust solution of monotone stochastic linear complementarity problems, *Math. Program.*, 2009.
4. C. Zhang and X. Chen, Smoothing projected gradient method and its application to stochastic linear complementarity problems, *SIAM J. Optim.*, 2009.
5. C. Zhang, X. Chen and A Sumalee, Robust Wardrop's user equilibrium assignment under stochastic demand and supply: expected residual minimization approach, *Transp. Res. B*, 2011.
6. X. Chen, R. J-B Wets and Y. Zhang, Stochastic variational inequalities: residual minimization smoothing/sample average approximations, *SIAM J. Optim.*, 2012.
7. X. Chen, H. Sun and R. Wets, Regularized mathematical programs with stochastic equilibrium constraints: estimating structural demand models, *SIAM J. Optim.*, 2015.
8. X. Chen, T.K. Pong and R. J-B Wets, Two-stage stochastic variational inequalities: an ERM-solution procedure, *Math. Program* 2017.

2017

1. X. Chen and R.J-B. Wets, Stochastic Equilibrium and Variational Inequalities, a special issue of Math. Program., 165(2017).
2. X. Chen, T.K. Pong and R. J-B Wets, Two-stage stochastic variational inequalities: an ERM-solution procedure, Math. Program., 165(2017), 71-111.
3. R.T. Rockafellar and R. J.-B. Wets, Stochastic variational inequalities: single-stage to multistage, Math. Program., 165(2017), 331-360.
4. R.T. Rockafellar and J. Sun, Solving monotone stochastic variational inequalities and complementarity problems by progressive hedging, (2017), to appear in Math. Program.
5. X. Chen, H. Sun and H. Xu, Discrete approximation of two-stage stochastic and distributionally robust linear complementarity problems, (2017), revised version for Math. Program.
6. X. Chen, A. Shapiro and H. Sun, Convergence analysis of sample average approximation of two-stage stochastic generalized equations, submitted (2017).

Thank You