

Sparse Solutions of Linear Complementarity Problems

Xiaojun Chen* Shuhuang Xiang[†]

September 15, 2015

Abstract

This paper considers the characterization and computation of sparse solutions and least- p -norm ($0 < p < 1$) solutions of the linear complementarity problems $\text{LCP}(q, M)$. We show that the number of non-zero entries of any least- p -norm solution of the $\text{LCP}(q, M)$ is less than or equal to the rank of M for any arbitrary matrix M and any number $p \in (0, 1)$, and there is $\bar{p} \in (0, 1)$ such that all least- p -norm solutions for $p \in (0, \bar{p})$ are sparse solutions. Moreover, we provide conditions on M such that a sparse solution can be found by solving convex minimization. Applications to the problem of portfolio selection within the Markowitz mean-variance framework are discussed.

Keywords: Linear complementarity problem, sparse solution, nonconvex optimization, restricted isometry property.

MSC2010 Classification: 90C33, 90C26.

1 Introduction

Given an $n \times n$ matrix M and an n -dimensional vector q , the linear complementarity problem (LCP) is to find $x \in R^n$ such that

$$Mx + q \geq 0, \quad x \geq 0 \quad \text{and} \quad x^T(Mx + q) = 0.$$

We denote the problem by $\text{LCP}(q, M)$, its solution set by $\text{SOL}(q, M)$ and its feasible set by $\text{FEA}(q, M) = \{x \mid Mx + q \geq 0, x \geq 0\}$. The LCP has many applications in engineering and economics. Moreover, the LCP plays a key role in optimization theory and presents optimality conditions for constrained quadratic programs [11, 15].

The solution set $\text{SOL}(q, M)$ often has an infinite number of solutions when it is nonempty. Finding a special solution in the solution set for different goals has a long and rich history. Most readers are familiar with the least norm solution, which is defined by

$$\begin{aligned} \min \quad & \|x\|_2^2 \\ \text{s.t.} \quad & x \in \text{SOL}(q, M). \end{aligned} \tag{1.1}$$

For the monotone LCP where M is positive semi-definite, it is known that the solution set $\text{SOL}(q, M)$ is a convex polyhedra and has a unique least norm solution. Algorithms for finding

*Department of Applied Mathematics, The Hong Kong Polytechnic University, Hong Kong, China. E-mail: maxjchen@polyu.edu.hk Xiaojun Chen's work is supported partly by Hong Kong Research Grant Council grant PolyU5003/11p.

[†]Department of Applied Mathematics and Software, Central South University, Changsha 410083, Hunan, China. E-mail: xiangsh@mail.csu.edu.cn Shuhuang Xiang's work is supported partly by NSF of China (No.11371376).

the least norm solution of the monotone LCP have been studied extensively [11]. It is worth noting that some attractive interior point methods are developed to find a maximal solution that has the number of positive components in (x, s) with $s = Mx + q$ is maximal [24].

In this paper, we consider the sparsity of solutions of the LCP. We call $\bar{x} \in \text{SOL}(q, M)$ a **sparse solution** of the LCP(q, M) if \bar{x} is a solution of the following optimization problem

$$\begin{aligned} \min \quad & \|x\|_0 \\ \text{s.t.} \quad & x \in \text{SOL}(q, M), \end{aligned} \tag{1.2}$$

where $\|x\|_0$ = number of nonzero components of x .

Sparse solutions of the Z-matrix LCP(q, M) have been studied [8, 11]. A square matrix is called a Z-matrix if its off-diagonal entries are non-positive. A vector $\bar{x} \in \text{SOL}(q, M)$ is called a least element solution of the LCP(q, M), if $\bar{x} \leq x$ for all $x \in \text{SOL}(q, M)$. If M is a Z-matrix, and $\text{SOL}(q, M) \neq \emptyset$, then $\text{SOL}(q, M)$ has a unique least element solution which is the unique sparse solution of the LCP(q, M) and is the unique solution of the following linear program [8, 11]

$$\begin{aligned} \min \quad & e^T x \\ \text{s.t.} \quad & Mx + q \geq 0, \quad x \geq 0, \end{aligned} \tag{1.3}$$

where e is the vector whose all entries are one. In other words, if M is a Z-matrix, then the unique least ℓ_1 norm solution in the feasible set $\text{FEA}(q, M)$ is the unique sparse solution in the solution set $\text{SOL}(q, M)$. Moreover, if M is a positive semi-definite Z-matrix, then the least element solution is the least norm solution of the LCP(q, M). The least element solution and the least norm solution have been used in the time stepping scheme to find stable solutions of dynamic linear complementarity systems in [9, 20]. However, few theoretical results and algorithms are known for sparse solutions of the LCP(q, M) when M is not a Z-matrix.

The function $\|x\|_0$ is discontinuous and brings difficulties to analyze the models and algorithms. The ℓ_p ($0 < p < 1$) norm¹

$$\|x\|_p^p = \sum_{i=1}^n |x_i|^p$$

has been used as a continuous approximation function to $\|x\|_0$ in sparse approximation and representation. The concavity of $\|x\|_p^p$ can provide desired sparsity (see the proof of Lemma 2.1). Hence it is interesting to study the relation between the sparse solutions of (1.2) and solutions of the following optimization problem

$$\begin{aligned} \min \quad & \|x\|_p^p \\ \text{s.t.} \quad & x \in \text{SOL}(q, M). \end{aligned} \tag{1.4}$$

We call a solution of (1.4) a **least- p -norm solution**.

It is known that finding a sparse solution of a system of linear equations is NP-hard [2, 4, 5]. Recently, Ge et al showed that finding a solution of

$$\begin{aligned} \min \quad & \|x\|_p^p \\ \text{s.t.} \quad & Ax = b, \quad x \geq 0 \end{aligned} \tag{1.5}$$

for $0 < p < 1$ is also NP-hard [17], where $A \in R^{m \times n}, b \in R^m$. From [2, 4, 5, 17], we can say that finding a sparse solution and a least- p -norm solution of the LCP(q, M) is NP-hard, since we can construct M, q such that

$$S := \{x \mid Mx + q = 0, x \geq 0\} \subseteq \text{SOL}(q, M),$$

¹When $p \in [0, 1)$, $\|x\|_p$ is only a pseudo norm since it fails to satisfy the triangle inequality (and thus convexity). For simplicity, without the confusion, we call $\|x\|_p$ a norm.

and solving $\min_{x \in S} \|x\|_0$ is NP-hard by using the argument in [17].

Sparse solutions of systems of linear equations have been studied extensively in the last decades [2, 4, 5, 16, 17, 21]. For example, sparse solutions are given by the following minimization problem

$$\begin{aligned} \min \quad & \|x\|_0 \\ \text{s.t.} \quad & Ax = b. \end{aligned} \tag{1.6}$$

Candes and Tao [5] introduced the restricted isometry property (RIP) and restricted orthogonality (RO) and proved that under the RIP and RO on the coefficient matrix A , a sparse solution of (1.6) can be found by the following ℓ_1 minimization problem

$$\begin{aligned} \min \quad & \|x\|_1 \\ \text{s.t.} \quad & Ax = b. \end{aligned} \tag{1.7}$$

It is known that in general problem (1.6) is NP-hard, but problem (1.7) is a convex minimization problem and can be further recast as a linear program. The RIP and RO classify a subclass of problem (1.6) which can be solved efficiently via linear programs, and thus they attract remarkable attention in compressed sensing.

In contrast with the fast development in sparse solutions of optimization and linear equations, sparse solutions of the LCP seem to lack theory and algorithms.

The aim of this paper is to present properties of the sparse solutions and least- p -norm solutions of the LCP(q, M) and computation methods for finding the sparse solutions. In particular, we show that the number of non-zero entries of any least- p -norm solution of the LCP(q, M) is less than or equal to the rank of M for any matrix M and any number $p \in (0, 1)$, and there is $\bar{p} \in (0, 1)$ such that all least- p -norm solutions of (1.4) for $p \in (0, \bar{p})$ are sparse solutions of (1.2). Moreover, we provide conditions on M such that a sparse solution can be found by solving convex minimization.

This paper is related to the problem of finding sparse solutions to quadratic programs. Due to the optimality conditions, sparse solutions of the LCP are sparse solutions of convex constrained quadratic programs, which have important applications in portfolio optimization. The classic Markowitz portfolio optimization is formulated as the following quadratic program [22]

$$\begin{aligned} \min \quad & w^T C w \\ \text{s.t.} \quad & e^T w = 1, \quad r^T w = \rho, \\ & w \geq 0, \end{aligned} \tag{1.8}$$

where C is the covariance matrix of the return on the assets in the portfolio, w is the vector of portfolio weights that represent the amount of capital to be invested in each asset, r is the vector of expected returns of the different assets and ρ is a given total return. Sparsity is important for investors who often want to select a limited number of assets for their investment. However, finding a sparse solution of (1.8) is a challenging problem for which many approaches have been proposed such as penalty regularized optimization, mixed integer quadratic programs, quadratic programs with constraints $\|w\|_0 \leq k$ for a given integer k [3, 6]. These approaches may find good approximate sparse solutions but have some drawback. For example, a solution of a penalty regularized optimization is not necessarily a solution of the original portfolio optimization; how to choose k and how to deal with the discontinuous constraint $\|w\|_0 \leq k$ are difficult. In this paper, we show that the LCP approach has various advantages for finding a sparse solution of (1.8).

Since C is a symmetric positive semi-definite matrix, problem (1.8) is equivalent to the

LCP(q, M) with

$$M = \begin{pmatrix} C & -B^T \\ B & 0 \end{pmatrix}, \quad B = \begin{pmatrix} e^T \\ r^T \\ -e^T \\ -r^T \end{pmatrix}, \quad q = \begin{pmatrix} \mathbf{0} \\ -1 \\ -\rho \\ 1 \\ \rho \end{pmatrix}, \quad x = \begin{pmatrix} w \\ y \end{pmatrix},$$

where $y \in R^4$ is the Lagrange multiplier. Hence, if $\bar{x} = (\bar{w}, \bar{y})$ is a sparse solution of the LCP(q, M), then \bar{w} is a solution of the Markowitz mean-covariance portfolio optimization (1.8) and $\|\bar{w}\|_0 \leq \|w^*\|_0 + 4$ for any solution w^* of (1.8).

It is easy to see that $q \geq 0$ if and only if $x = 0$ is the unique least- p -norm solution and the unique sparse solution. To avoid the triviality, we assume that $x = 0$ is not a solution of the LCP(q, M). Moreover, we assume the solution set $\text{SOL}(q, M)$ is nonempty.

In section 2, we show the sparsity of solutions of (1.2) and (1.4) for an arbitrary matrix M . In section 3, we study the sparsity of solutions of (1.2) and (1.4) for a column adequate matrix M and a positive semi-definite matrix M . It is shown in [11] that a positive semi-definite matrix may not be a column adequate matrix, although a symmetric positive semi-definite matrix is a column adequate matrix. In section 4, we show that if M is column adequate and satisfies restricted orthogonality or M is positive semi-definite and $M + M^T$ satisfies restricted orthogonality [5], then a sparse solution of the LCP(q, M) can be found by using an arbitrary solution of the LCP(q, M) and solving a linear program. We also extend such approach to the column sufficient matrix LCP. In section 5, we propose a two-phase method for finding a sparse solution of the LCP(q, M) by solving quadratic programs and linear programs.

For a solution $\tilde{x} \in \text{SOL}(q, M)$, we define the following index set:

$$J = \{i : \tilde{x}_i > 0\}.$$

We define the diagonal matrix D whose diagonal elements are

$$D_{ii} = \begin{cases} 1, & i \in J \\ 0, & \text{otherwise.} \end{cases} \quad (1.9)$$

Let J^c denote the complementarity set of J and $|J|$ the number of elements of J . Let e denote the vector whose all entries are one. For $x \in R^n$, let $|x| = (|x_1|, |x_2|, \dots, |x_n|)^T$. For simplicity, without the confusion, we use $x = (w, y)$ to denote $x = (w^T, y^T)^T$.

2 Arbitrary matrix M

From [11, p.98, 144], the solution set $\text{SOL}(q, M)$ of an arbitrary LCP(q, M) is a union of a finite number of convex polyhedra. Since a convex polyhedron has only finitely many extreme points, there are only finitely many extreme points in the solution set $\text{SOL}(q, M)$. We say x is an extreme point of $\text{SOL}(q, M)$ if x does not lie in any open line segment joining two points of $\text{SOL}(q, M)$. In general, $\text{SOL}(q, M)$ is not a convex set. If x is an extreme point of $\text{SOL}(q, M)$, then x is an extreme point of a convex polyhedron.

Lemma 2.1 *All least- p -norm ($0 < p < 1$) solutions of the LCP(q, M) are extreme points of $\text{SOL}(q, M)$.*

Proof: Let \tilde{x} be a least- p -norm solution. Suppose there exist $y, z \in \text{SOL}(q, M)$ such that $\tilde{x} = \lambda y + (1 - \lambda)z$ for some $0 < \lambda < 1$. Recall that t^p is strictly concave for $t \geq 0$. Then it follows

$$\|\tilde{x}\|_p^p = \sum_{j=1}^n (\lambda y_j + (1 - \lambda)z_j)^p \geq \lambda \sum_{j=1}^n y_j^p + (1 - \lambda) \sum_{j=1}^n z_j^p = \lambda \|y\|_p^p + (1 - \lambda) \|z\|_p^p \geq \|\tilde{x}\|_p^p,$$

where the last inequality uses that \tilde{x} is a least- p -norm solution. Furthermore, the above equalities hold if and only if $y = z = \tilde{x}$, which indicates that \tilde{x} is an extreme point of $\text{SOL}(q, M)$. \blacksquare

Theorem 2.1 *Let \tilde{x} and \bar{x} be a least- p -norm solution and a sparse solution of the LCP(q, M). Then $\|\tilde{x}\|_0 \leq \text{rank}(M)$ for $p \in (0, 1)$. Moreover, there is a $\bar{p} \in (0, 1)$ such that $\|\tilde{x}\|_0 = \|\bar{x}\|_0$ for all $p \in (0, \bar{p})$.*

Proof: Note that we can choose a permutation matrix $U \in R^{n \times n}$ such that

$$UDU^T = \begin{pmatrix} I_{J,J} & 0 \\ 0 & 0 \end{pmatrix} \quad \text{and} \quad UMU^T = \begin{pmatrix} M_{J,J} & M_{J,J^c} \\ M_{J^c,J} & M_{J^c,J^c} \end{pmatrix},$$

where the matrix D is defined in (1.9). Thus

$$U(I - D + DM)U^T = \begin{pmatrix} M_{J,J} & M_{J,J^c} \\ 0 & I_{J^c,J^c} \end{pmatrix}. \quad (2.1)$$

Note that the LCP(Uq, UM) and the LCP(q, M) are equivalent in the sense that x is a solution of the LCP(q, M) if and only if Ux is a solution of the LCP(Uq, UM). Without loss of generality, we assume $U = I$ in (2.1), $J = \{1, 2, \dots, k\}$ and

$$M = \begin{pmatrix} M_{J,J} & M_{J,J^c} \\ M_{J^c,J} & M_{J^c,J^c} \end{pmatrix}, \quad \tilde{x} = \begin{pmatrix} \tilde{x}_J \\ 0 \end{pmatrix}, \quad q = \begin{pmatrix} q_J \\ q_{J^c} \end{pmatrix}.$$

Note that $\tilde{x}_J > 0$. It follows $M_{J,J}\tilde{x}_J + q_J = 0$. If $\text{rank}(M_{\cdot,J}) < |J|$, then there exists a nonzero vector $h \in R^{|J|}$ such that $M_{\cdot,J}h = 0$, i.e., $M_{J,J}h = 0$ and $M_{J^c,J}h = 0$. Furthermore, by $\tilde{x}_J > 0$, we can choose a sufficiently small real positive number δ_0 such that for all $|\delta| \leq \delta_0$,

$$\begin{aligned} \tilde{x}_J + \delta h &> 0, \\ M_{J,J}(\tilde{x}_J + \delta h) + q_J &= M_{J,J}\tilde{x}_J + q_J = 0, \\ M_{J^c,J}(\tilde{x}_J + \delta h) + q_{J^c} &= M_{J^c,J}\tilde{x}_J + q_{J^c} \geq 0. \end{aligned}$$

Hence $([\tilde{x}_J + \delta h]^T, 0)^T \in R_+^n$ is also a solution of the LCP(q, M) for $|\delta| \leq \delta_0$. Notice that \tilde{x} is a least- p -norm solution of the LCP(q, M). It leads to

$$\|\tilde{x}\|_p^p = \min_{t \in (-\delta_0, \delta_0)} \|([\tilde{x}_J + th]^T, 0)^T\|_p^p =: f(t), \quad 0 < p < 1.$$

It is impossible since

$$f''(t) = p(p-1) \sum_{i=1}^{|J|} (\tilde{x}_i + th_i)^{p-2} h_i^2 < 0, \quad -\delta_0 < t < \delta_0.$$

Hence we have $\text{rank}(M_{\cdot,J}) \geq |J|$, which implies that $\|\tilde{x}\|_0 \leq \text{rank}(M)$.

Now we prove the second part of this theorem.

We show that there is a number $\bar{p} \in (0, 1)$ such that any least- p -norm solution \tilde{x} is a sparse solution for $p \in (0, \bar{p})$.

By Lemma 2.1, all least- p -norm solutions of the LCP(q, M) are extreme points of SOL(q, M) for $p \in (0, 1)$. Let $\{x^1, x^2, \dots, x^m\}$ be the set of extreme points of SOL(q, M). Then we have for all \bar{x}

$$\|\bar{x}\|_p^p \geq \min\{\|x^1\|_p^p, \|x^2\|_p^p, \dots, \|x^m\|_p^p\} = \|\tilde{x}\|_p^p. \quad (2.2)$$

If there is not a number $\bar{p} \in (0, 1)$ such that any least- p -norm solution \tilde{x} is a sparse solution for $p \in (0, \bar{p})$, then there are a sequence $\{p_i\}, p_i > 0, p_i \rightarrow 0$ as $i \rightarrow \infty$ and a sequence $\{x^{j_i}\}$ of extreme points of SOL(q, M) such that x^{j_i} is a least- p -norm solution and

$$\|x^{j_i}\|_0 > \|\bar{x}\|_0. \quad (2.3)$$

Since there are only finitely many extreme points in SOL(q, M), without loss of generality, we assume $x^{j_i} = x^j$. However, we cannot have (2.3), since (2.2) implies

$$\|\bar{x}\|_0 = \lim_{p_i \downarrow 0} \|\bar{x}\|_{p_i}^{p_i} \geq \lim_{p_i \downarrow 0} \|x^j\|_{p_i}^{p_i} = \|x^j\|_0.$$

Hence, the second part of this theorem holds. Moreover, this together with $\|\tilde{x}\|_0 \leq \text{rank}(M)$ for $p \in (0, 1)$ implies $\|\bar{x}\|_0 \leq \text{rank}(M)$. We complete the proof. \blacksquare

From Lemma 2.1 and Theorem 2.1, we can have the following corollary.

Corollary 2.1 *There is an extreme point \bar{x} of SOL(q, M) such that \bar{x} is a sparse solution of the LCP(q, M).*

We use the following example to explain Theorem 2.1.

Example 2.1 Consider the LCP(q, M) with

$$M = \begin{pmatrix} 1 & 3 & 0 \\ 1 & 3 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad q = \begin{pmatrix} -4 \\ -4 \\ -1 \end{pmatrix}.$$

The solution set is SOL(q, M) = $S_1 \cup S_2$ where

$$S_1 = \{(x_1, x_2, 0)^T : x_1 + 3x_2 = 4, x_1 > 1, x_2 \geq 0\}, \quad S_2 = \{(1, 1, x_3)^T : x_3 \geq 0\}.$$

The sparse solution is $\bar{x} = (4, 0, 0)^T$.

The least- p -norm solutions are $\tilde{x} = (1, 1, 0)^T$ for $p > \frac{1}{2}$; $\tilde{x} = (1, 1, 0)^T$ or $\tilde{x} = (4, 0, 0)^T$ for $p = \frac{1}{2}$; and $\tilde{x} = (4, 0, 0)^T$ for $0 < p < \frac{1}{2}$.

The number of non-zero components in the sparse solution and all least- p -norm solutions is one or two, which is less than or equal to $\text{rank}(M)=2$. Moreover, $\|\tilde{x}\|_0 = \|\bar{x}\|_0$ for all least- p -norm solutions with $p \in (0, \frac{1}{2})$.

Let us consider other LCP(q, M) with

$$M = \begin{pmatrix} 1 & 3 & 0 & 0 & 1 \\ 1 & 3 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 1 \\ 3 & 3 & 0 & 0 & 1 \end{pmatrix}, \quad q = \begin{pmatrix} -4 \\ -4 \\ -1 \\ -1 \\ -6 \end{pmatrix}.$$

The solution set is $\text{SOL}(q, M) = S_1 \cup S_2 \cup S_3$ where

$$S_1 = \left\{ (1, 1, x_3, x_4, 0)^T : x_3 \geq 0, x_4 \geq 0 \right\}, \quad S_2 = \left\{ (1, x_2, 0, 0, 3 - 3x_2)^T : 0 \leq x_2 \leq 1 \right\}$$

$$S_3 = \left\{ (0, 0, x_3, x_4, 6)^T : x_3 \geq 0, x_4 \geq 0 \right\}.$$

The sparse solution is $\bar{x} = (0, 0, 0, 0, 6)^T$.

Consider a threshold \bar{p} for $2 = 6^{\bar{p}}$. We find that the least- p -norm solutions are $\tilde{x} = (1, 1, 0, 0, 0)^T$ for $p > \frac{1}{\log_2 6}$; $\tilde{x} = (1, 1, 0, 0, 0)^T$ or $\tilde{x} = (0, 0, 0, 0, 6)^T$ for $p = \frac{1}{\log_2 6}$; and $\tilde{x} = (0, 0, 0, 0, 6)^T$ for $0 < p < \frac{1}{\log_2 6}$.

The number of non-zero components in the sparse solution and all least- p -norm solutions is less than $\text{rank}(M) = 3$. Moreover, $\|\tilde{x}\|_0 = \|\bar{x}\|_0$ for all least- p -norm solutions with $p \in (0, \frac{1}{\log_2 6})$.

Remark 2.1 *From the proof of Theorem 2.1, we can see that if $x^* \in \text{SOL}(q, M)$ and the column $M_{\cdot, j}$ is linearly dependent, then $([x_j^* + \delta h]^T, 0)^T \in \text{SOL}(q, M)$ for all sufficiently small $|\delta|$, which implies that $x^* = \frac{1}{2}(x^* + \delta h) + \frac{1}{2}(x^* - \delta h) \in \text{SOL}(q, M)$. Hence if $x^* \in \text{SOL}(q, M)$ is an extreme point, then the column $M_{\cdot, j}$ is linearly independent, and thus $\|x^*\|_0 \leq \text{rank}(M)$.*

Remark 2.2 *The Lemke algorithm is a popular algorithm for solving the LCP, which is a simplex-like vertex following algorithm [11]. Various schemes of the Lemke algorithm have stimulated remarkable research on the class of matrices M for which they can process the LCP(q, M). In [1], Adler and Verma present a sufficient condition for the processability of the Lemke algorithm for the E_0 -matrix LCP which unifies several sufficient conditions for a number of well known subclasses of the E_0 -matrix LCPs. A matrix M is called an E_0 -matrix if for any non-zero $x \geq 0$, there exists an index k such that $x_k > 0$ and $(Mx)_k \geq 0$. A matrix M is called a P_0 -matrix if for any non-zero x , there exists an index k such that $x_k \neq 0$ and $x_k(Mx)_k \geq 0$ [11]. By the definition, if M is a P_0 -matrix, then M is an E_0 -matrix. If M is an E_0 -matrix and the solution set $\text{SOL}(q, M) \neq \emptyset$, then the Lemke algorithm can find an extreme point x^* of $\text{SOL}(q, M)$ in a finitely many steps. By Remark 2.1, $\|x^*\|_0$ gives a sharper bound for the sparsity of the solution than $\text{rank}(M)$, that is, for any least- p -norm solution \tilde{x} for sufficiently small p and any sparse solution \bar{x} , we have $\max\{\|\tilde{x}\|_0, \|\bar{x}\|_0\} \leq \|x^*\|_0$.*

3 Column adequate matrix M

A matrix $M \in R^{n \times n}$ is called column adequate if $z_i(Mz)_i \leq 0$ for all $i = 1, 2, \dots, n$ implies $Mz = 0$ [11, 19]. The LCP(q, M) is said to be w -unique if there is a unique vector w such that $Mx + q = w$ for any $x \in \text{SOL}(q, M)$. Ingleton [19] proved that the LCP(q, M) is w -unique for any $q \in R^n$ if and only if M is column adequate.

Lemma 3.1 ([11, Theorem 3.4.4] and [23, Theorem 2.7]) *The following statements are equivalent.*

- (i) M is column adequate.
- (ii) M is P_0 -matrix and for each index set α for which $\det M_{\alpha, \alpha} = 0$, the columns of $M_{\cdot, \alpha}$ are linearly dependent.
- (iii) For all $q \in \text{FEA}(q, M)$, if z^1 and z^2 are any two solutions of the LCP(q, M), then $Mz^1 = Mz^2$.

From Theorem 3.1.7 [11], we see that if M is symmetric positive semi-definite, then M is adequate.

Theorem 3.1 Suppose that M is column adequate. Let \bar{x} be a sparse solution of the LCP(q, M). With the index set J and diagonal matrix D , the following statements hold.

(i) $M_{J,J}$ is nonsingular;

(ii) $\bar{x} = -(I - D + DM)^{-1}Dq$;

(iii) $\|\bar{x}\|_1 \leq L\|q\|_1$,

where $L = \max \left\{ \|M_{\alpha,\alpha}^{-1}\|_1 : M_{\alpha,\alpha} \text{ is nonsingular for } \alpha \subseteq \{1, \dots, n\} \right\}$;

(iv) there is no another solution $x \in \text{SOL}(q, M)$ with $\alpha = \{i : x_i > 0\}$ such that $\alpha \subseteq J$.

The statements also hold for any least- p -norm solution of the LCP(q, M).

Proof: (i) Following the proof of Theorem 2.1, without loss of generality, we assume $U = I$ in (2.1), $J = \{1, 2, \dots, k\}$ and

$$M = \begin{pmatrix} M_{J,J} & M_{J,J^c} \\ M_{J^c,J} & M_{J^c,J^c} \end{pmatrix}, \quad \bar{x} = \begin{pmatrix} \bar{x}_J \\ 0 \end{pmatrix}, \quad q = \begin{pmatrix} q_J \\ q_{J^c} \end{pmatrix}.$$

Note that $\bar{x}_J > 0$. It follows $M_{J,J}\bar{x}_J + q_J = 0$. If $M_{J,J}$ is singular, then from Lemma 3.1, the columns of $M_{\cdot,J}$ are linearly dependent, and there exists a nonzero vector $h \in R^{|J|}$ such that $M_{\cdot,J}h = 0$. Define

$$\tau = \min_{h_i \neq 0, 1 \leq i \leq |J|} \frac{\bar{x}_i}{|h_i|}.$$

It is easy to verify that

$$\bar{x}_J \pm \tau h \geq \bar{x}_J - \tau|h| \geq 0, \quad M_{J,J}(\bar{x}_J \pm \tau h) + q_J = 0, \quad M_{J^c,J}(\bar{x}_J \pm \tau h) + q_{J^c} \geq 0.$$

Hence $([\bar{x}_J \pm \tau h]^T, 0)^T \in R_+^n$ is also a solution of the LCP(q, M) but $\|\bar{x}_J - \tau h\|_0 < \|\bar{x}\|_0$ or $\|\bar{x}_J + \tau h\|_0 < \|\bar{x}\|_0$. Hence, these together imply that $M_{J,J}$ is nonsingular.

(ii) From the nonsingularity of $M_{J,J}$, expression (2.1) with $U = I$ implies that $I - D + DM$ is nonsingular and

$$(I - D + DM)^{-1}D = \begin{pmatrix} M_{J,J} & M_{J,J^c} \\ 0 & I \end{pmatrix}^{-1} D = \begin{pmatrix} M_{J,J}^{-1} & 0 \\ 0 & 0 \end{pmatrix}. \quad (3.1)$$

From $(I - D)\bar{x} + D(M\bar{x} + q) = 0$ and (3.1), we obtain the desired result.

(iii) From (3.1), we have

$$\|(I - D + DM)^{-1}D\|_1 \leq \max \left\{ \|M_{\alpha,\alpha}^{-1}\|_1 : M_{\alpha,\alpha} \text{ is nonsingular for } \alpha \subseteq \{1, \dots, n\} \right\},$$

which together with (ii) implies (iii).

(iv) Assume that there is another solution $\hat{x} \in \text{SOL}(q, M)$ with $\alpha = \{i : \hat{x}_i > 0\}$ such that $\alpha \subseteq J$. From the proof of (i), without loss of generality, assume

$$M = \begin{pmatrix} M_{\alpha,\alpha} & M_{\alpha,\beta} & M_{\alpha,J^c} \\ M_{\alpha,\beta} & M_{\beta,\beta} & M_{\beta,J^c} \\ M_{J^c,\alpha} & M_{J^c,\beta} & M_{J^c,J^c} \end{pmatrix}, \quad q = \begin{pmatrix} q_\alpha \\ q_\beta \\ q_{J^c} \end{pmatrix}, \quad J = \alpha \cup \beta.$$

It is easy to verify that both \hat{x}_J and \bar{x}_J are solutions of the LCP($q_J, M_{J,J}$). However $M_{J,J}$ is nonsingular and is a P-matrix, then the LCP($q_J, M_{J,J}$) has a unique solution. This is a contradiction.

Using the same argument and the proof of Theorem 2.1, we can see the same statements hold for any least- p -norm solution \tilde{x} of SOL(q, M). ■

Corollary 3.1 *Suppose that M is column adequate. Then all sparse solutions of the LCP(q, M) are extreme points of SOL(q, M).*

Proof: Let \bar{x} be a sparse solution. Assume that there exist $y, z \in \text{SOL}(q, M)$ such that $\bar{x} = \lambda y + (1 - \lambda)z$ for some $0 < \lambda < 1$. Since \bar{x} is a sparse solution, this means \bar{x}, y, z have the same support sets. However, from (iv) of Theorem 3.1 the support sets of \bar{x}, y, z are same if and only if $\bar{x} = y = z$. This is a contradiction. Hence \bar{x} is an extreme point of SOL(q, M). ■

Corollary 3.2 *Suppose that M is column adequate and $x^* \in \text{SOL}(q, M)$. Then x^* is an extreme point if and only if $\det M_{J(x^*), J(x^*)} \neq 0$.*

Proof: The necessity is from Remark 2.1 together with (ii) of Lemma 3.1. For the sufficiency, suppose there exist $y, z \in \text{SOL}(q, M)$ such that $x^* = \lambda y + (1 - \lambda)z$ for some $0 < \lambda < 1$. Then $J(y), J(z) \subseteq J(x^*)$. From the proof of (iv) of Theorem 3.1, it is easy to verify that $x_{J(x^*)}^*, y_{J(x^*)}$ and $z_{J(x^*)}$ are the solutions of LCP($q_{J(x^*)}, M_{J(x^*), J(x^*)}$). However, LCP($q_{J(x^*)}, M_{J(x^*), J(x^*)}$) has a unique solution since $M_{J(x^*), J(x^*)}$ is a P-matrix. This is a contradiction. Hence x^* is an extreme point. ■

From Theorem 2.1 and Theorem 3.1, if M is column adequate, the number of non-zero components of any sparse solution and least- p -norm solution of the LCP(q, M) is less than or equal to $\max \{\text{rank}(M_{\alpha, \alpha}) : \alpha \subseteq \{1, 2, \dots, n\}\}$. We use the following example to explain the sparsity.

Example 3.1 Consider the LCP(q, M) with

$$M = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}, \quad q = \begin{pmatrix} -1 \\ -1 \end{pmatrix}.$$

It is easy to see $\max \{\text{rank}(M_{\alpha, \alpha}) : \alpha \subseteq \{1, 2\}\} = 1$.

The solution set is $\text{SOL}(q, M) = \{(x_1, x_2)^T : x_1 + x_2 = 1, x_1, x_2 \geq 0\}$.

The sparse solution is $\bar{x} = \{(1, 0)^T, (0, 1)^T\}$.

The least- p -norm solution is $\tilde{x} = \{(1, 0)^T, (0, 1)^T\}$ for $0 < p < 1$.

For $p = 1$, each solution in SOL(q, M) is the least ℓ_1 norm solution. For $p > 1$, $(\frac{1}{2}, \frac{1}{2})^T$ is the least ℓ_p norm solution.

Let us consider other LCP(q, M) with

$$M = \begin{pmatrix} 5 & -1 & 1 \\ -1 & 1 & 1 \\ 1 & 1 & 2 \end{pmatrix}, \quad q = \begin{pmatrix} -4 \\ 0 \\ -2 \end{pmatrix},$$

and $\max \{\text{rank}(M_{\alpha, \alpha}) : \alpha \subseteq \{1, 2, 3\}\} = 2$.

The solution set is $\text{SOL}(q, M) = \{(x_1, x_2, x_3)^T : x_1 = \lambda + (1 - \lambda)\frac{2}{3}, x_2 = \lambda, x_3 = (1 - \lambda)\frac{2}{3}, 0 \leq \lambda \leq 1\}$.

The sparse solution is $\bar{x} = \{(\frac{2}{3}, 0, \frac{2}{3})^T, (1, 1, 0)^T\}$.

The least- p -norm solution is $\tilde{x} = \{(\frac{2}{3}, 0, \frac{2}{3})^T\}$ for $0 < p < 1$, which is also the least ℓ_1 norm solution and the least ℓ_p norm solution for $p \geq 1$.

It is worth noting that the least p -norm solution is unable to characterize all the sparse solutions. This is not surprising, since $\|x\|_p^p$ is also concerned with the value of each component $|x_i|$ unlike $\|x\|_0$.

Remark 3.1 *The sparsity of solutions of the LCP(q, M) is sensitive with the data (q, M). Consider the following LCP(q, M) with a symmetric positive semi-definite matrix*

$$M = \begin{pmatrix} 2 & 1 & 3 \\ 1 & 1 & 0 \\ 3 & 0 & 9 \end{pmatrix}, \quad q = \begin{pmatrix} -2 \\ -1 \\ -3 \end{pmatrix}.$$

The sparse solution is $\bar{x} = (1, 0, 0)^T$ with $\|\bar{x}\|_0 = 1$. However, for $q + \varepsilon e$ with $0 < \varepsilon < \frac{3}{2}$, the sparse solution is $\bar{x} = (1 - 2\varepsilon/3, 0, \varepsilon/9)^T$ with $\|\bar{x}\|_0 = 2$. It is known that the solvability of the monotone LCP(q, M) is stable for nonnegative noise in q , since the feasibility implies the solvability for the monotone LCP(q, M). However, the sparsity of solutions of the monotone LCP(q, M) can change with any small positive noise in q .

4 Computation of sparse solutions

In this section, we show that we can find a sparse solution of the LCP(q, M) by using an arbitrary solution of the LCP(q, M) and solving a linear program if the matrix M is column adequate and satisfies the s -restricted isometry property (RIP) for $1 \leq s \leq n$ and s, s' -restricted orthogonality (RO) for $s + s' \leq n$ [5] or M is positive semi-definite and $M + M^T$ satisfies s -RIP for $1 \leq s \leq n$ and s, s' -RO for $s + s' \leq n$.

An $m \times n$ matrix A is said to satisfy the s -RIP with a restricted isometry constant δ_s if for every $m \times |\Lambda|$ submatrix A_Λ of A and for every vector $z \in R^{|\Lambda|}$ with $|\Lambda| \leq s$,

$$(1 - \delta_s)\|z\|_2^2 \leq \|A_\Lambda z\|_2^2 \leq (1 + \delta_s)\|z\|_2^2. \quad (4.1)$$

Moreover, A is said to satisfy the s, s' -RO with a restricted orthogonality constant $\theta_{s, s'}$ for $s + s' \leq n$ if for all submatrices $A_\Lambda \in R^{m \times |\Lambda|}$, $A_{\Lambda'} \in R^{m \times |\Lambda'|}$ of A with $|\Lambda| \leq s$, $|\Lambda'| \leq s'$ and for all vectors $z \in R^{|\Lambda|}$, $z' \in R^{|\Lambda'|}$

$$|(A_\Lambda z, A_{\Lambda'} z')| \leq \theta_{s, s'} \|z\|_2 \|z'\|_2 \quad (4.2)$$

holds for all disjoint sets Λ and Λ' .

The concepts of s -RIP and s, s' -RO were introduced by Candes and Tao [5] and are used in many applications of sparse representations.

Using $x^T M x = \frac{1}{2} x^T (M + M^T) x$, the LCP(q, M) can be equivalently written as a quadratic program

$$\begin{aligned} \min \quad & \frac{1}{2} x^T (M + M^T) x + q^T x \\ \text{s.t.} \quad & M x + q \geq 0, x \geq 0 \end{aligned} \quad (4.3)$$

in the sense that x^* is a solution of the LCP(q, M) if and only if x^* is an optimal solution of (4.3) with the optimal value of zero. If M is a positive semi-definite matrix (need not be symmetric), then (4.3) is a convex quadratic program.

From Theorem 3.1.7 in [11], the solution set SOL(q, M) for a positive semi-definite matrix M equals to

$$\text{SOL}(q, M) = \left\{ x \in R_+^n \mid M x + q \geq 0, (M + M^T) x = c, q^T x = \gamma \right\}, \quad (4.4)$$

where $c = (M + M^T) x^*$, $\gamma = q^T x^*$ and x^* is an arbitrary solution of the LCP(q, M).

We consider the following linear program

$$\begin{aligned} \min \quad & e^T x \\ \text{s.t.} \quad & M x + q \geq 0, x \geq 0, (M + M^T) x = c, q^T x = \gamma. \end{aligned} \quad (4.5)$$

Theorem 4.1 Suppose that M is positive semi-definite. Let \hat{x} be a solution of the linear program (4.5) with $\|\hat{x}\|_0 \leq s$.

- (i) If $(M + M^T)$ satisfies the RIP with a restricted isometry constant $\delta_{2s} < 1$, then \hat{x} is the unique sparse solution of the LCP(q, M).
- (ii) If $(M + M^T)$ satisfies the RIP and RO with

$$\delta_s + \theta_{s,s'} + \theta_{s,2s'} < 1, \quad (4.6)$$

then \hat{x} is the unique solution of the linear program (4.5) and the unique sparse solution of the LCP(q, M).

Proof: (i) From (4.4), we know that \hat{x} is a solution of the LCP(q, M). Assume on contradiction that there is a sparse solution of the LCP(q, M) such that $\bar{x} \neq \hat{x}$. Then $\|\bar{x}\|_0 \leq \|\hat{x}\|_0 \leq s$ and $(M + M^T)(\hat{x} - \bar{x}) = 0$. Let the support set of $\hat{x} - \bar{x}$ be K . Then $|K| \leq 2s$. Hence $\|\hat{x} - \bar{x}\|_0 \leq 2s$, which together with the RIP, yields

$$\begin{aligned} (1 - \delta_{2s})\|\hat{x} - \bar{x}\|_2^2 &= (1 - \delta_{2s})\|(\hat{x} - \bar{x})_K\|_2^2 \\ &\leq \|(M + M^T)_{\cdot, K}(\hat{x} - \bar{x})_K\|_2^2 = \|(M + M^T)(\hat{x} - \bar{x})\|_2^2 = 0. \end{aligned}$$

This is a contradiction to $\hat{x} \neq \bar{x}$. Therefore \hat{x} is the unique sparse solution of the LCP(q, M).

- (ii) From Theorem 1.3 in [5], \hat{x} is the unique solution of the following linear program

$$\begin{aligned} \min \quad & \|x\|_1 \\ \text{s.t.} \quad & (M + M^T)x = c. \end{aligned} \quad (4.7)$$

Since the convex feasible set of (4.5) is contained in the convex set $\{x \mid (M + M^T)x = c\}$, \hat{x} is also the unique solution of the linear program (4.5).

From Lemma 1.1 in [5], the condition in (4.6) implies $\delta_{2s} < 1$. Hence, from (i) of this theorem, \bar{x} is the unique sparse solution of the LCP(q, M). ■

Theorem 4.2 Suppose that M is column adequate. Let \hat{x} with $\|\hat{x}\|_0 \leq s$ be a solution of the linear program

$$\begin{aligned} \min \quad & e^T x \\ \text{s.t.} \quad & x \geq 0, Mx = c, q^T x = \gamma, \end{aligned} \quad (4.8)$$

where $c = Mx^*$, $\gamma = q^T x^*$ and x^* is an arbitrary solution of the LCP(q, M).

- (i) If M satisfies the RIP with a restricted isometry constant $\delta_{2s} < 1$, then \hat{x} is the unique sparse solution of the LCP(q, M).
- (ii) If M satisfies the RIP and RO with (4.6), then \hat{x} is the unique solution of the linear program (4.8) and the unique sparse solution of the LCP(q, M).

Proof: From Lemma 3.1, the LCP(q, M) is w -unique for a column adequate matrix M . Hence the solution set SOL(q, M) for a column adequate matrix M is convex and equals to

$$\text{SOL}(q, M) = \left\{ x \in R_+^n \mid Mx = c, q^T x = \gamma \right\}. \quad (4.9)$$

Following the proof of Theorem 4.1, we can obtain the desired results. ■

A matrix $M \in R^{n \times n}$ is called column sufficient [11, 12] if

$$x_i(Mx)_i \leq 0 \quad \text{for } i = 1, 2, \dots, n \quad \Rightarrow \quad x_i(Mx)_i = 0 \quad \text{for } i = 1, 2, \dots, n. \quad (4.10)$$

Obviously, if M is column adequate or M is positive semi-definite, then M is column sufficient. In [12], Cottle et al show that the solution set $\text{SOL}(q, M)$ is convex for every $q \in R^n$ if and only if M is column sufficient. Moreover, there exist complementary index set α and α^c such that the solution set $\text{SOL}(q, M)$ for a column sufficient matrix equals to

$$\text{SOL}(q, M) = \left\{ x \in R_+^n \mid Mx + q \geq 0, \check{M}x + \check{q} = 0 \right\}, \quad (4.11)$$

where

$$\check{M} = \begin{pmatrix} M_{\alpha, \alpha} & 0 \\ 0 & I_{\alpha^c, \alpha^c} \end{pmatrix} \quad \check{q} = \begin{pmatrix} q_\alpha \\ 0 \end{pmatrix},$$

and $\alpha = \{i : (Mx^* + q)_i = 0\}$ for some solution x^* of the $\text{LCP}(q, M)$.

Following the proof of Theorem 4.1, we can derive the following corollary.

Corollary 4.1 *Suppose that M is column sufficient. Let \hat{x} with $\|\hat{x}\|_0 \leq s$ be a solution of the linear program*

$$\begin{aligned} \min \quad & e^T x \\ \text{s.t.} \quad & x \geq 0, Mx + q \geq 0, \check{M}x + \check{q} = 0. \end{aligned} \quad (4.12)$$

(i) *If \check{M} satisfies the RIP with a restricted isometry constant $\delta_{2s} < 1$, then \hat{x} is the unique sparse solution of the $\text{LCP}(q, M)$.*

(ii) *If \check{M} satisfies the RIP and RO with (4.6), then \hat{x} is the unique solution of the linear program (4.8) and the unique sparse solution of the $\text{LCP}(q, M)$.*

Example 4.1 *Consider the $\text{LCP}(q, M)$ with*

$$M = \begin{pmatrix} 0.4 & -0.3 & 0.1 \\ -0.3 & 0.3 & -0.3 \\ 0.1 & -0.3 & 0.7 \end{pmatrix}, \quad q = \begin{pmatrix} -0.4 \\ 0.3 \\ -0.1 \end{pmatrix}.$$

The solution set is $\text{SOL}(q, M) = \{(1, 0, 0)^T + \lambda(2, 3, 1)^T : \lambda \geq 0\}$.

The restricted isometry constants are $\delta_1 = 0.4901$ and $\delta_2 = 0.8421$. From (i) of Theorem 4.2, $\bar{x} = (1, 0, 0)^T$ is the unique sparse solution of the $\text{LCP}(q, M)$.

Remark 4.1 *For an $m \times n$ matrix A , the concept of $\text{Spark}(A)$ is also often used in the study of sparse solutions, which is defined as the smallest possible number such that there exists a subgroup of columns from A that are linearly dependent [13].*

Suppose that M is positive semi-definite. Let \hat{x} be a solution of the $\text{LCP}(q, M)$ with $\|\hat{x}\|_0 \leq \frac{1}{2}\text{Spark}(M + M^T)$. Then \hat{x} is a sparse solution of the $\text{LCP}(q, M)$. This statement can be shown as follows.

Suppose x' is another solution of the $\text{LCP}(q, M)$, then from (4.4), $(M + M^T)(\hat{x} - x') = 0$, which implies $\|\hat{x} - x'\|_0 \geq \text{Spark}(M + M^T)$ and

$$\|x'\|_0 \geq \text{Spark}(M + M^T) - \|\hat{x}\|_0 \geq \frac{1}{2}\text{Spark}(M + M^T) \geq \|\hat{x}\|_0.$$

Similarly, if $\|\hat{x}\|_0 < \frac{1}{2}\text{Spark}(M + M^T)$, then \hat{x} is the unique sparse solution of the $\text{LCP}(q, M)$.

For a column adequate matrix M , if \hat{x} is a solution of the $\text{LCP}(q, M)$ with $\|\hat{x}\|_0 \leq \frac{1}{2}\text{Spark}(M)$ then \hat{x} is a sparse solution of the $\text{LCP}(q, M)$. Moreover, the strict inequality implies the uniqueness.

Example 4.2 Consider the LCP(q, M) with

$$M = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}, \quad q = \begin{pmatrix} -1 \\ -1 \\ -1 \end{pmatrix}.$$

The solution set: $SOL(q, M) = \{(x_1, x_2, x_3)^T : x_1 + x_2 + x_3 = 1, x_1 \geq 0, x_2 \geq 0, x_3 \geq 0\}$.

$\text{Spark}(M) = 2$, and $(1, 0, 0)^T, (0, 1, 0)^T, (0, 0, 1)^T$ are sparse solutions of the LCP(q, M).

5 Numerical experiments

From Theorems 4.1-4.2 and Corollary 4.1, we propose a two phase method to find a sparse solution of the LCP(q, M), i.e. a solution of (1.2) as follows.

Phase 1: Find a solution x^* of the LCP(q, M).

Phase 2: Solve a linear program (4.12) if M is column sufficient, or a linear program (4.8) if M is column adequate, or a linear program (4.5) if M is positive semi-definite.

To test the two phase method, we apply Theorem 4.1 to sparse solutions of the following quadratic program

$$\begin{aligned} \min \quad & \frac{1}{2}z^T H z + c^T z \\ \text{s.t.} \quad & A z \geq b \\ & z \geq 0 \end{aligned} \tag{5.1}$$

where $H \in R^{m \times m}$ is positive semi-definite, $c \in R^m, A \in R^{k \times m}, b \in R^k$. This quadratic program includes the Markowitz mean-covariance portfolio optimization problem (1.8) as a special case. Let S_{QP} be the solution set of (5.1). We say \bar{z} is a sparse solution of the quadratic program (5.1) if

$$\|\bar{z}\|_0 = \min\{\|z\|_0 : z \in S_{QP}\}.$$

The quadratic program (5.1) is equivalent to the LCP(q, M) with

$$M = \begin{pmatrix} H & -A^T \\ A & 0 \end{pmatrix}, \quad q = \begin{pmatrix} c \\ -b \end{pmatrix}, \quad x = \begin{pmatrix} z \\ y \end{pmatrix},$$

where $y \in R^k$ is the Lagrange multiplier. Note that $M + M^T = \begin{pmatrix} H & 0 \\ 0 & 0 \end{pmatrix}$. The solution set $SOL(q, M)$ equals to

$$SOL(q, M) = \{x \in R_+^n \mid Mx + q \geq 0, Hz = w, q^T x = \gamma\},$$

where $w = Hz^*, \gamma = q^T x^*$ and $x^* = (z^*, y^*)$ is an arbitrary solution of the LCP(q, M).

We consider the following linear program

$$\begin{aligned} \min \quad & e^T z \\ \text{s.t.} \quad & Mx + q \geq 0, x \geq 0, Hz = w, q^T x = \gamma. \end{aligned} \tag{5.2}$$

Let $\hat{x} = (\hat{z}, \hat{y})$ be a solution of the linear program (5.2) with $\|\hat{z}\|_0 \leq s$. According to Theorem 4.1, we have the following statements.

- (i) If H satisfies the RIP with a restricted isometry constant $\delta_{2s} < 1$, then \hat{z} is the unique sparse solution of the quadratic program (5.1).
- (ii) If H satisfies the RIP and RO with (4.6) then \hat{z} is the unique sparse solution of the quadratic program (5.1) and all solutions $x^* = (z^*, y^*)$ of the linear program (5.2) have the same component $z^* = \hat{z}$.

Based on the statements above and (4.3), we propose the following procedure to find a sparse solution of (5.1).

1. Find a solution x^* of the LCP(q, M) by solving the following quadratic program

$$\begin{aligned} \min \quad & z^T H z + c^T z - b^T y \\ \text{s.t.} \quad & A z \geq b, H z - A^T y \geq -c, z, y \geq 0. \end{aligned} \quad (5.3)$$

2. Find a solution of the linear program (5.2).

We use the following code in Matlab to generate a solution $z \in R^m$ with $\|z\|_0 = s$ of (5.1), a positive semi-definite matrix H , a matrix $A \in R^{k \times m}$, and vectors $c \in R^m, b \in R^k$.

```
k=fix(m/5); s=fix(m/3); z=zeros(m,1); P=randperm(m);
z(P(1:2*s+m/10))=abs(randn(2*s+m/10,1)); H=randn(m,m); H=H*diag(z)*H';
A=randn(k-1,m); A=[A;-ones(1,m)]; z=zeros(m,1);
z(P(1:s))=abs(randn(s,1)); b=A*z; c=-H*z.
```

For each m, k, s , we generated 100 independent test problems by the code. The convex quadratic program (5.3) and the linear program (5.2) are solved by the Matlab code `quadprog` and `linprog` with initial iterate $x_0 = \text{zeros}(n, 1)$. Preliminary numerical results are reported in Table 1. In the last line of Table 1, we report $\|z_{LP}\|_0, n_1; n_2$ where z_{LP} is the numerical solution of the linear program (5.2), n_1 is the average of $\|z_{LP}\|_0$ for the 100 test problems and n_2 is the number of test problems with $\|z_{LP}\| \leq s$.

Table 1: 100 independent tests for each (m, k, s)

m	80	90	100	110	120	130	140	150
k	16	18	20	22	24	26	28	30
$\text{rank}(H)$	60	69	76	83	92	99	106	115
s	26	30	33	36	40	43	46	50
$\ z_{LP}\ _0$	25.9; 100	30; 100	33; 99	35.9; 100	40.2; 98	42.9; 100	46.2; 99	50.3; 99

The numerical testing is performed using MATLAB R2011b on a Lenovo PC (Intel Quad CPU Q9550, 2.83GHz, 4.00GB of RAM). The numerical results are encouraging for the study of sparse solutions of the LCP, although the matrix H generated by the Matlab code may not satisfy the RIP and RO conditions.

Acknowledgments. The authors are grateful for the comments of two very knowledgeable and thorough referees.

References

- [1] I. Adler and S. Verma, The linear complementarity problem, Lemke algorithm, perturbation, and the complexity class PPA, Industrial Engineering & Operations Research, University of California Berkeley 2011.
- [2] A.M. Bruckstein, D.L. Donoho and M. Elad, From sparse solutions of systems of equations to sparse modeling of signals and images, SIAM Rev., 51(2009), 34-81.
- [3] J. Brodie, I. Daubechies, C. De Mol, D. Giannone and I. Loris, Sparse and stable Markowitz portfolios, Proc. Natl. Acad. Sci., 106(2009), 12267-12272.
- [4] E.J. Candes, J. Romberg and T. Tao, Robust uncertainty principles: exact signal reconstruction from highly incomplete frequency information, IEEE T. Inform. Theory, 52(2006) 489-509.
- [5] E. Candes and T. Tao, Decoding by linear programming, IEEE T. Inform. Theory, 51(2005) 4203-4215.
- [6] F. Cesarone, A. Scozzari and F. Tardella, Efficient algorithms for mean-variance portfolio optimization with hard real-world constraints, Giornale dell'Istituto Italiano degli Attuari, 72(2009), 37-56.
- [7] X. Chen, D. Ge, Z. Wang and Y. Ye, Complexity of unconstrained L_2 - L_p minimization, Math. Program., 143(2014), 371-383.
- [8] X. Chen and S. Xiang, Implicit solution function of P_0 and Z matrix linear complementarity constraints, Math. Program., 128(2011), 1-18.
- [9] X. Chen and S. Xiang, Newton iterations in implicit time-stepping scheme for differential linear complementarity systems, Math. Program., 138(2013), 579-606.
- [10] X. Chen, F. Xu and Y. Ye, Lower bound theory of nonzero entries in solutions of l_2 - l_p minimization, SIAM J. Sci. Comput., 32(2010), 2832-2852.
- [11] R.W. Cottle, J.-S. Pang and R.E. Stone, The Linear Complementarity Problem, Academic Press, Boston, MA, 1992.
- [12] R.W. Cottle, J.-S. Pang and V. Venkateswaran, Sufficient matrices and the linear complementarity problem, Linear Algebra Appl. 114/115(1989), 231-249.
- [13] D.L. Donoho and M. Elad, Optimally sparse representation in general (non-orthogonal) dictionaries via L_1 minimization, Proc. Natl. Acad. Sci., 100(2003), 2197-2202.
- [14] J. Fan and R. Li, Variable selection via nonconcave penalized likelihood and its oracle properties, J. Amer. Statist. Assoc., 96(2001), 1348-1360.
- [15] M.C. Ferris and J.-S. Pang, Engineering and economic applications of complementarity problems, SIAM Rev., 39 (1997), 669-713.
- [16] S. Foucart and H. Rauhut, A Mathematical Introduction to Compressive Sensing, Springer, Birkhuser Basel, 2013.
- [17] D. Ge, X. Jiang and Y. Ye, A note on the complexity of L_p minimization, Math. Program., 129(2011), 285-299.

- [18] D. Geman and G. Reynolds, Constrained restoration and the recovery of discontinuities, *IEEE Trans. Pattern Anal. Mach. Intell.*, 14(1992), 357-383.
- [19] A.W. Ingleton, A problem in linear inequalities, *Proc. London Math. Soc.*, 16(1966), 519-536.
- [20] L. Han, A. Tiwari, M.K. Camlibel and J.-S. Pang, Convergence of time-stepping schemes for passive and extended linear complementarity systems, *SIAM J. Numer. Anal.*, 47(2009), 3768-3796.
- [21] B.K. Natarajan, Sparse approximate solutions to linear systems, *SIAM J. Computing*, 24(1995), 227-234.
- [22] H.M. Markowitz, *Portfolio selection: efficient diversification of investments*, New York, Wiley, 1959.
- [23] S. Xu, On local w-uniqueness of solutions to linear complementarity problem, *Linear Algebra Appl.*, 290(1999), 23-29.
- [24] Y. Ye, On homogeneous and self-dual algorithms for LCP, *Math. Program.*, 76(1996), 211-221.