

42 There exists a large amount of literature on numerical solutions and numerical analysis
 43 of the NLS equation, see [10, 33, 28, 22, 1, 2, 4, 5, 6, 36, 21, 25, 38, 14, 28, 13]. To the best of
 44 our knowledge, all the existing mass- and energy-conserving methods have only second-order
 45 accuracy in time and is of the Crank–Nicolson type. No higher-order time-stepping schemes,
 46 which conserve both mass and energy, have been reported in the literature. Moreover, the
 47 existing error estimates for nonlinearly implicit schemes for the NLS equation generally
 48 require certain grid-ratio conditions. The standard grid-ratio conditions in the literature are
 49 $\tau = o(h^{\frac{d}{4}})$ for the cubic NLS equation and $\tau = o(h^{\frac{d}{2}})$ for general nonlinearity, where h and τ
 50 denote the spatial and temporal mesh sizes. Karakashian and Makridakis [22, 23] proposed
 51 some continuous and discontinuous space-time Galerkin finite element methods for the cubic
 52 NLS equation and proved optimal-order convergence under a weaker grid-ratio condition
 53 $\tau^{k-1} |\ln h| \rightarrow 0$ in two dimensions, where $k \geq 2$ is the degree of finite elements in time. For
 54 the defocusing cubic NLS equation (or the focusing cubic NLS equation with sufficiently
 55 small initial data), using the energy conservation of the numerical scheme, error estimates
 56 were established without grid-ratio condition in [17, 37]. For general nonlinearity (possibly
 57 focusing), Wang [36] established an error estimate for a linearized semi-implicit scheme
 58 without grid-ratio condition; Henning and Peterseim [20] established an error estimate for
 59 the nonlinearly implicit Crank–Nicolson finite element method without grid-ratio condition.
 60 Both [36] and [20] used an error splitting technique in which they proved boundedness of the
 61 numerical solutions by establishing an L^∞ -norm error estimate between the fully discrete
 62 and the semidiscrete-in-time numerical solutions. The error splitting technique allows to
 63 avoid grid-ratio conditions in using the inverse inequality.

64 The objective of this paper is to develop a family of arbitrarily higher-order mass-
 65 and energy-conserving fully discrete space-time finite element methods based on the scalar
 66 auxiliary variable (SAV) formulation of the NLS equation, and to establish the existence,
 67 uniqueness and optimal order convergence of numerical solutions without grid-ratio con-
 68 dition. Two key ideas are utilized in our construction of the method. First, the SAV
 69 reformulation of the NLS equation is used. This approach was introduced in [31, 30] as an
 70 enhanced version of the invariant energy quadratization (IEQ) approach [39, 40, 41, 42],
 71 for developing energy-decay methods for dissipative (gradient flow) systems. Here we adapt
 72 the SAV approach to the dispersive NLS equation, and the SAV reformulation is essential
 73 to enable our methods to maintain the energy conservation property at the discrete level.
 74 Second, the Gauss collocation method is used for time discretization in the SAV formula-
 75 tion of the NLS equation. The method can be viewed as an efficient implementation of the
 76 space-time finite element methods for the SAV formulation with Gauss quadrature in time.
 77 The Gauss collocation method was combined with IEQ and SAV to preserve energy decay
 78 in solving phase field equations in [3, 18, 19]. We adopt this method here to preserve mass
 79 conservation without affecting the energy conservation structure of the SAV formulation.

80 The SAV formulation introduces new difficulties to error analysis for the NLS equation
 81 due to the presence of $\partial_t u$ in the equation of r , see equation (2.2b), which leads to a con-
 82 sistency error of sub-optimal order in time and introduces new difficulty in obtaining the
 83 stability estimate. These difficulties are overcome by combining three techniques. First,
 84 inspired by the error analysis of Karakashian and Makridakis [23], our proof makes use of
 85 properties of the Legendre polynomials on each interval I_n , rewriting the Gauss collocation
 86 method into a space-time Galerkin finite element method, which makes it easier to choose
 87 suitable test functions in the error estimation. Second, we introduce a temporal Ritz projec-
 88 tion and use a super-approximation result of the temporal local L^2 projection to eliminate
 89 the sub-optimal temporal consistency error caused by $\partial_t u$ in the equation of r . Third,
 90 we estimate the time derivative of the error in $H^{-1}(\Omega)$ using a duality argument, which
 91 leads to an optimal-order H^1 -norm error estimate. We prove the existence, uniqueness and
 92 optimal-order convergence of numerical solutions based on Schaefer’s fixed point theorem in
 93 an L^∞ -neighborhood of the exact solution. This allows us to avoid grid-ratio conditions for

94 the NLS equation with general nonlinearity.

95 The rest of this paper is organized as follows. In Section 2, we present the SAV reformulation of the NLS equation and introduce our SAV space-time Gauss collocation finite element method. In Section 3, we first present an integral reformulation of the proposed method and then establish its mass and energy conservation properties. We also derive a consistency error estimate for the method, which is vitally used to prove an error estimate in the subsequent section. In Section 4, we first establish the well-posedness of the numerical method and then prove an error bound of the form $O(h^p + \tau^{k+1})$ in the energy norm, where τ and h denote the temporal and spatial mesh sizes, respectively, with (p, k) denoting the degree of polynomials in the space-time finite element method. Finally, in Section 5, we present a few numerical tests to validate the theoretical results, and to demonstrate the effectiveness of the proposed method in preserving the shape of a soliton wave.

104 Throughout this paper, unless stated otherwise, C will be used to denote a generic positive constant which is independent of τ , h , n and N , but may depend on T and the regularity of solution.

109 **2. Formulation of the SAV–Gauss collocation finite element method.** In this section, we construct a Gauss collocation finite element method based on the SAV reformulation of the NLS equation.

110 **2.1. Function spaces.** Let $H^k(\Omega)$, $k \geq 0$, be the conventional complex-valued Sobolev space of functions on Ω , and denote

$$L^2(\Omega) = H^0(\Omega) \quad \text{and} \quad H_0^1(\Omega) = \{v \in H^1(\Omega) : v = 0 \text{ on } \partial\Omega\}.$$

We denote by (\cdot, \cdot) and $\|\cdot\|$ the inner product and norm of the complex-valued Hilbert space $L^2(\Omega)$, respectively, defined by

$$(u, v) := \int_{\Omega} u \bar{v} \, dx \quad \text{and} \quad \|u\| := \sqrt{(u, u)}.$$

112 For $m, s \geq 0$ and $1 \leq p \leq \infty$, the notation $W^{m,p}(0, T; H^s(\Omega))$ stands for the space-time Sobolev space of functions which are $W^{m,p}$ in time and H^s in space; see [11, Chapter 5.9]. We abbreviate the norms of $H^s(\Omega)$ and $W^{m,p}(0, T; H^s(\Omega))$ as $\|\cdot\|_{H^k}$ and $\|\cdot\|_{W^{m,p}(I_n; H^s)}$, respectively, omitting the dependence on Ω in the subscripts.

116 **2.2. The SAV reformulation of (1.1).** The SAV formulation of the NLS equation (cf. [30]) introduces a scalar auxiliary variable

$$118 \quad (2.1) \quad r = \sqrt{\int_{\Omega} \frac{1}{2} F(|u|^2) dx + c_0} \quad \text{with} \quad g(u) = \frac{f(|u|^2)}{\sqrt{\int_{\Omega} \frac{1}{2} F(|u|^2) dx + c_0}},$$

119 with a positive c_0 (which guarantees that the function r has a positive lower bound), and reformulate (1.1) as

$$122 \quad (2.2a) \quad i\partial_t u - \Delta u - rg(u)u = 0 \quad \text{in } \Omega \times (0, T],$$

$$123 \quad (2.2b) \quad \frac{dr}{dt} = \text{Re}\left(\frac{1}{2}g(u)u, \partial_t u\right) \quad \text{in } \Omega \times (0, T],$$

$$124 \quad (2.2c) \quad u = 0 \quad \text{on } \partial\Omega \times (0, T],$$

$$125 \quad (2.2d) \quad u = u_0, \quad r = r_0 \quad \text{in } \Omega \times \{0\},$$

127 where $r_0 = \sqrt{\int_{\Omega} \frac{1}{2} F(|u_0|^2) dx + c_0}$. The mass and energy conservation in the SAV formulation are

$$129 \quad (2.3) \quad \frac{d}{dt} \int_{\Omega} |u|^2 dx = 0, \quad \text{and} \quad \frac{d}{dt} \left(\frac{1}{2} \int_{\Omega} |\nabla u|^2 dx - r^2 + c_0 \right) = 0.$$

131 **2.3. Space-time finite element spaces.** Let \mathcal{T}_h be a shape-regular and quasi-uniform triangulation of Ω with mesh size $h \in (0, 1)$ and $\{t_n\}_{n=0}^N$ be a uniform partition of $[0, T]$

133 with the time step size $\tau \in (0, 1)$, where N is a positive integer and hence $\tau = \frac{T}{N}$. For an
 134 integer $p \geq 1$ we denote by \mathbb{Q}^p the space of complex-valued polynomials of degree $\leq p$ in
 135 space, and we denote by S_h the complex-valued Lagrange finite element space subject to
 136 the triangulation of Ω , defined by

$$137 \quad S_h = \{v \in C(\overline{\Omega}) : v|_K \in \mathbb{Q}^p \text{ for all } K \in \mathcal{T}_h, v = 0 \text{ on } \partial\Omega\},$$

139 where $C(\overline{\Omega})$ denotes the space of complex-valued uniformly continuous functions on Ω . Then
 140 S_h is a complex Hilbert spaces with the inner product (\cdot, \cdot) and norm $\|\cdot\|$.

141 For an integer $k \geq 1$, let \mathbb{P}^k denote the space of real-valued polynomials of degree $\leq k$
 142 in t . For a Banach space X , such as $X = L^2(\Omega)$ or $X = S_h$, we define the following
 143 tensor-product space:

$$144 \quad (2.4) \quad \mathbb{P}^k \otimes X := \text{span}\{p(t)\phi(x) : p \in \mathbb{P}^k, \phi \in X\} = \left\{ \sum_{j=0}^k t^j \phi_j : \phi_j \in X \right\}.$$

Moreover, let $P_h : L^2(\Omega) \rightarrow S_h$ denote the L^2 projection operator defined by

$$(w - P_h w, v_h) = 0 \quad \forall v_h \in S_h, \forall w \in L^2(\Omega).$$

146 The following stability properties are well-known (cf. [8]):

$$147 \quad (2.5a) \quad \|P_h w\| \leq \|w\| \quad \forall w \in L^2(\Omega),$$

$$148 \quad (2.5b) \quad \|P_h w\|_{H^1} \leq C \|w\|_{H^1} \quad \forall w \in H_0^1(\Omega),$$

150 where C depends only on the shape-regularity and quasi-uniformity of the mesh.

151 We also introduce the global space-time finite element spaces

$$152 \quad (2.6) \quad X_{\tau,h} = \{v_h \in C([0, T]; S_h) : v_h|_{I_n} \in \mathbb{P}^k \otimes S_h \text{ for } n = 1, \dots, N\},$$

$$153 \quad (2.7) \quad Y_{\tau,h} = \{q_h \in C([0, T]) : q_h|_{I_n} \in \mathbb{P}^k \text{ for } n = 1, \dots, N\}.$$

155 **2.4. SAV–Gauss collocation finite element method.** Let c_j and w_j , $j = 1, \dots, k$,
 156 be the nodes and weights of the k -point Gauss quadrature rule in the interval $[-1, 1]$ (see
 157 [32, Table 3.1]), and let $t_{nj} = t_{n-1} + (1 + c_j)\tau/2$, $j = 1, \dots, k$ denote the Gauss points in the
 158 interval $I_n = [t_{n-1}, t_n]$. We define the following Gauss collocation finite element method for
 159 (2.2).

160 Main Algorithm

161 *Step 1:* Set $u_h^0 := I_h u_0$ and $r_h^0 := r_0$, where I_h is the Lagrange interpolation operator
 162 onto the finite element space. Determine $(u_h, r_h) \in X_{\tau,h} \times Y_{\tau,h}$ by the following two steps.

163 *Step 2:* For $n = 1, 2, \dots, N$, define $\{(u_h(t_{nj}), r_h(t_{nj}))\}_{j=1}^k \subset S_h \times \mathbb{R}$ by solving recur-
 164 sively (in n) the following nonlinear (algebraic) system:

$$165 \quad (2.8a) \quad i(\partial_t u_h(t_{nj}), v_h) + (\nabla u_h(t_{nj}), \nabla v_h) \\ 166 \quad - (r_h(t_{nj})g(u_h(t_{nj}))u_h(t_{nj}), v_h) = 0, \quad \forall v_h \in S_h,$$

$$167 \quad (2.8b) \quad \partial_t r_h(t_{nj}) = \frac{1}{2} \text{Re}(g(u_h(t_{nj}))u_h(t_{nj}), \partial_t u_h(t_{nj})),$$

$$168 \quad (2.8c) \quad u_h(t_{n-1}) = u_h^{n-1} \quad \text{and} \quad r_h(t_{n-1}) = r_h^{n-1}.$$

170 *Step 3:* Set $u_h^n := u_h(t_n)$ and $r_h^n := r_h(t_n)$.

171 **REMARK 2.1.** (a) We note that in (2.8a) and (2.8b), $\partial_t u_h(t_{nj}) = \partial_t u_h(t)|_{t=t_{nj}}$ and
 172 $\partial_t r_h(t_{nj}) = \partial_t r_h(t)|_{t=t_{nj}}$. Main Algorithm actually computes $\{(u_h(t_{nj}), r_h(t_{nj}))\}_{j=1}^k$ for
 173 each $n \geq 1$, however, since any k th order polynomial on I_n is uniquely determined by its
 174 initial value at t_{n-1} and its values at the k Gauss points t_{nj} , $j = 1, \dots, k$, then the Gauss-
 175 point values generated by Main Algorithm uniquely determine the pair $(u_h, r_h) \in X_{\tau,h} \times Y_{\tau,h}$.

176 (b) Each of (2.8a) and (2.8b) consists of nonlinear algebraic equations, note that the
 177 test function v_h can be different for different j , and one ‘‘initial condition’’ is prescribed for

178 each of u_h and r_h . The number of equations imposed is the same as the degree of freedoms
 179 which equals the dimension of the space $\mathbb{P}^k \otimes S_h$ for each n .

180 (c) Main Algorithm can be obtained by applying the Gauss quadrature rule (in time)
 181 to a (continuous) space-time finite element method for (2.2); see Section 3.1.

(d) In practical computation, we solve the solution of the nonlinear scheme (2.8) by
 Newton's method: For given $\{(u_h^{\ell-1}(t_{nj}), r_h^{\ell-1}(t_{nj}))\}_{j=1}^k \subset S_h \times \mathbb{R}$, find

$$\{(u_h^\ell(t_{nj}), r_h^\ell(t_{nj}))\}_{j=1}^k \subset S_h \times \mathbb{R}$$

182 satisfying the linearized equations

$$\begin{aligned} \text{(2.9a)} \quad & i(\partial_t u_h^\ell(t_{nj}), v_h) + (\nabla u_h^\ell(t_{nj}), \nabla v_h) \\ & = (r_h^\ell(t_{nj})g(u_h^{\ell-1}(t_{nj}))u_h^{\ell-1}(t_{nj}), v_h) \\ & \quad + (r_h^{\ell-1}(t_{nj})g_1(u_h^{\ell-1}(t_{nj}))(u_h^\ell(t_{nj}) - u_h^{\ell-1}(t_{nj})), v_h) \\ & \quad + (r_h^{\ell-1}(t_{nj})g_2(u_h^{\ell-1}(t_{nj}))(\bar{u}_h^\ell(t_{nj}) - \bar{u}_h^{\ell-1}(t_{nj})), v_h), \quad \forall v_h \in S_h, \end{aligned}$$

$$\begin{aligned} \text{(2.9b)} \quad & \partial_t r_h^\ell(t_{nj}) = \frac{1}{2} \text{Re}(g(u_h^{\ell-1}(t_{nj}))u_h^{\ell-1}(t_{nj}), \partial_t u_h^\ell(t_{nj})) \\ & \quad + \frac{1}{2} \text{Re}(g_1(u_h^{\ell-1}(t_{nj}))(u_h^\ell(t_{nj}) - u_h^{\ell-1}(t_{nj})), \partial_t u_h^{\ell-1}(t_{nj})) \\ & \quad + \frac{1}{2} \text{Re}(g_2(u_h^{\ell-1}(t_{nj}))(\bar{u}_h^\ell(t_{nj}) - \bar{u}_h^{\ell-1}(t_{nj})), \partial_t u_h^{\ell-1}(t_{nj})) \end{aligned}$$

$$\text{(2.9c)} \quad u_h^\ell(t_{n-1}) = u_h^{n-1} \quad \text{and} \quad r_h^\ell(t_{n-1}) = r_h^{n-1},$$

189 where

$$g_1(u) := \partial_u[g(u)u] \quad \text{and} \quad g_2(u) := \partial_{\bar{u}}[g(u)u],$$

and $\partial_{\bar{u}}$ denotes the differentiation with respect to \bar{u} in the expression of

$$g(u)u = \frac{f(u\bar{u})u}{\sqrt{\int_{\Omega} \frac{1}{2} F(u\bar{u}) dx + c_0}}.$$

192 The iteration in ℓ is set to stop when the desired tolerance error is achieved.

193 3. Conservation, stability and consistency analysis.

194 **3.1. A reformulation of scheme (2.8a)–(2.8b).** In this subsection, we present several
 195 integral identities and inequalities, including a reformulation of Main Algorithm. These
 196 identities and inequalities will be used in the subsequent analysis of existence, uniqueness
 197 and convergence of numerical solutions.

198 Consider the interval $I_n = [t_{n-1}, t_n]$, then we define $P_\tau^n : L^2(I_n; L^2(\Omega)) \rightarrow \mathbb{P}^{k-1} \otimes L^2(\Omega)$
 199 to be the L^2 projection defined by

$$\text{(3.1)} \quad \int_{I_n} (u - P_\tau^n u, v) dt = 0 \quad \forall v \in \mathbb{P}^{k-1} \otimes L^2(\Omega).$$

200 Thus $u - P_\tau^n u$ is orthogonal to all temporal polynomials of degree $\leq k - 1$, which means
 201 that if $u \in \mathbb{P}^k \otimes L^2(\Omega)$ then

$$\text{(3.2)} \quad u - P_\tau^n u = \phi_{n-1} L_k,$$

202 where $\phi_{n-1} \in L^2(\Omega)$ and

$$\text{(3.3)} \quad L_k(t) := \widehat{L}_k\left(\frac{2t - t_{n-1} - t_n}{\tau}\right)$$

203 is the shifted Legendre polynomial (orthogonal to polynomials of lower degree on I_n). The

210 temporal L^2 projection operator P_τ^n has the following approximation property (cf. [9]):

$$211 \quad (3.4) \quad \max_{t \in I_n} \|v - P_\tau^n v\|_X \leq C\tau^m \max_{t \in I_n} \|\partial_t^m v\|_X, \quad 0 \leq m \leq k,$$

212 for all $v \in C^k([0, T]; X)$, where $X = \mathbb{R}$ or $X = H^s(\Omega)$ for some $s \in \mathbb{R}$.

213 Since the k -point Gauss quadrature holds exactly for polynomials of degree $2k - 1$ (cf.
214 [16, p. 222]), and the Gauss points $t_{nj}, j = 1, \dots, k$, are the roots of the Legendre polynomial
215 $L_k(t)$ (cf. [24, p. 33]), it follows that the following two identities hold:
216

$$217 \quad (3.5) \quad \int_{I_n} v(t) dt = \frac{\tau}{2} \sum_{j=1}^k v(t_{nj}) w_j \quad \forall v \in \mathbb{P}^{2k-1} \otimes S_h,$$

$$218 \quad (3.6) \quad v(t_{nj}) = P_\tau^n v(t_{nj}) \quad \forall v \in \mathbb{P}^k \otimes S_h.$$

219 Setting $v_h = \frac{\tau}{2} v_h(t_{nj}) w_j$ in (2.8a) and summing up the results for $j = 1, \dots, k$, and
220 using (3.5)–(3.6) in the first two terms yield the following integral identity:
221

$$222 \quad (3.7) \quad \int_{I_n} i(\partial_t u_h, v_h) dt + \int_{I_n} (\nabla P_\tau^n u_h, \nabla v_h) dt \\ 223 \quad - \frac{\tau}{2} \sum_{j=1}^k w_j (r_h(t_{nj}) g(u_h(t_{nj})) u_h(t_{nj}), v_h(t_{nj})) = 0 \quad \forall v_h \in \mathbb{P}^k \otimes S_h.$$

224 Similarly, multiplying (2.8b) by $\frac{\tau}{2} q_h(t_{nj}) w_j$ and summing up the results for $j = 1, \dots, k$,
225 and using (3.5) in the first term, we have
226

$$227 \quad (3.8) \quad \int_{I_n} \partial_t r_h q_h dt = \frac{\tau}{2} \sum_{j=1}^k \frac{w_j}{2} \operatorname{Re}(g(u_h(t_{nj})) u_h(t_{nj}), \partial_t u_h(t_{nj}) q_h(t_{nj})) \quad \forall q_h \in \mathbb{P}^k.$$

228 (3.7)–(3.8) provides a reformulation of Main Algorithm. The above reformulation will be
229 crucially used later to show mass and energy conservations, as well as existence, uniqueness
230 and convergence of numerical solutions.
231

232 From (3.2) we get

$$233 \quad \|\phi_{n-1}\| = \frac{1}{|L_k(t_{n-1})|} \|u_h(t_{n-1}) - P_\tau^n u_h(t_{n-1})\| \\ 234 \quad \leq C \|u_h(t_{n-1})\| + C \left(\frac{1}{\tau} \int_{I_n} \|P_\tau^n u_h(t)\|^2 dt \right)^{\frac{1}{2}},$$

235 where we have used the inverse inequality in time. Thus, by using (3.2) again, we obtain
236 the following inequality:
237

$$238 \quad (3.9) \quad \int_{I_n} \|u_h\|^2 dt \leq C \int_{I_n} \|P_\tau^n u_h\|^2 dt + C\tau \|u_h(t_{n-1})\|^2 \quad \forall u_h \in \mathbb{P}^k \otimes S_h.$$

239 By using the two identities (3.5)–(3.6), one can also prove the following inequality:
240

$$241 \quad (3.10) \quad \frac{\tau}{2} \sum_{j=1}^k w_j \|v_h(t_{nj})\|^2 = \int_{I_n} \|P_\tau^n v_h(t)\|^2 dt \leq \int_{I_n} \|v_h(t)\|^2 dt \quad \forall v_h \in \mathbb{P}^k \otimes S_h.$$

242 The inequalities (3.9)–(3.10) will be frequently used in the subsequent error analysis.
243

244 **3.2. Mass and energy conservation properties.** In this subsection, we prove the
245 following conservation properties of the numerical solution, which comprise of the first main
246 theorem of this paper.

247 **THEOREM 3.1.** *Let $(u_h, r_h) \in X_{\tau,h} \times Y_{\tau,h}$ be a solution of Main Algorithm, then the*

248 following mass and energy conservations hold:

$$\begin{aligned} \frac{1}{2}\|u_h(t_n)\|^2 &= \frac{1}{2}\|u_h(t_0)\|^2 && \text{for } n \geq 1, \\ \frac{1}{2}\|\nabla u_h(t_n)\|^2 - |r_h(t_n)|^2 + c_0 &= \frac{1}{2}\|\nabla u_h(t_0)\|^2 - |r_h(t_0)|^2 + c_0 && \text{for } n \geq 1. \end{aligned}$$

251 *Proof.* Setting $v_h = u_h \in \mathbb{P}^k \otimes S_h$ in (3.7) and taking the imaginary part yield

$$\begin{aligned} (3.11) \quad \operatorname{Im} \int_{I_n} i(\partial_t u_h, u_h) dt &= -\operatorname{Im} \int_{I_n} (\nabla P_\tau^n u_h, \nabla u_h) dt \\ &+ \operatorname{Im} \left[\frac{\tau}{2} \sum_{j=1}^k w_j(r_h(t_{n_j})g(u_h(t_{n_j})), |u_h(t_{n_j})|^2) \right] = 0, \end{aligned}$$

252 where we have used the definition of the projection operator P_τ^n , which implies

$$\operatorname{Im} \int_{I_n} (\nabla P_\tau^n u_h, \nabla u_h) dt = \operatorname{Im} \int_{I_n} (\nabla P_\tau^n u_h, \nabla P_\tau^n u_h) dt = 0.$$

253 Then the mass conservation follows from (3.11) and the identity

$$\operatorname{Im} \int_{I_n} i(\partial_t u_h, u_h) dt = \frac{1}{2}\|u_h(t_n)\|^2 - \frac{1}{2}\|u_h(t_{n-1})\|^2.$$

255 Alternatively, setting $v_h = \partial_t u_h$ and $q_h = 2r_h$ in (3.7) and (3.8), respectively, and taking
256 the real parts yield

$$(3.12) \quad \operatorname{Re} \int_{I_n} (\nabla P_\tau^n u_h, \nabla \partial_t u_h) dt = \frac{\tau}{2} \operatorname{Re} \sum_{j=1}^k w_j(r_h(t_{n_j})g(u_h(t_{n_j})), u_h(t_{n_j}), \partial_t u_h(t_{n_j}))$$

$$(3.13) \quad |r_h(t_n)|^2 - |r_h(t_{n-1})|^2 = \frac{\tau}{2} \operatorname{Re} \sum_{j=1}^k w_j(r_h(t_{n_j})g(u_h(t_{n_j})), u_h(t_{n_j}), \partial_t u_h(t_{n_j})).$$

258 Since

$$\begin{aligned} \operatorname{Re} \int_{I_n} (\nabla P_\tau^n u_h, \nabla \partial_t u_h) dt &= \operatorname{Re} \int_{I_n} (P_\tau^n \nabla u_h, \nabla \partial_t u_h) dt = \operatorname{Re} \int_{I_n} (\nabla u_h, \nabla \partial_t u_h) dt \\ &= \frac{1}{2}\|\nabla u_h(t_n)\|^2 - \frac{1}{2}\|\nabla u_h(t_{n-1})\|^2, \end{aligned}$$

262 it follows that

$$\begin{aligned} (3.14) \quad \frac{1}{2}\|\nabla u_h(t_n)\|^2 - \frac{1}{2}\|\nabla u_h(t_{n-1})\|^2 \\ = \frac{\tau}{2} \operatorname{Re} \sum_{j=1}^k w_j(r_h(t_{n_j})g(u_h(t_{n_j})), u_h(t_{n_j}), \partial_t u_h(t_{n_j})). \end{aligned}$$

263 Subtracting (3.13) from (3.14) yields

$$(3.15) \quad \frac{1}{2}\|\nabla u_h(t_n)\|^2 - |r_h(t_n)|^2 = \frac{1}{2}\|\nabla u_h(t_{n-1})\|^2 - |r_h(t_{n-1})|^2 \quad \text{for } n \geq 1.$$

264 Thus, the energy conservation holds. The proof is complete. \square

272 **3.3. An upper bound of mass at internal stages.** In this subsection, we prove
273 that the average mass of numerical solutions at internal stages has an upper bound uncondi-
274 tionally (independent of the regularity of solutions). This property furthermore strengthens
275 the stability of numerical solutions when the exact solution is not smooth (for example, close
276 to blow up).

277 **THEOREM 3.2.** *Let $(u_h, r_h) \in X_{\tau,h} \times Y_{\tau,h}$ be a solution of Main Algorithm, then the*

278 following inequalities hold:

$$279 \quad (3.16a) \quad \max_{1 \leq n \leq N} \frac{1}{\tau} \int_{I_n} \|P_\tau^n u_h\|^2 dt \leq \|u_h(0)\|^2,$$

$$280 \quad (3.16b) \quad \max_{1 \leq n \leq N} \max_{1 \leq j \leq k} \|u_h(t_{nj})\| \leq C \|u_h(0)\|,$$

281 where C is a constant independent of τ , h and the regularity of the solution.

282 *Proof.* By the definition of the temporal L^2 projection P_τ^n , we get

$$284 \quad (3.17) \quad \int_{I_n} \|P_\tau^n u_h(t)\|^2 dt = \operatorname{Re} \int_{I_n} (u_h(t), P_\tau^n u_h(t)) dt$$

$$285 \quad = \operatorname{Re} (u_h(t_{n-1}), P_\tau^n u_h(t_{n-1}))\tau + \operatorname{Re} \int_{I_n} (\partial_t u_h(t), (t_n - t)P_\tau^n u_h(t)) dt$$

$$286 \quad + \operatorname{Re} \int_{I_n} (u_h(t), (t_n - t)\partial_t P_\tau^n u_h(t)) dt =: J_1 + J_2 + J_3,$$

287 where we have interchanged the order of integration in deriving the second to last equality.

288 It can be shown that (cf. [12]) that $J_2 = 0$ and

$$290 \quad J_1 \leq \frac{\tau}{2} \|u_h(t_{n-1})\|^2 + \frac{\tau}{2} \|P_\tau^n u_h(t_{n-1})\|^2,$$

$$291 \quad J_3 = -\frac{\tau}{2} \|P_\tau^n u_h(t_{n-1})\|^2 + \int_{I_n} \frac{1}{2} \|P_\tau^n u_h(t)\|^2 dt.$$

292 Substituting the estimates of J_1 , J_2 and J_3 into (3.17) gives (3.16a).

293 Substituting (3.16a) into (3.9) and using the mass conservation property again, we
294 obtain $\int_{I_n} \|u_h\|^2 dt \leq C\tau \|u_h(0)\|^2$, which and an application of the inverse inequality yield
295 (3.16b). The proof is complete. \square

296 **3.4. Temporal and spatial Ritz projections.** Let $I_\tau^n u$ and $I_\tau^n r$ be the temporal
297 Lagrange interpolation polynomials of u and r , respectively, interpolated at the $k+1$ points
298 t_{n-1} and t_{nj} , $j = 1, \dots, k$. It is well known that the following approximation property (cf.
299 [9]):

$$301 \quad (3.18) \quad \max_{t \in I_n} (\|v - I_\tau^n v\|_X + \tau \|\partial_t(v - I_\tau^n v)\|_X) \leq C\tau^{m+1} \max_{t \in I_n} \|\partial_t^{m+1} v\|_X$$

302 for all $v \in C^{m+1}([0, T]; X)$, $0 \leq m \leq k$, and $X = \mathbb{R}$ or $X = H^s(\Omega)$ for some $s \in \mathbb{R}$. We also
303 define a temporal Ritz projection operator $R_\tau^n : W^{1,\infty}(I_n; L^2(\Omega)) \rightarrow \mathbb{P}^k \otimes L^2(\Omega)$ as follows:

$$305 \quad (3.19) \quad \int_{I_n} (\partial_t(u - R_\tau^n u), v) dt = 0 \quad \forall v \in \mathbb{P}^{k-1} \otimes L^2(\Omega),$$

$$306 \quad (3.20) \quad u(t_{n-1}) - R_\tau^n u(t_{n-1}) = 0.$$

307 By using this property and the shifted Legendre polynomials defined in (3.3), we can express
308 the temporal Ritz projection as

$$310 \quad (3.21) \quad R_\tau^n u(t) = u(t_{n-1}) + \sum_{j=0}^{k-1} \frac{\int_{I_n} L_j(s) \partial_s u(s) ds}{\int_{I_n} |L_j(s)|^2 ds} \int_{t_{n-1}}^t L_j(s) ds,$$

311 which implies that if $X \subset L^2(\Omega)$ is a Banach space and $u \in W^{1,\infty}(I_n; X)$, then $R_\tau^n u$ is
312 automatically in $\mathbb{P}^k \otimes X$. It can be shown that R_τ^n satisfies the following approximation
313 property, see [12, Lemma 3.3].

314 **LEMMA 3.3.** *Let $X = \mathbb{R}$ or $H^s(\Omega)$ for some $s \geq 0$. For $u \in W^{m+1,\infty}(I_n; X)$, with
315 $0 \leq m \leq k$, the following approximation property holds:*

$$316 \quad \|u - R_\tau^n u\|_{L^\infty(I_n; X)} + \tau \|\partial_t(u - R_\tau^n u)\|_{L^\infty(I_n; X)} \leq C\tau^{m+1} \|u\|_{W^{m+1,\infty}(I_n; X)}.$$

317 In addition to the above optimal-order approximation result, we also have the following
318

320 super-convergence result.

321 LEMMA 3.4 (A super-approximation property). *Let $X = \mathbb{R}$ or $H^s(\Omega)$ for some $s \geq 0$.*
 322 *If $w \in W^{k,\infty}(I_n; W^{s,\infty}(\Omega))$ and $v \in \mathbb{P}^{k-1} \otimes X$, then*

$$323 \quad \|wv - P_\tau^n(wv)\|_{L^2(I_n; X)} \leq C\tau \|v\|_{L^2(I_n; X)}.$$

325 *Proof.* We only give a proof for the case $X = H^s(\Omega)$ because the other cases are similar.
 326 By applying (3.4) with $m = k$, we have

$$\begin{aligned} 327 \quad \|wv - P_\tau^n(wv)\|_{L^2(I_n; H^s)} &\leq C\tau^{\frac{1}{2}} \|wv - P_\tau^n(wv)\|_{L^\infty(I_n; H^s)} \\ 328 \quad &\leq C\tau^{k+\frac{1}{2}} \|\partial_t^k(wv)\|_{L^\infty(I_n; H^s)} \\ 329 \quad &\leq C \sum_{m=0}^{k-1} \tau^{k+\frac{1}{2}} \|\partial_t^{k-m} w \partial_t^m v\|_{L^\infty(I_n; H^s)} \quad (\text{since } \partial_t^k v = 0) \\ 330 \quad &\leq C \sum_{m=0}^{k-1} \tau^{k+\frac{1}{2}} \|\partial_t^{k-m} w\|_{L^\infty(I_n; W^{s,\infty})} \|\partial_t^m v\|_{L^\infty(I_n; H^s)} \\ 331 \quad &\leq C \sum_{m=0}^{k-1} \tau^{k+\frac{1}{2}-m} \|v\|_{L^\infty(I_n; H^s)} \\ 332 \quad &\leq C\tau^{\frac{3}{2}} \|v\|_{L^\infty(I_n; H^s)} \\ 333 \quad &\leq C\tau \|v\|_{L^2(I_n; H^s)}. \end{aligned}$$

335 here we have used the inverse inequality in time twice above. The proof is complete. \square

Finally, we also recall the (spatial) Ritz projection operator $R_h : H_0^1(\Omega) \rightarrow S_h$ defined by

$$(\nabla(w - R_h w), \nabla v_h) = 0 \quad \forall v_h \in S_h, \quad \forall w \in H_0^1(\Omega),$$

336 and the discrete Laplacian operator $\Delta_h : S_h \rightarrow S_h$ defined by

$$337 \quad (3.22) \quad (\Delta_h \phi_h, \chi_h) := -(\nabla \phi_h, \nabla \chi_h) \quad \forall \phi_h, \chi_h \in S_h.$$

339 It is known [8] that there hold the following identities:

$$340 \quad (3.23a) \quad P_h \Delta v = \Delta_h R_h v \quad \forall v \in H_0^1(\Omega),$$

$$341 \quad (3.23b) \quad R_\tau^n R_h v = R_h R_\tau^n v \quad \forall v \in W^{1,\infty}(I_n; H_0^1(\Omega)),$$

$$342 \quad (3.23c) \quad R_\tau^n \Delta_h v_h = \Delta_h R_\tau^n v_h \quad \forall v \in W^{1,\infty}(I_n; S_h).$$

344 Moreover, there holds the following approximation property (cf. [8]):

$$345 \quad (3.24) \quad \|v - R_h v\|_{H^1} \leq Ch^p \|v\|_{H^{p+1}} \quad \forall v \in H_0^1(\Omega) \cap H^{p+1}(\Omega).$$

3.5. Consistency of scheme (2.8a)–(2.8b). We define a pair of intermediate solutions (for comparison with the numerical solutions)

$$u_h^* = R_\tau^n R_h u \quad \text{and} \quad r_h^* = R_\tau^n r,$$

347 and the following consistency error functions:

$$348 \quad (3.25) \quad d_u^n := i\partial_t R_\tau^n (R_h u - u) + \Delta_h R_h (u - R_\tau^n u) + rg(u)u - I_\tau^n [r_h^* g(u_h^*) u_h^*],$$

$$349 \quad (3.26) \quad d_r^n := \frac{1}{2} \text{Re} [(g(u)u, \partial_t u) - I_\tau^n (g(u_h^*) u_h^*, \partial_t u_h^*)].$$

350

351 It is easy to check that there hold

$$352 \quad (3.27) \quad \int_{I_n} i(\partial_t u_h^*, v_h) dt + \int_{I_n} (\nabla P_\tau^n u_h^*, \nabla v_h) dt$$

$$353 \quad - \frac{\tau}{2} \sum_{j=1}^k w_j (r_h^*(t_{nj}) g(u_h^*(t_{nj})) u_h^*(t_{nj}), v_h(t_{nj})) = \int_{I_n} (P_\tau^n d_u^n, v_h) dt,$$

$$354 \quad (3.28) \quad \int_{I_n} \partial_t r_h^* q_h dt = \frac{\tau}{4} \sum_{j=1}^k w_j \operatorname{Re}(q_h(t_{nj}) g(u_h^*(t_{nj})) u_h^*(t_{nj}), \partial_t u_h^*(t_{nj})) + \int_{I_n} P_\tau^n d_r^n q_h dt.$$

355

356 **THEOREM 3.5.** *Suppose that the solution of (1.1) is sufficiently smooth, then $d_u^n \in$*
 357 *$C(I_n; H_0^1(\Omega))$ and there hold*

$$358 \quad (3.29) \quad \sup_{t \in I_n} \|d_u^n\|_{H^1} \leq C(h^p + \tau^{k+1}) \quad \text{and} \quad \sup_{t \in I_n} |P_\tau^n d_r^n| \leq C(h^p + \tau^{k+1}).$$

359

360 *Proof.* Since the spatial Ritz projection R_h maps $H_0^1(\Omega)$ into $S_h \subset H_0^1(\Omega)$, and the
 361 temporal Ritz projection R_τ^n maps $W^{1,\infty}(I_n; H_0^1(\Omega))$ into $\mathbb{P}^k \otimes H_0^1(\Omega)$, it follows that every
 362 term in (3.25) is in $C(I_n; H_0^1(\Omega))$. This implies $d_u^n \in C(I_n; H_0^1(\Omega))$.

363 By using the triangle inequality, from (3.25) we get

$$364 \quad (3.30) \quad \max_{t \in I_n} \|d_u^n\|_{H^1} \leq \max_{t \in I_n} (\|\partial_t R_\tau^n (R_h u - u)\|_{H^1} + \|\Delta_h R_h (u - R_\tau^n u)\|_{H^1})$$

$$365 \quad + \max_{t \in I_n} (\|rg(u)u - I_\tau^n [rg(u)u]\|_{H^1} + \|rg(u)u - r_h^* g(u_h^*) u_h^*\|_{H^1})$$

$$366 \quad =: D_1^u + D_2^u + D_3^u + D_4^u.$$

368 Choosing $m = 0$ in Lemma 3.3, we obtain the following stability result:

$$369 \quad (3.31) \quad \|R_\tau^n u\|_{W^{1,\infty}(I_n; H^s)} \leq C \|u\|_{W^{1,\infty}(I_n; H^s)}.$$

371 Using (3.31) and (3.24), we can estimate D_1^u as follows:

$$372 \quad D_1^u = \max_{t \in I_n} \|\partial_t R_\tau^n (R_h u - u)\|_{H^1} \leq \|R_h u - u\|_{W^{1,\infty}(I_n; H^1)}$$

$$373 \quad \leq Ch^p \|R_h u - u\|_{W^{1,\infty}(I_n; H^{p+1})}.$$

375 Similarly, using identity (3.23) and Lemma 3.3, we have

$$376 \quad D_2^u = \max_{t \in I_n} \|\Delta_h R_h (u - R_\tau^n u)\|_{H^1} \leq \max_{t \in I_n} \|u - R_\tau^n u\|_{H^3}$$

$$377 \quad \leq C\tau^{k+1} \|u\|_{W^{k+1,\infty}(I_n; H^3)},$$

379 and

$$380 \quad D_3^u = \max_{t \in I_n} \|rg(u)u - I_\tau^n [rg(u)u]\|_{H^1} \leq C\tau^{k+1}.$$

382 By using the triangle inequality, we decompose D_4^u into two parts,

$$383 \quad D_4^u \leq \max_{t \in I_n} (\|rg(u)u - rg(R_h u) R_h u\|_{H^1} + \|rg(R_h u) R_h u - R_\tau^n rg(R_\tau^n R_h u) R_\tau^n R_h u\|_{H^1})$$

$$384 \quad \leq Ch^p + C\tau^{k+1}.$$

386 Then, substituting the estimates of D_j^u , $j = 1, 2, 3, 4$, into (3.30), we obtain the desired
 387 estimate for $\|d_u^n\|_{H^1}$.

388 To estimate $|P_\tau^n d_r^n|$, we rewrite (3.26) as

$$389 \quad d_r^n = \frac{1}{2} \operatorname{Re} \left[(g(u)u, \partial_t (u - u_h^*)) + (g(u)u - g(u_h^*) u_h^*, \partial_t u_h^*) \right]$$

$$390 \quad + \frac{1}{2} \operatorname{Re} \left[(g(u_h^*) u_h^*, \partial_t u_h^*) - I_\tau^n (g(u_h^*) u_h^*, \partial_t u_h^*) \right]$$

391

392 and test this expression by $P_\tau^n v$ in the time interval I_n , with $v \in \mathbb{P}^k$. This yields

$$\begin{aligned}
393 \quad (3.32) \quad & \int_{I_n} P_\tau^n d_\tau^n v \, dt = \int_{I_n} d_\tau^n P_\tau^n v \, dt \\
394 \quad & \leq \frac{1}{2} \operatorname{Re} \int_{I_n} (g(u)u, \partial_t(u - u_h^*)) P_\tau^n v \, dt \\
395 \quad & \quad + C\tau^{\frac{1}{2}} \|(g(u)u - g(u_h^*)u_h^*, \partial_t u_h^*)\|_{L^\infty(I_n)} \|v\|_{L^2(I_n)} \\
396 \quad & \quad + C\tau^{k+\frac{3}{2}} \|\partial_t^{k+1}(g(u_h^*)u_h^*, \partial_t u_h^*)\|_{L^\infty(I_n)} \|v\|_{L^2(I_n)} \\
397 \quad & \leq \frac{1}{2} \operatorname{Re} \int_{I_n} (g(u)u, \partial_t(u - u_h^*)) P_\tau^n v \, dt + C\tau^{\frac{1}{2}}(h^p + \tau^{k+1}) \|v\|_{L^2(I_n)}. \\
398 \quad &
\end{aligned}$$

399 The first term on the right-hand side of (3.32) can be estimated as follows.

$$\begin{aligned}
400 \quad (3.33) \quad & \frac{1}{2} \operatorname{Re} \int_{I_n} (g(u)u, \partial_t(u - u_h^*)) P_\tau^n v \, dt = \int_{I_n} (g(u)u, \partial_t(u - R_\tau^n u)) P_\tau^n v \, dt \\
401 \quad & \quad + \int_{I_n} (g(u)u, \partial_t R_\tau^n(u - R_h u)) P_\tau^n v \, dt \\
402 \quad & =: D_1^r + D_2^r, \\
403 \quad &
\end{aligned}$$

$$\begin{aligned}
405 \quad D_1^r &= \int_{I_n} (g(u)u P_\tau^n v, \partial_t(u - R_\tau^n u)) \, dt \\
406 \quad &= \int_{I_n} (g(u)u P_\tau^n v - P_\tau^n(g(u)u P_\tau^n v), \partial_t(u - R_\tau^n u)) \, dt \\
407 \quad &\leq C\tau^{\frac{1}{2}} \|g(u)u P_\tau^n v - P_\tau^n(g(u)u P_\tau^n v)\|_{L^2(I_n; L^2)} \|\partial_t(u - R_\tau^n u)\|_{L^\infty(I_n; L^2)} \\
408 \quad &\leq C\tau^{\frac{3}{2}} \|P_\tau^n v\|_{L^2(I_n; L^2)} \|\partial_t(u - R_\tau^n u)\|_{L^\infty(I_n; L^2)} \quad (\text{we have used Lemma 3.4}) \\
409 \quad &\leq C\tau^{k+\frac{3}{2}} \|v\|_{L^2(I_n)} \|\partial_t^{k+1} u\|_{L^\infty(I_n; L^2)} \quad (\text{we have used Lemma 3.3}),
\end{aligned}$$

$$\begin{aligned}
410 \quad D_2^r &\leq C\tau^{\frac{1}{2}} \|g(u)u\|_{L^\infty(I_n; L^2)} \|u - R_h u\|_{W^{1, \infty}(I_n; L^2)} \|v\|_{L^2(I_n)} \\
411 \quad &\leq C\tau^{\frac{1}{2}} h^p \|v\|_{L^2(I_n)} \|u\|_{W^{1, \infty}(I_n; H^{p+1})}.
\end{aligned}$$

413 Substituting these estimates into (3.32), we obtain

$$\left| \int_{I_n} P_\tau^n d_\tau^n v \, dt \right| \leq C\tau^{\frac{1}{2}}(h^p + \tau^{k+1}) \|v\|_{L^2(I_n)}.$$

416 Since this inequality holds for arbitrary $v \in L^2(I_n)$, it follows that

$$\|P_\tau^n d_\tau^n\|_{L^2(I_n)} \leq C\tau^{\frac{1}{2}}(h^p + \tau^{k+1}).$$

419 Then, using the inverse inequality in time, we obtain the desired estimate for $|P_\tau^n d_\tau^n|$. \square

420 **4. Well-posedness and convergence analysis.** We define the error functions $e_h^u =$
421 $u_h - u_h^*$ and $e_h^r = r_h - r_h^*$, with the following abbreviations:

$$\begin{aligned}
422 \quad & e_{nj}^u = e_h^u(t_{nj}), \quad e_{nj}^r = e_h^r(t_{nj}), \quad u_{nj} = u_h(t_{nj}), \quad r_{nj} = r_h(t_{nj}), \\
423 \quad & u_{nj}^* = u_h^*(t_{nj}), \quad r_{nj}^* = r_h^*(t_{nj}), \quad v_{nj} = v_h(t_{nj}), \quad q_{nj} = q_h(t_{nj}).
\end{aligned}$$

424 Subtracting (3.27)–(3.28) from (3.7)–(3.8), we obtain the following error equations:

$$425 \quad i \int_{I_n} (\partial_t e_h^u, v_h) dt = - \int_{I_n} (\nabla P_\tau^n e_h^u, \nabla v_h) dt + \frac{\tau}{2} \sum_{j=1}^k w_j (e_{n_j}^r g(u_{n_j}) u_{n_j}, v_{n_j})$$

$$426 \quad (4.1a) \quad + \frac{\tau}{2} \sum_{j=1}^k w_j (r_{n_j}^* [g(u_{n_j}) u_{n_j} - g(u_{n_j}^*) u_{n_j}^*], v_{n_j}) - \int_{I_n} (P_\tau^n d_u^n, v_h) dt,$$

$$427 \quad \int_{I_n} \partial_t e_h^r q_h dt = \frac{\tau}{4} \sum_{j=1}^k w_j \operatorname{Re}(q_{n_j} (g(u_{n_j}) u_{n_j} - g(u_{n_j}^*) u_{n_j}^*), \partial_t u_h^*(t_{n_j}))$$

$$428 \quad (4.1b) \quad + \frac{\tau}{4} \sum_{j=1}^k w_j \operatorname{Re}(q_{n_j} g(u_{n_j}) u_{n_j}, \partial_t e_h^u(t_{n_j})) - \int_{I_n} P_\tau^n d_r^n q_h dt,$$

$$429$$

430 which hold for all test functions $v_h \in \mathbb{P}^k \otimes S_h$ and $q_h \in \mathbb{P}^k$.

431 **REMARK 4.1.** If (4.1) has a solution $(e_h^u, e_h^r) \in X_{\tau,h} \times Y_{\tau,h}$, then $u_h = u_h^* + e_h^u$ and
 432 $r_h = r_h^* + e_h^r$ give a solution of the numerical scheme (2.8). In the following, we prove
 433 existence of (e_h^u, e_h^r) to (4.1) by using Schaefer's Fixed Point Theorem, which is quoted
 434 below.

435 **THEOREM 4.1** (Schaefer's Fixed Point Theorem [11, Chapter 9.2, Theorem 4]). *Let B*
 436 *be a Banach space and $M : B \rightarrow B$ be a continuous and compact mapping. If the set*

$$437 \quad (4.2) \quad \{\phi \in B : \exists \theta \in [0, 1] \text{ such that } \phi = \theta M(\phi)\}$$

438 *is bounded in B , then the mapping M has at least one fixed point.*

439 We define

$$440 \quad (4.3) \quad X_{\tau,h}^* = \left\{ v_h \in X_{\tau,h} : \max_{1 \leq n \leq N} \max_{1 \leq j \leq k} \|v_h(t_{n_j}) - u_h^*(t_{n_j})\|_{L^\infty \cap H^1} \leq \frac{1}{2} \right\},$$

$$441 \quad (4.4) \quad Y_{\tau,h}^* = \left\{ q_h \in Y_{\tau,h} : \max_{1 \leq n \leq N} \max_{1 \leq j \leq k} |q_h(t_{n_j}) - r_h^*(t_{n_j})| \leq \frac{1}{2} \right\},$$

$$442$$

where the norm $\|\cdot\|_{L^\infty \cap H^1}$ is defined as

$$\|\phi_h\|_{L^\infty \cap H^1} := \max(\|\phi_h\|_{L^\infty}, \|\phi_h\|_{H^1}).$$

443 For any element $(\phi_h, \varphi_h) \in X_{\tau,h} \times Y_{\tau,h}$, we define two associated numbers

$$444 \quad (4.5a) \quad \rho[\phi_h] := \min \left(\frac{1}{\max_{1 \leq n \leq N} \max_{1 \leq j \leq k} \|\phi_h(t_{n_j})\|_{L^\infty \cap H^1}}, 1 \right),$$

$$445 \quad (4.5b) \quad \rho[\varphi_h] := \min \left(\frac{1}{\max_{1 \leq n \leq N} \max_{1 \leq j \leq k} |\varphi_h(t_{n_j})|}, 1 \right),$$

$$446$$

447 which are continuous with respect to (ϕ_h, φ_h) (because all norms are equivalent in the finite-
 448 dimensional space $X_{\tau,h} \times Y_{\tau,h}$). Furthermore, the two numbers defined above satisfy the
 449 following estimates:

$$450 \quad (4.6) \quad \max_{1 \leq n \leq N} \max_{1 \leq j \leq k} \|\rho[\phi_h] \phi_h(t_{n_j})\|_{L^\infty \cap H^1} \leq 1,$$

$$451 \quad (4.7) \quad \max_{1 \leq n \leq N} \max_{1 \leq j \leq k} |\rho[\varphi_h] \varphi_h(t_{n_j})| \leq 1.$$

$$452$$

453 Then we define

$$454 \quad (4.8) \quad u^\phi := u_h^* + \rho[\phi_h] \phi_h \quad \text{and} \quad r^\varphi := r_h^* + \rho[\varphi_h] \varphi_h,$$

$$455$$

456 with the following abbreviations:

$$457 \quad u_{nj}^\phi = u_h^\phi(t_{nj}) \quad \text{and} \quad \varphi_{nj} = \varphi_h(t_{nj}),$$

458 and define $(e_h^u, e_h^r) \in X_{\tau,h} \times Y_{\tau,h}$ to be the solution of the following linear equations:

$$460 \quad (4.9) \quad i \int_{I_n} (\partial_t e_h^u, v_h) dt + \int_{I_n} (\nabla P_\tau^n e_h^u, \nabla v_h) dt = \frac{\tau}{2} \sum_{j=1}^k w_j (\varphi_{nj} g(u_{nj}^\phi) u_{nj}^\phi, v_{nj}) \\ 461 \quad + \frac{\tau}{2} \sum_{j=1}^k w_j (r_{nj}^* [g(u_{nj}^\phi) u_{nj}^\phi - g(u_{nj}^*) u_{nj}^*], v_{nj}) - \int_{I_n} (P_\tau^n d^u, v_h) dt$$

462 and

$$464 \quad (4.10) \quad \int_{I_n} \partial_t e_h^r q_h dt = \frac{\tau}{4} \sum_{j=1}^k w_j \operatorname{Re}(q_{nj} (g(u_{nj}^\phi) u_{nj}^\phi - g(u_{nj}^*) u_{nj}^*), \partial_t u_h^*(t_{nj})) \\ 465 \quad + \frac{\tau}{4} \sum_{j=1}^k w_j \operatorname{Re}(q_{nj} g(u_{nj}^\phi) u_{nj}^\phi, \partial_t \phi_h(t_{nj})) - \int_{I_n} P_\tau^n d^r q_h dt$$

466 for all $v_h \in \mathbb{P}^k \otimes S_h$ and $q_h \in \mathbb{P}^k$, $n = 1, \dots, N$. We denote by $M : X_{\tau,h} \times Y_{\tau,h} \rightarrow X_{\tau,h} \times Y_{\tau,h}$

468 the mapping from (ϕ_h, φ_h) to (e_h^u, e_h^r) , and define the set

$$469 \quad (4.11) \quad \mathfrak{B} = \{(\phi_h, \varphi_h) \in X_{\tau,h} \times Y_{\tau,h} : \exists \theta \in [0, 1] \text{ such that } (\phi_h, \varphi_h) = \theta M(\phi_h, \varphi_h)\},$$

470 and the following norm on $X_{\tau,h} \times Y_{\tau,h}$: for any $(\phi_h, \varphi_h) \in X_{\tau,h} \times Y_{\tau,h}$

$$471 \quad (4.12) \quad \|(\phi_h, \varphi_h)\|_{X_{\tau,h} \times Y_{\tau,h}} := \|\phi_h\|_{L^\infty(0,T;H^1)} + \|\varphi_h\|_{L^\infty(0,T)}.$$

472 It is straightforward to show the following result (see [12, Proof of Lemma 4.2]).

473 LEMMA 4.2. *The mapping $M : X_{\tau,h} \times Y_{\tau,h} \rightarrow X_{\tau,h} \times Y_{\tau,h}$ is well defined, continuous*
474 *and compact.*

475 Moreover, there holds the following key technical lemma.

476 LEMMA 4.3. *Let $1 \leq d \leq 3$ and assume that the solution of the NLS equation (1.1) is*
477 *sufficiently smooth. Then there exist positive constants τ_0 and h_0 such that when $\tau \leq \tau_0$ and*
478 *$h \leq h_0$, the following statement holds: If $(\phi_h, \varphi_h) \in \mathfrak{B}$ and $(e_h^u, e_h^r) = M(\phi_h, \varphi_h)$, then*

$$479 \quad (4.13) \quad \|e_h^u\|_{L^\infty(0,T;H^1)} + \|e_h^r\|_{L^\infty(0,T)} \leq C(\|e_h^u(0)\|_{H^1} + |e_h^r(0)|) \\ 480 \quad + C \max_{1 \leq n \leq N} \max_{t \in I_n} (\|d_u^n\|_{H^1} + |P_\tau^n d_r^n|),$$

$$481 \quad (4.14) \quad \max_{1 \leq n \leq N} \max_{1 \leq j \leq k} \|e_h^u(t_{nj})\|_{L^\infty \cap H^1} \leq \frac{1}{2} \quad \text{and} \quad \max_{1 \leq n \leq N} \max_{1 \leq j \leq k} |e_h^r(t_{nj})| \leq \frac{1}{2},$$

$$482 \quad (4.15) \quad \rho[\phi_h] = 1, \quad \rho[\varphi_h] = 1.$$

484 *Proof.* Since the proof is very long and technical, below we only outline the main steps
485 and ingredients of the proof and refer the interested reader to [12] for the details.

If $(\phi_h, \varphi_h) \in \mathfrak{B}$ and $(e_h^u, e_h^r) = M(\phi_h, \varphi_h)$, then

$$(\phi_h, \varphi_h) = \theta M(\phi_h, \varphi_h) = (\theta e_h^u, \theta e_h^r),$$

486 which implies $\phi_h = \theta e_h^u$ and $\varphi_h = \theta e_h^r$. In this case, (4.9)–(4.10) can be rewritten as

$$487 \quad (4.16) \quad i \int_{I_n} (\partial_t e_h^u, v_h) dt = - \int_{I_n} (\nabla P_\tau^n e_h^u, \nabla v_h) dt + \frac{\theta\tau}{2} \sum_{j=1}^k w_j (e_{n_j}^r g(u_{n_j}^\phi) u_{n_j}^\phi, v_{n_j})$$

$$488 \quad + \frac{\tau}{2} \sum_{j=1}^k w_j (r_{n_j}^* [g(u_{n_j}^\phi) u_{n_j}^\phi - g(u_{n_j}^*) u_{n_j}^*], v_{n_j}) - \int_{I_n} (P_\tau^n d_u^n, v_h) dt,$$

$$489 \quad (4.17) \quad \int_{I_n} \partial_t e_h^r q_h dt = \frac{\tau}{4} \sum_{j=1}^k w_j \operatorname{Re}(q_{n_j} (g(u_{n_j}^\phi) u_{n_j}^\phi - g(u_{n_j}^*) u_{n_j}^*), \partial_t u_h^*(t_{n_j}))$$

$$490 \quad + \frac{\theta\tau}{4} \sum_{j=1}^k w_j \operatorname{Re}(q_{n_j} g(u_{n_j}^\phi) u_{n_j}^\phi, \partial_t e_h^u(t_{n_j})) - \int_{I_n} P_\tau^n d_r^n q_h dt,$$

$$491$$

492 which hold for all $v_h \in \mathbb{P}^k \otimes S_h$ and $q_h \in \mathbb{P}^k$, $n = 1, \dots, N$. It remains to derive estimates
493 for e_h^u and e_h^r based on the above equations.

494 From (4.6)–(4.7) and definition (4.8) we get

$$495 \quad (4.18) \quad \max_{1 \leq n \leq N} \max_{1 \leq j \leq k} \|u^\phi(t_{n_j})\|_{L^\infty \cap H^1} + \max_{1 \leq n \leq N} \max_{1 \leq j \leq k} |r^\varphi(t_{n_j})|$$

$$496 \quad \leq \max_{1 \leq n \leq N} \max_{1 \leq j \leq k} \|u_h^*(t_{n_j})\|_{L^\infty \cap H^1} + \max_{1 \leq n \leq N} \max_{1 \leq j \leq k} |r_h^*(t_{n_j})|$$

$$497 \quad + \max_{1 \leq n \leq N} \max_{1 \leq j \leq k} \|\rho[\phi_h] \phi_h(t_{n_j})\|_{L^\infty \cap H^1} + \max_{1 \leq n \leq N} \max_{1 \leq j \leq k} |\rho[\varphi_h] \varphi_h(t_{n_j})|$$

$$498 \quad \leq \|u_h^*\|_{L^\infty(0, T; L^\infty \cap H^1)} + \|r_h^*\|_{L^\infty(0, T)} + 2.$$

499 Thus $\|u^\phi(t_{n_j})\|_{L^\infty \cap H^1}$ and $|r^\varphi(t_{n_j})|$ are bounded uniformly with respect to τ and h .

500 The major part of the remaining proof is devoted to proving the following three inequal-
501 ities:
502

$$503 \quad (4.19) \quad \int_{I_n} \|e_h^u\|_{H^1}^2 dt \leq C\tau \|e_h^u(t_{n-1})\|_{H^1}^2 + C\tau^2 \int_{I_n} |e_h^r|^2 dt + C\tau^3 \max_{t \in I_n} \|d_u^n\|_{H^1}^2.$$

$$504 \quad (4.20) \quad \int_{I_n} |e_h^r|^2 dt \leq C\tau [\|e_h^u(t_{n-1})\|_{H^1}^2 + |e_h^r(t_{n-1})|^2] + \tau^2 \max_{t \in I_n} (\|d_u^n\|_{H^1}^2 + |P_\tau^n d_r^n|^2).$$

$$505 \quad (4.21) \quad \|\nabla e_h^u(t_n)\|^2 + |e_h^r(t_n)|^2 - \|\nabla e_h^u(t_{n-1})\|^2 - |e_h^r(t_{n-1})|^2 + \int_{I_n} \|\partial_t e_h^u\|_{H^{-1}}^2 dt$$

$$506 \quad \leq C \int_{I_n} (\|e_h^u\|_{H^1}^2 + |e_h^r|^2) dt + C \int_{I_n} (\|d_u^n\|_{H^1}^2 + |P_\tau^n d_r^n|^2) dt.$$

$$507$$

508 In particular, (4.19) can be obtained by substituting $v_h = (-\Delta_h) P_\tau^n [P_\tau^n e_h^u(t)(t_n - t)]$ into
509 (4.16) and considering the imaginary part; (4.20) can be obtained by substituting $q_h =$
510 $P_\tau^n [P_\tau^n e_h^r(t)(t_n - t)]$ into (4.17); (4.21) is obtained by setting $v_h = \partial_t e_h^u$ in (4.16) and
511 considering the real part, setting $q_h = 2e_h^r$ in (4.17), and estimating $\int_{I_n} \|\partial_t e_h^u\|_{H^{-1}}^2 dt$ via a
512 duality argument using (4.16). More details can be found in [12, Proof of Lemma 4.3].

513 To complete the proof, substituting (4.19)–(4.20) into (4.21), we obtain

$$514 \quad (4.22) \quad (\|\nabla e_h^u(t_n)\|^2 + |e_h^r(t_n)|^2) - (\|\nabla e_h^u(t_{n-1})\|^2 + |e_h^r(t_{n-1})|^2) + \int_{I_n} \|\partial_t e_h^u\|_{H^{-1}}^2 dt$$

$$515 \quad \leq C\tau (\|\nabla e_h^u(t_{n-1})\|^2 + |e_h^r(t_{n-1})|^2) + C \int_{I_n} (\|d_u^n\|_{H^1}^2 + |P_\tau^n d_r^n|^2) dt.$$

$$516$$

517 It follows from Gronwall's inequality that

$$518 \quad (4.23) \quad \max_{1 \leq n \leq N} (\|\nabla e_h^u(t_n)\|^2 + |e_h^r(t_n)|^2) + C \int_0^T \|\partial_t e_h^u\|_{H^{-1}}^2 dt$$

$$519 \quad \leq C(\|\nabla e_h^u(0)\|^2 + |e_h^r(0)|^2) + C \sum_{n=1}^N \int_{I_n} (\|d_u^n\|_{H^1}^2 + |P_\tau^n d_r^n|^2) dt.$$

521 Then, substituting the above inequality into (4.19)–(4.20) and using temporal inverse in-
522 equality, we obtain

$$523 \quad (4.24) \quad \max_{t \in [0, T]} (\|e_h^u(t)\|_{H^1}^2 + |e_h^r(t)|^2) \leq C(\|e_h^u(0)\|_{H^1}^2 + |e_h^r(0)|^2)$$

$$524 \quad + C \max_{1 \leq n \leq N} \max_{t \in I_n} (\|d_u^n\|_{H^1}^2 + |P_\tau^n d_r^n|^2).$$

526 Hence, (4.13) holds.

527 When τ and h are sufficiently small, inequality (4.24) implies that

$$528 \quad (4.25) \quad \max_{t \in [0, T]} \|e_h^u(t)\|_{H^1} \leq \frac{1}{2} \quad \text{and} \quad \max_{t \in [0, T]} |e_h^r(t)| \leq \frac{1}{2}.$$

530 On the one hand, by the inverse inequality, we have

$$531 \quad (4.26) \quad \max_{t \in [0, T]} \|e_h^u(t)\|_{L^\infty} \leq C \ell_h \max_{t \in [0, T]} \|e_h^u(t)\|_{H^1}$$

$$532 \quad \leq C \ell_h \left[\|e_h^u(0)\|_{H^1} + |e_h^r(0)| + \max_{1 \leq n \leq N} \max_{t \in I_n} (\|d_u^n\|_{H^1} + |P_\tau^n d_r^n|) \right],$$

533 where

$$\ell_h = \begin{cases} 1 & \text{if } d = 1, \\ \ln(2 + 1/h) & \text{if } d = 2, \\ h^{-\frac{1}{2}} & \text{if } d = 3. \end{cases}$$

534 On the other hand, by choosing a test function v in (4.16) satisfying the properties $v(t_{nj}) = 1$
535 and $v(t_{ni}) = 0$ for $i \neq j$, and using property (3.6), we obtain

$$536 \quad (4.27) \quad \|\Delta_h e_{nj}^u\| = \left\| i \partial_t e_{nj}^u - \theta P_h [e_{nj}^r g(u_{nj}^\phi) u_{nj}^\phi] + P_h d_{nj}^u \right.$$

$$537 \quad \left. - P_h [r_{nj}^* (g(u_{nj}^\phi) u_{nj}^\phi - g(u_{nj}^*) u_{nj}^*)] \right\|$$

$$538 \quad \leq C \tau^{-1} \left[\|e_h^u(0)\|_{H^1} + |e_h^r(0)| + \max_{1 \leq n \leq N} \max_{t \in I_n} (\|d_u^n\|_{H^1} + |P_\tau^n d_r^n|) \right],$$

540 where we have used (4.23)–(4.24) and an inverse inequality in time in estimating $\partial_t e_{nj}^u$. By
541 the discrete Sobolev embedding inequality, for $1 \leq d \leq 3$ we have

$$542 \quad (4.28) \quad \|e_{nj}^u\|_{L^\infty} \leq C \|e_{nj}^u\|_{H^1}^{\frac{1}{2}} \|\Delta_h e_{nj}^u\|_{H^1}^{\frac{1}{2}}$$

$$543 \quad \leq C \tau^{-\frac{1}{2}} \left[\|e_h^u(0)\|_{H^1} + |e_h^r(0)| + \max_{1 \leq n \leq N} \max_{t \in I_n} (\|d_u^n\|_{H^1} + |P_\tau^n d_r^n|) \right],$$

545 where we have used (4.24) and (4.27) in the last inequality. Then, combining (4.26) and
546 (4.28) yields

$$547 \quad \max_{1 \leq n \leq N} \max_{1 \leq j \leq k} \|e^u(t_{nj})\|_{L^\infty} \leq C \min(\ell_h, \tau^{-\frac{1}{2}}) \left[\|e_h^u(0)\|_{H^1} + |e_h^r(0)| \right.$$

$$548 \quad \left. + \max_{1 \leq n \leq N} \max_{t \in I_n} (\|d_u^n\|_{H^1} + |P_\tau^n d_r^n|) \right]$$

$$549 \quad \leq C (h^{p-\frac{1}{2}} + \tau^{k+\frac{1}{2}}),$$

551 where we have used the consistency estimate from Theorem 3.5. When τ and h are suffi-

552 ciently small, the inequality above implies

$$553 \quad (4.29) \quad \max_{1 \leq n \leq N} \max_{1 \leq j \leq k} \|e^u(t_{nj})\|_{L^\infty} \leq \frac{1}{2}.$$

554 This together with (4.25) gives (4.14).

555 Furthermore, since $\phi_h = \theta e_h^u$ and $\varphi_h = \theta e_h^r$, it follows that

$$\max_{1 \leq n \leq N} \max_{1 \leq j \leq k} \|\phi_h(t_{nj})\|_{L^\infty \cap H^1} \leq \frac{1}{2} \quad \text{and} \quad \max_{1 \leq n \leq N} \max_{1 \leq j \leq k} |\varphi_h(t_{nj})| \leq \frac{1}{2},$$

556 which imply $\rho[\phi_h] = \rho[\varphi_h] = 1$ in view of the definition in (4.5). This proves (4.15). \square

557 We now are ready to state and prove existence, uniqueness and convergence of numerical
558 solutions, which comprise of the second main theorem of this paper.

559 **THEOREM 4.4.** *Let $1 \leq d \leq 3$ and assume that the solution of the NLS equation (1.1) is*
560 *sufficiently smooth. Then there exist positive constants τ_0 and h_0 such that when $\tau \leq \tau_0$ and*
561 *$h \leq h_0$, the numerical method (2.8) has a unique solution $(u_h, r_h) \in X_{\tau,h}^* \times Y_{\tau,h}^*$. Moreover,*
562 *this solution satisfies the following error estimate:*

$$563 \quad (4.30) \quad \max_{t \in [0, T]} \left(\|u_h(t) - u_h^*(t)\|_{H^1} + |r_h(t) - r_h^*(t)| \right) \leq C(h^p + \tau^{k+1}).$$

564 *Proof. Step 1: Existence.* By the definition of \mathfrak{B} , if $(\phi_h, \varphi_h) \in \mathfrak{B}$ and $(e_h^u, e_h^r) =$
565 $M(\phi_h, \varphi_h)$ then $\phi_h = \theta e_h^u$ and $\varphi_h = \theta e_h^r$. Thus (4.13) implies

$$566 \quad (4.31) \quad \|(\phi_h, \varphi_h)\|_{X_{\tau,h} \times Y_{\tau,h}} = \|\phi_h\|_{L^\infty(0, T; H^1)} + \|\varphi_h\|_{L^\infty(0, T)} \leq C,$$

567 which together with Schaefer's fixed point theorem imply the existence of a fixed point
(ϕ_h, φ_h) for the mapping M (corresponding to $\theta = 1$), with

$$(e_h^u, e_h^r) = (\phi_h, \varphi_h), \quad u^\phi = u_h^* + \phi_h \quad \text{and} \quad r^\phi = r_h^* + \varphi_h,$$

568 satisfying (4.9)–(4.10), where we have used (4.15) in the expression (4.8). Consequently,
569 (e_h^u, e_h^r) is a solution of (4.1) with $(u_h, r_h) = (u_h^\phi, r_h^\phi) = (u_h^* + e_h^u, r_h^* + e_h^r)$. Hence, in view
570 of the discussions in Remark 4.1, (u_h, r_h) is a solution of the numerical scheme (2.8), and
571 (4.14) implies (u_h, r_h) is in the set $X_{\tau,h}^* \times Y_{\tau,h}^*$ defined in (4.3)–(4.4). This proves existence
572 of a numerical solution in $X_{\tau,h}^* \times Y_{\tau,h}^*$.

573 *Step 2: Uniqueness.* Suppose that (u_h, r_h) and $(\tilde{u}_h, \tilde{r}_h)$ in $X_{\tau,h}^* \times Y_{\tau,h}^*$ are two pairs
574 of numerical solutions, and set $e_h^u = u_h - \tilde{u}_h$ and $e_h^r = r_h - \tilde{r}_h$ (abusing the notation).
575 Subtracting the corresponding equations satisfied by (u_h, r_h) and $(\tilde{u}_h, \tilde{r}_h)$ shows that (e_h^u, e_h^r)
576 satisfies equations (4.1) with $e_h^u(0) = e_h^r(0) = 0$ and $d_u^n = d_r^n = 0$. In the meantime, the
577 definition in (4.3)–(4.4) implies

$$578 \quad (4.32) \quad \|e_h^u(t_{nj})\|_{L^\infty \cap H^1} \leq 1 \quad \text{and} \quad |e_h^r(t_{nj})| \leq 1.$$

580 Accordingly, (e_h^u, e_h^r) is a fixed point of the mapping M (corresponding to $\theta = 1$ in \mathfrak{B}) in
581 the case $e_h^u(0) = e_h^r(0) = 0$ and $d_u^n = d_r^n = 0$. Hence, an application of (4.13) yields

$$582 \quad \|e_h^u\|_{L^\infty(0, T; H^1)} + \|e_h^r\|_{L^\infty(0, T)} \leq C \left[\|e_h^u(0)\|_{H^1} + |e_h^r(0)| \right. \\ \left. + \max_{1 \leq n \leq N} \max_{t \in I_n} (\|d_u^n\|_{H^1} + |P_\tau^n d_r^n|) \right] = 0.$$

585 Thus, $(u_h, r_h) = (\tilde{u}_h, \tilde{r}_h)$ and the uniqueness of the numerical solution is proved.

586 *Step 3: Error estimate.* Since the error functions $e_h^u = u_h - u_h^*$ and $e_h^r = r_h - r_h^*$ satisfy
587 (4.1) and (4.32), it follows that (e_h^u, e_h^r) is a fixed point of the mapping M (corresponding
588 to $\theta = 1$ in \mathfrak{B}). Hence, an application of (4.13) yields

$$589 \quad \|e_h^u\|_{L^\infty(0, T; H^1)} + \|e_h^r\|_{L^\infty(0, T)} \leq C \left[\|e_h^u(0)\|_{H^1} + |e_h^r(0)| + \max_{1 \leq n \leq N} \max_{t \in I_n} (\|d_u^n\|_{H^1} + |P_\tau^n d_r^n|) \right].$$

591 Substituting the consistency error estimates from Theorem 3.5 into the above inequality

592 yields the desired estimate (4.30). The proof is complete. □

593 **REMARK 4.2.** For the periodic and Neumann boundary conditions, the mass and energy
594 conservations in Theorem 3.1 and the error estimate in Theorem 4.4 can be proved similarly.

595 **5. Numerical experiments.** In this section, we present some one-dimensional nu-
596 merical tests to validate the theoretical results proved in Theorems 3.1 and 4.4 about the
597 mass and energy conservations, and the convergence rates of the proposed method. All the
598 computations are performed using the software package FEniCS (<https://fenicsproject.org>).
599 We consider the cubic nonlinear Schrödinger equation

$$600 \quad (5.1) \quad \begin{aligned} i\partial_t u - \partial_{xx} u - 2|u|^2 u &= 0 && \text{in } (-L, L) \times (0, T], \\ u|_{t=0} &= u_0 && \text{in } (-L, L), \quad \text{with } L = 20, \end{aligned}$$

602 subject to the periodic boundary condition. We choose $u_0 = \text{sech}(x) \exp(2ix)$ so that the
603 exact solution is given by

$$604 \quad (5.2) \quad u(x, t) = \text{sech}(x + 4t) \exp(i(2x + 3t)).$$

606 This example contains a soliton wave and is often used as a benchmark for measuring the
607 effectiveness of numerical methods for the NLS equation; see [34, 38, 26].

608 **5.1. Convergence rates.** We solve problem (5.1) by the proposed method (2.8) and
609 compare the numerical solutions with the exact solution (5.2). Newton’s method is used to
610 solve the nonlinear system. The iteration is set to stop when the error is below 10^{-10} .

611 The time discretization errors are presented in Table 1, where we have used finite ele-
612 ments of degree 3 with a sufficiently spatial mesh $h = 2L/5000$ so that the error from spatial
613 discretization is negligibly small in observing the temporal convergence rates. From Table
614 1 we see that the error of time discretization is $O(\tau^{k+1})$, which is consistent with the result
615 proved in Theorem 4.4.

616 The spatial discretization errors are presented in Table 2, where we have chosen $k = 3$
617 with a sufficiently small time stepsize $\tau = 1/1000$ so that the time discretization error
618 is negligibly small compared to the spatial error. From Table 2 we see that the spatial
619 discretization errors are $O(h^p)$ in the H^1 norm. This is also consistent with the result
620 proved in Theorem 4.4.

621 **5.2. Mass and energy conservations.** We denote the mass and SAV energy of a
622 numerical solution by

$$623 \quad (5.3) \quad M_h(t) = \int_{\Omega} |u_h(t)|^2 dx \quad \text{and} \quad E_h(t) = \frac{1}{2} \int_{\Omega} |\nabla u_h(t)|^2 dx - r_h(t)^2,$$

624 respectively. The evolution of mass and SAV energy of the numerical solutions is presented
in Figure 1 with $\tau = 0.2$ and $h = 0.2$. It is shown that

$$\text{mass} = 2 + O(10^{-12}) \quad \text{and} \quad \text{SAV energy} = -7.33358048516 + O(10^{-12}),$$

625 which are much smaller than the error of the numerical solutions, as shown in Figure 2. This
626 shows the effectiveness of the proposed method in preserving mass and energy (independent
627 of the error of numerical solutions). The number of iterations at each time level is presented
628 in Figure 3 to show the effectiveness of the Newton’s method.

629 **5.3. Comparison of different methods in preserving the shape of a soliton.**

630 The graph of $|u(x, t)|$ is a soliton propagating towards left. Its shape remains unchanged for
631 all $t \geq 0$. The graphs of numerical solutions given by several different numerical methods
632 using the same mesh sizes are presented in Figures 4 and 5. All the methods preserve mass
633 and energy conservations. The numerical results show the effectiveness of the proposed
634 method in preserving the shape of the soliton.

TABLE 1
Time discretization errors of the proposed method, with $h = \frac{2L}{5000}$ and $T = 1$.

k	τ	$p = 3$	
		$\ u(x, t) - u_h(x, t)\ _{L^\infty(0, T; H^1)}$	order
2	1/60	3.7964E-05	–
	1/70	2.3429E-05	3.1312
	1/80	1.5460E-05	3.1132
	1/90	1.0733E-05	3.0985
	1/100	7.7542E-06	3.0853
3	1/20	3.4019E-05	–
	1/25	1.3821E-05	4.0364
	1/30	6.6322E-06	4.0275
	1/35	3.5689E-06	4.0200
	1/40	2.0886E-06	4.0123
4	1/8	1.2291E-04	–
	1/12	1.5120E-05	5.1681
	1/14	6.8492E-06	5.1369
	1/16	3.4634E-06	5.1067
	1/20	1.1555E-06	4.9192

TABLE 2
Spatial discretization errors of the proposed method, with $\tau = \frac{1}{1000}$ and $T = 1$.

p	M	$k = 3$	
		$\ u(x, t) - u_h(x, t)\ _{L^\infty(0, T; H^1)}$	order
1	1400	5.8670E-02	–
	1600	5.1134E-02	1.0295
	1800	4.5330E-02	1.0229
	2000	4.0719E-02	1.0183
	2200	3.6964E-02	1.0149
2	240	1.9306E-02	–
	260	1.6438E-02	2.0094
	280	1.4167E-02	2.0062
	300	1.2338E-02	2.0041
	320	1.0842E-02	2.0027
3	90	1.6147E-02	–
	100	1.1661E-02	3.0894
	110	8.7112E-03	3.0599
	120	6.6844E-03	3.0436
	130	5.2435E-03	3.0334

635 **5.4. Capability of solving focusing nonlinearity.** We consider the cubic nonlinear
636 Schrödinger equation

$$637 \quad (5.4) \quad \begin{aligned} i\partial_t u - \partial_{xx} u - \partial_{yy} u + 2|u|^2 u &= 0 && \text{in } \Omega \times (0, T], \\ u|_{t=0} &= u_0 && \text{in } \Omega, \end{aligned}$$

639 in two-dimensional space $\Omega = [0, 1] \times [0, 1]$ subject to the periodic boundary condition. We
640 choose $u_0 = \exp(2\pi i(x + y))$ so that the exact solution is given by

$$641 \quad (5.5) \quad u(x, t) = \exp(i(2\pi x + 2\pi y + (2 + 8\pi^2)t)),$$

643 which admits a progressive plane wave solution; see [38].

644 We solve problem (5.4) by the proposed method (2.8) and compare the numerical solu-




FIG. 1. Evolution of mass $M_h(t) - M_h(0)$ and SAV energy $E_h(t) - E_h(0)$, with $p = 3$ and $\tau = h = 0.2$.




FIG. 2. Evolution of error of the numerical solution, with $p = 3$ and $\tau = h = 0.2$.




FIG. 3. Number of iterations at each time level, with $p = 3$ and $\tau = h = 0.2$.

645 tions with the exact solution (5.5). Newton’s method is used to solve the nonlinear system.
 646 The iteration is stopped when the error is below 10^{-10} .

647 The time discretization errors are presented in Table 3, where we have used finite ele-
 648 ments of degree 3 with a sufficiently spatial mesh $h = 1/80$ so that the error from spatial
 649 discretization is negligibly small in observing the temporal convergence rates. From Table
 650 3 we see that the error of time discretization is $O(\tau^{k+1})$, which is consistent with the result
 651 proved in Theorem 4.4.

652 The spatial discretization errors are presented in Table 4, where we have chosen $k = 3$
 653 with a sufficiently small time stepsize $\tau = 1/1000$ so that the time discretization error
 654 is negligibly small compared to the spatial error. From Table 4 we see that the spatial
 655 discretization errors are $O(h^p)$ in the H^1 norm. This is also consistent with the result
 656 proved in Theorem 4.4.

The evolution of mass and SAV energy of the numerical solutions is presented in Figure




FIG. 4. Soliton propagation when $t \in [0, 2]$: numerical solutions with $p = 1$, $M = 1200$ and $\Delta t = 0.1$.




FIG. 5. Soliton propagation when $t \in [0, 2]$: numerical solutions with $p = 1$, $M = 1200$ and $\Delta t = 0.05$.

TABLE 3
Time discretization errors of the proposed method, with $h = \frac{1}{80}$ and $T = 0.1$.

k	τ	$p = 3$	
		$\ u(x, t) - u_h(x, t)\ _{L^\infty(0, T; H^1)}$	order
2	1/460	5.0023E-04	–
	1/480	4.3780E-04	3.1321
	1/500	3.8572E-04	3.1027
	1/520	3.4198E-04	3.0686
	1/540	3.0504E-04	3.0290
3	1/60	1.6206E-02	–
	1/80	4.9792E-03	4.1022
	1/100	2.0173E-03	4.0490
	1/120	9.6960E-04	4.0183
	1/140	5.2530E-04	3.9761
4	1/30	3.6941E-02	–
	1/40	8.0993E-03	5.2750
	1/50	2.5534E-03	5.1731
	1/60	1.0078E-03	5.0989
	1/70	4.6554E-04	5.0104

TABLE 4
Spatial discretization errors of the proposed method, with $\tau = \frac{1}{1000}$ and $T = 0.1$.

p	h	$k = 3$	
		$\ u(x, t) - u_h(x, t)\ _{L^\infty(0, T; H^1)}$	order
1	1/70	5.6297E-01	–
	1/80	4.8304E-01	1.1466
	1/90	4.2346E-01	1.1178
	1/100	3.7726E-01	1.0964
	1/110	3.4035E-01	1.0803
2	1/10	4.9467E-01	–
	1/15	2.0992E-01	2.1141
	1/20	1.1748E-01	2.0178
	1/25	7.5177E-02	2.0005
	1/30	5.2233E-02	1.9972
3	1/12	2.1955E-02	–
	1/14	1.3738E-02	3.0412
	1/16	9.1747E-03	3.0236
	1/18	6.4327E-03	3.0144
	1/20	4.6849E-03	3.0092

6 with $\tau = 0.01$ and $h = 0.1$. It is shown that

$$\text{mass} = 1.000397142598 + O(10^{-12}) \quad \text{and} \quad \text{SAV energy} = 80.45628698537 + O(10^{-11}),$$

657 which are much smaller than the error of the numerical solutions, as shown in Figure 7. This
658 shows the effectiveness of the proposed method in preserving mass and energy (independent
659 of the error of numerical solutions). The number of iterations at each time level is presented
660 in Figure 8 to show the effectiveness of the Newton's method.

661

REFERENCES

662 [1] G. D. Akrivis, V. A. Dougalis and O. A. Karakashian, *On fully discrete Galerkin methods of second-*




FIG. 6. Evolution of mass $M_h(t) - M_h(0)$ and SAV energy $E_h(t) - E_h(0)$, with $p = 3$, $\tau = 0.01$ and $h = 0.1$.




FIG. 7. Evolution of error of the numerical solution, with $p = 3$, $\tau = 0.01$ and $h = 0.1$.




FIG. 8. Number of iterations at each time level, with $p = 3$, $\tau = 0.01$ and $h = 0.1$.

- 663 order temporal accuracy for the NLS equation, *Numer. Math.*, 59(1991), pp. 31–53. pages 2
 664 [2] G. Akrivis, *Finite difference discretization of the cubic Schrödinger equation*, *IMA J. Numer. Anal.*
 665 13(1993), pp. 115–124. pages 2
 666 [3] G. Akrivis, B. Li and D. Li, *Energy-decaying extrapolated RK-SAV methods for the Allen–Cahn and*
 667 *Cahn–Hilliard equations*, *SIAM J. Sci. Comput.*, 41(2019), pp. A3703–A3727. pages 2
 668 [4] X. Antoine, W. Bao and C. Besse, *Computational methods for the dynamics of the nonlinear*
 669 *Schrödinger/Gross–Pitaevskii equations*, *Comput. Phys. Commun.*, 184(2013), pp. 2621–2633.
 670 pages 2
 671 [5] W. Bao and Y. Cai, *Optimal error estimates of finite difference methods for the Gross–Pitaevskii*
 672 *equation with angular momentum rotation*, *Math. Comp.*, 82(2013), pp. 99–128. pages 2
 673 [6] W. Bao, Q. Tang and Z. Xu, *Numerical methods and comparison for computing dark and bright*
 674 *solitons in the NLS equation*, *J. Comput. Phys.*, 235(2013), pp. 423–445. pages 2
 675 [7] J. Bourgain, *Global Solutions of Nonlinear Schrödinger Equations*, vol. 46, American Mathematical
 676 Society, 1999. pages 1
 677 [8] S. C. Brenner and L.R. Scott, *The Mathematical Theory of FEMs*, Third edition. Texts in Applied
 678 Mathematics, Vol. 15, Springer, New York, 2008. pages 4, 9
 679 [9] C. Canuto, M. Y. Hussaini, A. Quarteroni and T. A. Zang, *Spectral Methods: Fundamentals in Single*

- 680 *Domains*, Springer, Berlin, 2007. pages 6, 8
- 681 [10] M. Delfour, M. Fortin and G. Payre, *Finite-difference solutions of a non-linear Schrödinger equation*,
682 J. Comput. Phys., 44(1981), pp.277–288. pages 2
- 683 [11] L. C. Evans, *Partial Differential Equations*, second edition, Graduate Studies in Mathematics 19,
684 AMS, Providence, RI, 2010. pages 3, 12
- 685 [12] X. Feng, B. Li and S. Ma, *High-order mass- and energy-conserving SAV-Gauss collocation finite*
686 *element methods for the nonlinear Schrödinger equation*, arXiv:2006.05073 pages 8, 13, 14
- 687 [13] X. Feng, H. Liu and S. Ma, *Mass- and energy-conserved numerical schemes for nonlinear Schrödinger*
688 *equations*, Commun. Comput. Phys., 26(2019), pp. 1365–1396. pages 2
- 689 [14] Z. Gao and S. Xie, *Fourth-order alternating direction implicit compact finite difference schemes for*
690 *two-dimensional Schrödinger equations*, Appl. Numer. Math., 61(2011), pp. 593–614. pages 2
- 691 [15] D. Gilbarg and N. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer, New
692 York, 2001. pages
- 693 [16] G. H. Golub and J. H. Welsch, *Calculation of Gauss quadrature rules*, Math. Comput., 23(1969), pp.
694 221–230. pages 6
- 695 [17] Y. Gong, Q. Wang and Y. Wang, *A conservative Fourier pseudo-spectral method for the nonlinear*
696 *Schrödinger equation*, J. Comput. Phys., 328(2017), pp. 354–370. pages 2
- 697 [18] Y. Gong and J. Zhao, *Energy-stable Runge–Kutta schemes for gradient flow models using the energy*
698 *quadrization approach*, Appl. Math. Letters, 94(2019), pp. 224–231. pages 2
- 699 [19] Y. Gong, J. Zhao and Q. Wang, *Arbitrarily high-order unconditionally energy stable schemes for*
700 *thermodynamically consistent gradient flow models*, SIAM J. Sci. Comput. 42(2020), pp. B135–
701 156. pages 2
- 702 [20] P. Henning and D. Peterseim, *Crank–Nicolson Galerkin approximations to nonlinear Schrödinger*
703 *equations with rough potentials*, Math. Models Meth. Appl. Sci., 27(2017), pp. 2147–2184. pages 2
- 704 [21] J. Hong, Y. Liu, H. Munthe-Kaas and A. Zanna, *Globally conservative properties and error esti-*
705 *mation of a multi-symplectic scheme for Schrödinger equations with variable coefficients*, Appl.
706 Numer.Math., 56(2006), pp. 814–843. pages 2
- 707 [22] O. Karakashian and C. Makridakis, *A space-time finite element method for the nonlinear Schrödinger*
708 *equation: the discontinuous Galerkin method*, Math. Comp., 67(1998), pp. 479–499. pages 2
- 709 [23] O. Karakashian and C. Makridakis, *A space-time finite element method for the nonlinear Schrödinger*
710 *equation: the continuous Galerkin method*, SIAM J. Numer. Anal., 36(1999), pp. 1779–1807. pages
711 2
- 712 [24] D. A. Kopriva, *Implementing spectral methods for partial differential equations: Algorithms for scien-*
713 *tists and engineers*, Springer Science & Business Media, 2009. pages 6
- 714 [25] H. Liu, Y. Huang, W. Lu and N. Yi, *On accuracy of the mass-preserving DG method to multi-*
715 *dimensional Schrödinger equations*, IMA J. Numer. Anal., 39(2019), pp. 760–791 pages 2
- 716 [26] W. Lu, and Y. Huang and H. Liu, *Mass preserving discontinuous Galerkin methods for Schrödinger*
717 *equations*, J. Comput. Phys., 282(2015), pp. 210–226. pages 17
- 718 [27] D. E. Pelinovsky, V. V. Afanasjev and Y. S. Kivshar, *Nonlinear theory of oscillating, decaying, and*
719 *collapsing solitons in the generalized nonlinear Schrödinger equation*, Phys. Rev. E, 53(1996), pp.
720 1940–1953. pages 1
- 721 [28] J. M. Sanz-Serna, *Methods for the numerical solution of the nonlinear Schrödinger equation*, Math.
722 Comp., 43(1984), pp. 21–27 pages 2
- 723 [29] H. W. Schürmann, *Traveling-wave solutions of the cubic-quintic nonlinear Schrödinger equation*, Phys.
724 Rev. E, 54 (1996), pp. 4312–4320. pages 1
- 725 [30] J. Shen, J. Xu and J. Yang, *A new class of efficient and robust energy stable schemes for gradient*
726 *flows*, SIAM Rev., 61(2019), pp. 474–506. pages 2, 3
- 727 [31] J. Shen, J. Xu and J. Yang, *The scalar auxiliary variable (SAV) approach for gradient flows*, J.
728 Comput. Phys., 353(2018), pp. 407–416. pages 2
- 729 [32] J. Shen, T. Tang and L. L. Wang, *Spectral methods: algorithms, analysis and applications*, Springer
730 Science & Business Media, 41(2011). pages 4
- 731 [33] W. A. Strauss and L. Vazquez, *Numerical solution of a nonlinear Klein–Gordon equation*, J. Comput.
732 Phys., 28(1978), pp.271–278. pages 2
- 733 [34] N. Taghizadeh, M. Mirzazadeh and F. Farahrooz, *Exact solutions of the nonlinear Schrödinger equation*
734 *by the first integral method*, J. Math. Anal. Appl., 374(2011), pp. 549–553. pages 17
- 735 [35] T. Tao, *Nonlinear Dispersive Equations: Local and Global Analysis*, American Mathematical Society,
736 2006. pages 1
- 737 [36] J. Wang, *A new error analysis of Crank–Nicolson Galerkin FEMs for a generalized nonlinear*
738 *Schrödinger equation*, J. Sci. Comput., 60(2014), pp. 390–407. pages 2
- 739 [37] T. Wang, B. Guo and Q. Xu, *Fourth-order compact and energy conservative difference schemes for*
740 *the nonlinear Schrödinger equation in two dimensions*, J. Comput. Phys., 243(2013), pp. 382–399.
741 pages 2
- 742 [38] Y. Xu and C. W. Shu, *Local discontinuous Galerkin methods for nonlinear Schrödinger equations*, J.
743 Comput. Phys., 205(2005), pp. 72–97. pages 2, 17, 18
- 744 [39] X. Yang, *Linear, first and second-order, unconditionally energy stable numerical schemes for the phase*
745 *field model of homopolymer blends*, J. Comput. Phys. 327(2016), pp. 294–316. pages 2

- 746 [40] X. Yang and L. Ju, *Efficient linear schemes with unconditional energy stability for the phase field*
747 *elastic bending energy model*, *Comput. Meth. Appl. Mech. Engrg.*, 315(2017), pp. 691–712. pages
748 2
- 749 [41] X. Yang and L. Ju, *Linear and unconditionally energy stable schemes for the binary fluid-surfactant*
750 *phase field model*, *Comput. Meth. Appl. Mech. Engrg.*, 318(2017), pp. 1005–1029. pages 2
- 751 [42] X. Yang, J. Zhao, Q. Wang and J. Shen, *Numerical approximations for a three components Cahn–*
752 *Hilliard phase-field model based on the invariant energy quadratization method*, *Math. Models*
753 *Methods Appl. Sci.*, 27(2017), pp. 1993–2030. pages 2
- 754 [43] N. J. Zabusky and M. D. Kruskal, *Interaction of "solitons" in a collisionless plasma and the recurrence*
755 *of initial states*, *Phys. Rev. Lett.*, 15(1965), pp. 240–243. pages 1