

# Global dynamics and spatio-temporal patterns of predator–prey systems with density-dependent motion

HAI-YANG JIN<sup>1</sup> and ZHI-AN WANG<sup>2</sup>

<sup>1</sup>*Department of Mathematics, South China University of Technology, Guangzhou 510640, China  
email: mahyjin@scut.edu.cn*

<sup>2</sup>*Department of Applied Mathematics, Hong Kong Polytechnic University, Hung Hom, Hong Kong  
email: mawza@polyu.edu.hk*

(Received 8 August 2018; revised 19 June 2019; accepted 6 July 2020; first published online 12 August 2020)

In this paper, we investigate the global boundedness, asymptotic stability and pattern formation of predator–prey systems with density-dependent prey-taxis in a two-dimensional bounded domain with Neumann boundary conditions, where the coefficients of motility (diffusion) and mobility (prey-taxis) of the predator are correlated through a prey density-dependent motility function. We establish the existence of classical solutions with uniform-in time bound and the global stability of the spatially homogeneous prey-only steady states and coexistence steady states under certain conditions on parameters by constructing Lyapunov functionals. With numerical simulations, we further demonstrate that spatially homogeneous time-periodic patterns, stationary spatially inhomogeneous patterns and chaotic spatio-temporal patterns are all possible for the parameters outside the stability regime. We also find from numerical simulations that the temporal dynamics between linearised system and nonlinear systems are quite different, and the prey density-dependent motility function can trigger the pattern formation.

**Key words:** Predator–prey system, prey-taxis, boundedness, global stability, Lyapunov functional

**2020 Mathematics Subject Classification:** Primary: 35K51; Secondary: 35A01, 35B40, 35K57, 35Q92

## 1 Introduction

The foraging is the searching for wild food resources by hunting, fishing, consuming or the gathering of plant matter. It plays an important role in an organism's ability to survive and reproduce. The non-random foraging strategies in the predator–prey dynamics, such as the area-restricted search, are often observed to result in populations of predators moving (or flowing) towards regions of higher prey density (see Refs. [7, 30, 31]). Such movement is referred to as prey-taxis which has important roles in biological control or ecological balance such as regulating prey (pest) population to avoid incipient outbreaks of prey or forming large-scale aggregation for survival (cf. Refs. [10, 30, 34]). To understand the dynamics of predator–prey systems with prey-taxis, Kareiva and Odell [16] put individual foraging behaviours into a biased random walk

model which, upon passage to a continuum limit, leads to the following prey-taxis system (see equations (55) and (56) in Ref. [16]):

$$\begin{cases} \partial_t u = \nabla \cdot (d(v)\nabla u) - \nabla \cdot (u\chi(v)\nabla v) + \mathcal{F}(u, v), \\ \partial_t v = D\Delta v + \mathcal{G}(u, v), \end{cases} \quad (1.1)$$

where  $u = u(x, t)$  and  $v = v(x, t)$  denote the population density of predators and preys at position  $x$  and time  $t$ , respectively, and  $D > 0$  is a constant denoting the diffusivity of preys. The term  $\nabla \cdot (d(v)\nabla u)$  describes the diffusion (motility) of predators with coefficient  $d(v)$ , and  $-\nabla \cdot (u\chi(v)\nabla v)$  accounts for the prey-taxis (mobility) with coefficient  $\chi(v)$ , where both motility and mobility coefficients are related to individual foraging behaviours. The source terms  $\mathcal{F}(u, v)$  and  $\mathcal{G}(u, v)$  represent the predator–prey interactions to be discussed below. By fitting the abstract model (1.1) to field experiment data of area-restricted search behaviour exhibited by individual ladybugs (predators) and aphids (prey) with appropriate predator–prey interactions (see Ref. [16]), Karevia and Odell showed that the area-restricted non-random foraging yield heterogeneous aggregative patterns observed in the field experiment.

In a special case  $\chi(v) = -d'(v)$ , the system (1.1) becomes

$$\begin{cases} \partial_t u = \Delta(d(v)u) + \mathcal{F}(u, v), \\ \partial_t v = D\Delta v + \mathcal{G}(u, v), \end{cases}$$

where the diffusion term  $\Delta(d(v)u)$  with  $d'(v) < 0$  has been interpreted as ‘density-suppressed motility’ in Refs. [9, 24] (see mathematical results in Refs. [13, 26, 35]), and  $d(v)$  is called the motility function. This means that the predator will reduce its motility when encountering the prey, which is a rather reasonable assumption and has very sound applications in the predator–prey systems.

As mentioned in Ref. [16], the model (1.1) was tailored to study the non-random foraging behaviour (or prey-taxis) not only for ladybugs and aphids but also for general organisms living in the predator–prey system. Formally, the model (1.1) can be regarded as a variant of the Keller–Segel chemotaxis model [17], where  $u(x, t)$  denotes the cell density and  $v(x, t)$  the chemical concentration. However the prey-taxis model (1.1) has two striking features that the Keller–Segel models have not considered yet. First the model (1.1) characterises the non-random population dispersal and aggregation (i.e. both diffusion and prey-taxis coefficients depend on the prey density). Second, the source terms in (1.1) have the inter-specific interactions. These two features distinguish the prey-taxis model from the Keller–Segel type chemotaxis models.

Ecological/biological interactions can be defined as either intra-specific or inter-specific. The former occurs between individuals of the same species, while the latter between two or more species. There are three types of basic interspecific interactions (see Refs. [8, 14, 23]): predator–prey, competition and mutualism, which can be encapsulated in  $\mathcal{F}(u, v)$  and  $\mathcal{G}(u, v)$  by the following typical form:

$$\mathcal{F}(u, v) = c_1 u F(v) + h(u), \quad \mathcal{G}(u, v) = f(v) - c_2 u F(v), \quad (1.2)$$

where  $h(u)$  and  $f(v)$  are functions representing the intra-specific interactions of predators and preys, respectively. Parameters  $c_1, c_2 \in \mathbb{R}$  denote the coefficients of inter-specific interactions between predators and preys, where  $F(v)$  is commonly called the functional response function fulfilling  $F(0) = 0$ ,  $F'(v) > 0$ . This paper is interested in the predator–prey interaction where

$c_1 > 0, c_2 > 0$ . The function forms given in (1.2) have represented most of ecologically meaningful examples of predator–prey population interactions used in the literature by assigning appropriate expressions to  $h(u), f(v)$  and  $F(v)$ . Typically, the predator kinetics  $h(u)$  may include density-dependent death  $h(u) = -u(\theta + \alpha u), \theta > 0, \alpha \geq 0$ , the prey kinetics  $f(v)$  could be linear, logistic or Allee effect (bistable) type and  $F(v)$  may be of Lotka–Volterra type [25, 41] or Holling’s type [11]. We refer the readers to the excellent surveys [29, 40] for an exhaustive list of  $h(u), f(v)$  and  $F(v)$ . Hence, in this paper, we consider the following system:

$$\begin{cases} u_t = \nabla \cdot (d(v)\nabla u) - \nabla \cdot (u\chi(v)\nabla v) + \gamma uF(v) - \theta u - \alpha u^2, & x \in \Omega, \quad t > 0, \\ v_t = D\Delta v - uF(v) + f(v), & x \in \Omega, \quad t > 0, \\ \frac{\partial u}{\partial \nu} = \frac{\partial v}{\partial \nu} = 0, & x \in \partial\Omega, \quad t > 0, \\ u(x, 0) = u_0(x), v(x, 0) = v_0(x), & x \in \Omega, \end{cases} \tag{1.3}$$

where  $D, \gamma > 0, \theta > 0, \alpha \geq 0$  are constants, and  $d(v), \chi(v), F(v)$  and  $f(v)$  satisfy the following conditions:

- (H1)  $d(v), \chi(v) \in C^2([0, \infty)), d(v) > 0, \chi(v) \geq 0$  and  $d'(v) \leq 0$  on  $[0, \infty)$ .
- (H2)  $F(v) \in C^1([0, \infty)), F(0) = 0, F(v) > 0$  in  $(0, \infty)$  and  $F'(v) > 0$ .
- (H3)  $f : [0, \infty) \rightarrow \mathbb{R}$  is in  $C^1$  with  $f(0) = 0$ , and there exist two constants  $\mu, K > 0$  such that  $f(v) \leq \mu v$  for any  $v \geq 0, f(K) = 0$  and  $f(v) < 0$  for all  $v > K$ .

We remark that the above assumptions for  $F(v)$  and  $f(v)$  have covered a large class of interesting and meaningful examples encountered in the literature as mentioned above. Our first result on the global boundedness of solutions of (1.3) is the following.

**Theorem 1.1** (Global boundedness) *Assume  $(u_0, v_0) \in [W^{1,p}(\Omega)]^2$  with  $p > 2$  and  $u_0, v_0 \geq 0 (\neq 0)$ . Let  $\Omega \subset \mathbb{R}^2$  be a bounded domain with smooth boundary and the hypotheses (H1)–(H3) hold. If  $\alpha > 0$  or  $\chi(v) = -d'(v)$ , then there is a unique classical solution  $(u, v) \in [C([0, \infty) \times \bar{\Omega}) \cap C^{2,1}((0, \infty) \times \bar{\Omega})]^2$  solving the problem (1.3). Moreover there is constant  $C > 0$  independent of  $t$  such that*

$$\|u(\cdot, t)\|_{L^\infty} + \|v(\cdot, t)\|_{W^{1,\infty}} \leq C,$$

where in particular  $0 \leq v \leq K_0$  with

$$K_0 := \max\{\|v_0\|_{L^\infty}, K\}. \tag{1.4}$$

Next, we will study the large time behaviour of solutions. One can easily compute that the system (1.3) has three homogeneous steady states  $(u_s, v_s)$ :

$$(u_s, v_s) = \begin{cases} (0, 0) \text{ or } (0, K), & \text{if } \gamma F(K) \leq \theta, \\ (0, 0) \text{ or } (0, K) \text{ or } (u_*, v_*), & \text{if } \gamma F(K) > \theta \end{cases}$$

with  $u_*, v_* > 0$  determined by the following algebraic equations:

$$u_* = \frac{f(v_*)}{F(v_*)}, \quad \gamma F(v_*) = \theta + \alpha u_*, \tag{1.5}$$

where  $(0, 0)$  is the extinction steady state,  $(0, K)$  is the prey-only steady state and  $(u_*, v_*)$  is the coexistence steady state. As in Ref. [12], for the global stability, along with the hypotheses (H1)–(H3), we need another condition for the following compound function:

$$\phi(v) = \frac{f(v)}{F(v)},$$

as follows:

(H4) The function  $\phi(v)$  is continuously differentiable on  $(0, \infty)$ ,  $\phi(0) = \lim_{v \rightarrow 0} \phi(v) > 0$  and  $\phi'(v) < 0$  for any  $v \geq 0$ .

Then, the global stability results are given as follows:

**Theorem 1.2** (Global stability) *Let the hypotheses (H1)–(H4) and assumptions in Theorem 1.1 hold. Then, the solution  $(u, v)$  obtained in Theorem 1.1 has the following properties:*

(1) *If the parameters  $\theta, \gamma, K$  satisfy  $\gamma F(K) \leq \theta$  where ‘=’ holds iff  $\alpha > 0$ , then*

$$\|u\|_{L^\infty} + \|v - K\|_{L^\infty} \rightarrow 0 \text{ as } t \rightarrow \infty,$$

*exponentially if  $\gamma F(K) < \theta$  or algebraically if  $\gamma F(K) = \theta, \alpha > 0$ .*

(2) *If the parameters  $\theta, \gamma, K$  satisfy  $\gamma F(K) > \theta$  and*

$$D \geq \max_{0 \leq v \leq K_0} \frac{u_* |F(v)|^2 |\chi(v)|^2}{4\gamma F(v_*) F'(v) d(v)},$$

*then*

$$\|u - u_*\|_{L^\infty} + \|v - v_*\|_{L^\infty} \rightarrow 0 \text{ as } t \rightarrow \infty,$$

*where the convergence is exponential if  $\alpha > 0$ .*

We remark if  $d(v)$  and  $\chi(v)$  are constant, the system (1.3) has been studied from various aspects (cf. Refs. [2, 6, 22, 23, 37, 46]), and in particular the global existence and stability of solutions have been established in a previous work by the authors in Ref. [12]. The results Theorems 1.1 and 1.2 extend the results of Ref. [12] to non-constant  $d(v)$  and  $\chi(v)$ . Moreover, the proof of Theorem 1.2 uses a similar idea (method of Lyapunov functional) as in Ref. [12]. Except these analogies, we would like to stress some essential differences between the present paper and [12] below.

- The method used in Ref. [12] to prove the global existence of solutions was based on a *a priori* estimate for the energy functional  $\int_\Omega u \ln u dx$  to attain the  $L^2$ -estimate of solutions. However, such *a priori* estimates are attainable only for case where the motility function  $d(v)$  is constant. Hence, the method of Ref. [12] is inapplicable to the model (1.3). In this paper, we estimate  $L^2$ -norm of solutions directly to obtain the global existence of solutions (see Section 3). With this new direct  $L^2$ -estimate method, the concavity of  $F(v)$  required in Ref. [12] is no longer needed. That is, we not only use a different method to prove the global boundedness of solutions but also remove the condition  $F''(v) \leq 0$  imposed in Ref. [12].
- If  $d(v)$  and  $\chi(v)$  are non-constant as considered in this paper, we find that the system (1.3) can generate pattern formation as shown in Section 5. This is different from Ref. [12] where

no pattern formation can be founded for constant  $d(v)$  and  $\chi(v)$ . Our result indicates that the density-dependent motility is a trigger for pattern formation, which is a new finding.

- In the proof of global stability shown in Section 4, we need the estimate of  $\|\nabla u\|_{L^4}$  which can be easily obtained for constant  $d(v)$  and  $\chi(v)$  but is not clear if  $d(v)$  and  $\chi(v)$  are not constant. Hence, we present a new proof in Lemma 4.2.

In Section 5, we shall detail the results of Theorems 1.1 and 1.2 for specific and often used forms of  $F(v)$ . In Theorem 1.2, the conditions on parameter values are identified to ensure the global stability of homogeneous steady states. However, it is unknown whether non-homogeneous (i.e. non-constant) steady states exist outside the stability regimes found in Theorem 1.2. In the final Section 5, we shall use linear stability analysis to find the conditions on parameters for the instability of equilibria of (1.3) and then perform numerical simulations to illustrate that indeed spatially inhomogeneous patterns and time-periodic patterns can be found under certain conditions. We also demonstrate that the non-constant motility function  $d(v)$  plays an important role in generating the pattern formation.

### 2 Local existence and Preliminaries

In what follows, we shall use  $c_i$  or  $C_i (i = 1, 2, 3, \dots)$  to denote a generic constant which may vary in the context. We first state the existence of local-in-time classical solutions of system (1.3) by using the abstract theory of quasilinear parabolic systems in Ref. [5].

**Lemma 2.1** (Local existence) *Let  $\Omega$  be a bounded domain in  $\mathbb{R}^2$  with smooth boundary and the hypotheses (H1)–(H3) hold. Assume  $(u_0, v_0) \in [W^{1,p}(\Omega)]^2$  with  $u_0, v_0 \geq 0 (\neq 0)$  and  $p > 2$ . Then there exists  $T_{max} \in (0, \infty]$  such that the problem (1.3) has a unique classical solution  $(u, v) \in [C([0, T_{max}) \times \bar{\Omega}) \cap C^{2,1}((0, T_{max}) \times \bar{\Omega})]^2$  satisfying  $u, v \geq 0$  for all  $t > 0$ . Moreover, we have*

$$\text{either } T_{max} = \infty, \text{ or } \limsup_{t \nearrow T_{max}} (\|u(\cdot, t)\|_{L^\infty} + \|v(\cdot, t)\|_{L^\infty}) = \infty. \tag{2.1}$$

**Proof** We shall apply the theory developed by Amann [5] to prove this lemma. With  $\omega = (u, v)$ , the system (1.3) can be reformulated as

$$\begin{cases} \omega_t = \nabla \cdot (\mathbb{A}(\omega)\nabla\omega) + \Phi(\omega), & x \in \Omega, t > 0, \\ \frac{\partial \omega}{\partial \nu} = 0, & x \in \partial\Omega, t > 0, \\ \omega(\cdot, 0) = (u_0, v_0), & x \in \Omega, \end{cases}$$

where

$$\mathbb{A}(\omega) = \begin{bmatrix} d(v) & -\chi(v)u \\ 0 & 1 \end{bmatrix} \text{ and } \Phi(\omega) = \begin{bmatrix} \gamma u F(v) - \theta u - \alpha u^2 \\ f(v) - u F(v) \end{bmatrix}.$$

Since the given initial conditions satisfy  $0 \leq (u_0, v_0) \in [W^{1,p}(\Omega)]^2$  with  $p > 2$  and hence the matrix  $\mathbb{A}(\omega)$  is positively definite at  $t = 0$ . Hence, the system (1.3) is normally parabolic and the local existence of solutions follows from Ref. [4, Theorem 7.3]. That is, there exists a  $T_{max} > 0$  such that the system (1.3) admits a unique solution  $(u, v) \in [C([0, T_{max}) \times \bar{\Omega}) \cap C^{2,1}((0, T_{max}) \times \bar{\Omega})]^2$ .

Next, we will use the maximum principle to prove that  $u, v \geq 0$ . To this end, we rewrite the first equation of (1.3) as

$$u_t = d(v)\Delta u + [d'(v)\nabla v - \chi(v)\nabla v] \cdot \nabla u - \chi'(v)u|\nabla v|^2 - \chi(v)u\Delta v + \gamma uF(v) - \theta u - \alpha u^2. \tag{2.2}$$

Then applying the strong maximum principle to (2.2) with the Neumann boundary condition asserts that  $u > 0$  for all  $(x, t) \in \Omega \times (0, T_{max})$  due to  $u_0 \not\equiv 0$ . Similarly, we can show  $v > 0$  for any  $(x, t) \in \Omega \times (0, T_{max})$  by applying the strong maximum principle to the second equation of system (1.3). Since  $\mathbb{A}(\omega)$  is an upper triangular matrix, the assertion (2.1) follows from Ref. [3, Theorem 5.2] directly. Then the proof of Lemma 2.1 is completed.  $\square$

**Lemma 2.2** ([12]) *Under the conditions in Lemma 2.1, the solution  $(u, v)$  of (1.3) satisfies*

$$0 \leq v(x, t) \leq K_0, \text{ for all } x \in \Omega \text{ and } t \in (0, T_{max}), \tag{2.3}$$

where  $K_0$  is defined by (1.4).

Next, we present a basic boundedness property of the solutions to (1.3).

**Lemma 2.3** *Let  $(u, v)$  be a solution of system (1.3). Then there exists a constant  $C > 0$  such that*

$$\int_{\Omega} u dx \leq C, \text{ for all } t \in (0, T_{max}). \tag{2.4}$$

Moreover, if  $\alpha > 0$  or  $\chi(v) = -d'(v)$ , one has

$$\int_t^{t+\tau} \int_{\Omega} u^2 dx ds \leq C, \text{ for all } t \in (0, \tilde{T}_{max}), \tag{2.5}$$

where

$$\tau := \min \left\{ 1, \frac{1}{2} T_{max} \right\} \text{ and } \tilde{T}_{max} := \begin{cases} T_{max} - \tau, & \text{if } T_{max} < \infty, \\ \infty, & \text{if } T_{max} = \infty. \end{cases} \tag{2.6}$$

**Proof** Multiplying the second equation (1.3) by  $\gamma$  and adding the resulting equation into the first equation of (1.3), then integrating the result over  $\Omega \times (0, t)$ , one has

$$\frac{d}{dt} \left( \int_{\Omega} u dx + \gamma \int_{\Omega} v dx \right) + \int_{\Omega} u(\theta + \alpha u) dx = \gamma \int_{\Omega} f(v) dx,$$

which, along with the hypotheses (H2) and (H3) and the fact that  $0 \leq v \leq K_0$ , gives

$$\begin{aligned} \frac{d}{dt} \left( \int_{\Omega} u dx + \gamma \int_{\Omega} v dx \right) + \theta \left( \int_{\Omega} u dx + \gamma \int_{\Omega} v dx \right) + \alpha \int_{\Omega} u^2 dx &\leq (\gamma \mu + \theta \gamma) \int_{\Omega} v dx \\ &\leq (\gamma \mu + \theta \gamma) K_0 |\Omega|. \end{aligned} \tag{2.7}$$

Then, the Gronwall's inequality applied to (2.7) yields (2.4). If  $\alpha > 0$ , then integrating (2.7) over  $(t, t + \tau)$ , we have (2.5) directly.

Next, we consider the case  $\chi(v) = -d'(v)$ . In this case, the first equation of system (1.3) can be written as

$$u_t = \Delta(d(v)u) + \gamma uF(v) - \theta u - \alpha u^2. \tag{2.8}$$

Let  $0 < \delta < \frac{\theta}{d(0)}$  and let  $\mathcal{A}$  denote the self-adjoint realisation of  $-\Delta + \delta$  under homogeneous Neumann boundary conditions in  $L^2(\Omega)$ . Since  $\delta > 0$ ,  $\mathcal{A}$  possesses an order-preserving bounded inverse  $\mathcal{A}^{-1}$  on  $L^2(\Omega)$  and hence we can find a constant  $c_1 > 0$  such that

$$\|\mathcal{A}^{-1}\psi\|_{L^2} \leq c_1 \|\psi\|_{L^2} \quad \text{for all } \psi \in L^2(\Omega), \tag{2.9}$$

and

$$\|\mathcal{A}^{-\frac{1}{2}}\psi\|_{L^2}^2 = \int_{\Omega} \psi \cdot \mathcal{A}^{-1}\psi \, dx \leq c_1 \|\psi\|_{L^2}^2 \quad \text{for all } \psi \in L^2(\Omega). \tag{2.10}$$

From the second equation of (1.3) with (2.8), we have

$$(u + \gamma v)_t = \Delta(d(v)u + D\gamma v) - \theta u - \alpha u^2 + \gamma f(v),$$

which can be rewritten as

$$\begin{aligned} (u + \gamma v)_t + \mathcal{A}(d(v)u + D\gamma v) &= \delta[d(v)u + D\gamma v] - \theta u - \alpha u^2 + \gamma f(v) \\ &= (\delta d(v) - \theta - \alpha u)u + \gamma f(v) + D\delta\gamma v. \end{aligned} \tag{2.11}$$

Noting the facts  $0 < \delta < \frac{\theta}{d(0)}$  and (2.3), one can derive that

$$(\delta d(v) - \theta - \alpha u)u + \gamma f(v) + D\delta\gamma v \leq (\delta d(0) - \theta)u + c_2 \leq c_2. \tag{2.12}$$

Hence, multiplying (2.11) by  $\mathcal{A}^{-1}(u + \gamma v) \geq 0$ , and using the fact (2.12), one has

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} |\mathcal{A}^{-\frac{1}{2}}(u + \gamma v)|^2 \, dx + \int_{\Omega} (d(v)u + D\gamma v)(u + \gamma v) \, dx \leq c_2 \int_{\Omega} \mathcal{A}^{-1}(u + \gamma v) \, dx,$$

which together with the fact  $0 < d(K_0) \leq d(v)$ , gives a constant  $c_3 := \min\{d(K_0), D\}$  such that

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} |\mathcal{A}^{-\frac{1}{2}}(u + \gamma v)|^2 \, dx + c_3 \int_{\Omega} (u + \gamma v)^2 \, dx \leq c_2 \int_{\Omega} \mathcal{A}^{-1}(u + \gamma v) \, dx. \tag{2.13}$$

Using (2.9) and (2.10), we can derive that

$$\begin{aligned} \frac{c_3}{4c_1} \int_{\Omega} |\mathcal{A}^{-\frac{1}{2}}(u + \gamma v)|^2 \, dx + c_2 \int_{\Omega} \mathcal{A}^{-1}(u + \gamma v) \, dx &\leq \frac{c_3}{4} \int_{\Omega} (u + \gamma v)^2 \, dx + c_2 |\Omega|^{\frac{1}{2}} \|u + \gamma v\|_{L^2} \\ &\leq \frac{c_3}{2} \int_{\Omega} (u + \gamma v)^2 \, dx + \frac{c_2^2 |\Omega|}{c_3}. \end{aligned} \tag{2.14}$$

Adding (2.14) and (2.13), and letting  $y_1(t) := \int_{\Omega} |\mathcal{A}^{-\frac{1}{2}}(u + \gamma v)|^2$ , one has

$$y_1'(t) + \frac{c_3}{2c_1} y_1(t) + c_3 \int_{\Omega} (u + \gamma v)^2 \, dx \leq \frac{2c_2^2 |\Omega|}{c_3}.$$

Then one has  $y_1(t) \leq c_4$  and

$$\int_t^{t+\tau} \int_{\Omega} u^2 dx ds \leq \int_t^{t+\tau} \int_{\Omega} (u + \gamma v)^2 dx ds \leq \frac{y_1(t)}{c_3} + \frac{2c_2^2 |\Omega| \tau}{c_3^2} \leq \frac{c_4}{c_3} + \frac{2c_2^2 |\Omega|}{c_3^2} \text{ for all } t \in (0, \tilde{T}_{max}),$$

which gives (2.5). Then the proof of this lemma is completed. □

Moreover, we can thereupon deduce the following result as a consequence of Lemma 2.3.

**Lemma 2.4** *Let  $(u, v)$  be the solution of system (1.3), then there exists a constant  $C > 0$  independent of  $t$  such that*

$$\|\nabla v\|_{L^2} \leq C, \text{ for all } t \in (0, T_{max}), \tag{2.15}$$

and

$$\int_t^{t+\tau} \int_{\Omega} |\Delta v|^2 dx ds \leq C, \text{ for all } t \in (0, \tilde{T}_{max}), \tag{2.16}$$

where  $\tau$  and  $\tilde{T}_{max}$  are defined by (2.6).

**Proof** Multiplying the second equation of system (1.3) by  $-\Delta v$ , integrating the result by part and using the Cauchy–Schwarz inequality and the boundedness of  $v$  in (2.3), we end up with

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \int_{\Omega} |\nabla v|^2 dx + D \int_{\Omega} |\Delta v|^2 dx &= \int_{\Omega} u F(v) \Delta v dx - \int_{\Omega} f(v) \Delta v dx, \\ &\leq \frac{D}{2} \int_{\Omega} |\Delta v|^2 dx + \frac{1}{D} \int_{\Omega} F^2(v) u^2 dx + \frac{1}{D} \int_{\Omega} f^2(v) dx \\ &\leq \frac{D}{2} \int_{\Omega} |\Delta v|^2 dx + \frac{F^2(K_0)}{D} \int_{\Omega} u^2 dx + c_1, \end{aligned}$$

which yields

$$\frac{d}{dt} \int_{\Omega} |\nabla v|^2 dx + D \int_{\Omega} |\Delta v|^2 dx \leq \frac{2F^2(K_0)}{D} \int_{\Omega} u^2 dx + 2c_1. \tag{2.17}$$

Using the Gagliardo–Nirenberg inequality and noting the fact  $\|v\|_{L^2} \leq K_0 |\Omega|^{\frac{1}{2}}$ , one has

$$\int_{\Omega} |\nabla v|^2 dx = \|\nabla v\|_{L^2}^2 \leq c_2 (\|\Delta v\|_{L^2} \|v\|_{L^2} + \|v\|_{L^2}^2) \leq \frac{D}{2} \|\Delta v\|_{L^2}^2 + c_3. \tag{2.18}$$

Substituting (2.18) into (2.17), we get

$$\frac{d}{dt} \int_{\Omega} |\nabla v|^2 dx + \int_{\Omega} |\nabla v|^2 dx + \frac{D}{2} \int_{\Omega} |\Delta v|^2 dx \leq \frac{2F^2(K_0)}{D} \int_{\Omega} u^2 dx + c_4. \tag{2.19}$$



Denote  $y(t) := \int_{\Omega} |\nabla v(\cdot, t)|^2 dx$ ,  $t \in [0, T_{max})$  and  $z(t) := \frac{2F^2(K_0)}{D} \int_{\Omega} u^2 dx + c_4$ , from (2.19) we have

$$y'(t) + y(t) \leq z(t) \text{ for all } t \in (0, T_{max}), \tag{2.20}$$

which gives (2.15) by noting (2.5). On the other hand, integrating (2.19) over  $(t, t + \tau)$  along with (2.5) and (2.15), we obtain (2.16). □

### 3 Boundedness of solutions

We will derive the *a priori*  $L^2$ -estimate of the solution component  $u$ .

**Lemma 3.1** *Let the conditions in Theorem 1.1 hold. Then, the solution of (1.3) satisfies*

$$\|u(\cdot, t)\|_{L^2} \leq C, \text{ for all } t \in (0, T_{max}), \tag{3.1}$$

where  $C > 0$  is a constant independent of  $t$ .

**Proof** Multiplying the first equation of (1.3) by  $2u$ , integrating the result with respect to  $x$  over  $\Omega$ , one has

$$\begin{aligned} & \frac{d}{dt} \int_{\Omega} u^2 dx + 2 \int_{\Omega} d(v) |\nabla u|^2 dx + 2\alpha \int_{\Omega} u^3 dx + 2\theta \int_{\Omega} u^2 dx \\ & = 2 \int_{\Omega} u \chi(v) \nabla u \cdot \nabla v dx + 2\gamma \int_{\Omega} u^2 F(v) dx. \end{aligned} \tag{3.2}$$

With the assumptions in (H1)–(H2) and the fact (2.3), one has  $d(v) \geq d(K_0) > 0$ ,  $0 < F(v) \leq F(K_0)$  and  $0 \leq \chi(v) \leq c_1$ . Then using the Young’s inequality and Hölder inequality, we have from (3.2) that

$$\begin{aligned} & \frac{d}{dt} \int_{\Omega} u^2 dx + 2d(K_0) \int_{\Omega} |\nabla u|^2 dx + 2\alpha \int_{\Omega} u^3 dx + 2\theta \int_{\Omega} u^2 dx \\ & \leq 2c_1 \int_{\Omega} u |\nabla u| |\nabla v| dx + 2\gamma F(K_0) \int_{\Omega} u^2 dx \\ & \leq d(K_0) \int_{\Omega} |\nabla u|^2 dx + \frac{c_1^2}{d(K_0)} \int_{\Omega} u^2 |\nabla v|^2 dx + 2\gamma F(K_0) \int_{\Omega} u^2 dx \\ & \leq d(K_0) \|\nabla u\|_{L^2}^2 + \frac{c_1^2}{d(K_0)} \|u\|_{L^4}^2 \|\nabla v\|_{L^4}^2 + 2\gamma F(K_0) \|u\|_{L^2}^2. \end{aligned} \tag{3.3}$$

Using the Gagliardo–Nirenberg inequality, one has

$$\|u\|_{L^4}^2 \leq c_2 \left( \|\nabla u\|_{L^2} \|u\|_{L^2} + \|u\|_{L^2}^2 \right), \tag{3.4}$$

and the following estimate (cf. Ref. [13, Lemma 2.5]):

$$\|\nabla v\|_{L^4}^2 \leq c_3 \left( \|\Delta v\|_{L^2} \|\nabla v\|_{L^2} + \|\nabla v\|_{L^2}^2 \right) \leq c_3 c_4 \left( \|\Delta v\|_{L^2} + c_4 \right), \tag{3.5}$$

where the fact  $\|\nabla v\|_{L^2} \leq c_4$  has been used. Combining (3.4)–(3.5) and using the Young’s inequality, we obtain

$$\begin{aligned} \frac{c_1^2}{d(K_0)} \|u\|_{L^4}^2 \|\nabla v\|_{L^4}^2 &\leq c_5 (\|\nabla u\|_{L^2} \|u\|_{L^2} + \|u\|_{L^2}^2) (\|\Delta v\|_{L^2} + c_4) \\ &\leq c_5 \|\nabla u\|_{L^2} \|u\|_{L^2} \|\Delta v\|_{L^2} + c_4 c_5 \|\nabla u\|_{L^2} \|u\|_{L^2} \\ &\quad + c_5 \|u\|_{L^2}^2 \|\Delta v\|_{L^2} + c_4 c_5 \|u\|_{L^2}^2 \\ &\leq d(K_0) \|\nabla u\|_{L^2}^2 + \frac{c_5^2}{d(K_0)} \|u\|_{L^2}^2 \|\Delta v\|_{L^2}^2 + c_6 \|u\|_{L^2}^2, \end{aligned} \tag{3.6}$$

where  $c_5 := \frac{c_1^2 c_2 c_3 c_4}{d(K_0)}$  and  $c_6 := \frac{c_4^2 c_5^2 + d^2(K_0) + 2c_4 c_5 d(K_0)}{2d(K_0)}$ . Then with (3.6), we update (3.3) as

$$\frac{d}{dt} \|u\|_{L^2}^2 \leq \frac{c_5^2}{d(K_0)} \|u\|_{L^2}^2 \|\Delta v\|_{L^2}^2 + (2\gamma F(K_0) + c_6) \|u\|_{L^2}^2 \leq c_7 \|u\|_{L^2}^2 (\|\Delta v\|_{L^2}^2 + 1), \tag{3.7}$$

with  $c_7 := \frac{c_5^2}{d(K_0)} + 2\gamma F(K_0) + c_6$ . For any  $t \in (0, T_{max})$  and in the case of either  $t \in (0, \tau)$  or  $t \geq \tau$  with  $\tau = \min \left\{ 1, \frac{1}{2} T_{max} \right\}$ , from (2.5) we can find a  $t_0 = t_0(t) \in ((t - \tau)_+, t)$  such that  $t_0 \geq 0$  and

$$\int_{\Omega} u^2(x, t_0) dx \leq c_8. \tag{3.8}$$

On the other hand, from (2.16) in Lemma 2.4, we can find a constant  $c_9 > 0$  such that

$$\int_{t_0}^{t_0+\tau} \int_{\Omega} |\Delta v(x, s)|^2 dx ds \leq c_9 \text{ for all } t_0 \in (0, \tilde{T}_{max}). \tag{3.9}$$

Then integrating (3.7) over  $(t_0, t)$ , and using (3.8), (3.9) and the fact  $t \leq t_0 + \tau \leq t_0 + 1$ , we derive

$$\|u(\cdot, t)\|_{L^2}^2 \leq \|u(\cdot, t_0)\|_{L^2}^2 \cdot e^{c_7 \int_{t_0}^t (\|\Delta v(\cdot, s)\|_{L^2}^2 + 1) ds} \leq c_8 e^{c_7(c_9+1)},$$

which yields (3.1) and completes the proof of Lemma 3.1. □

Next, we will derive the boundedness of  $\|u(\cdot, t)\|_{L^\infty}$  by the Moser iteration.

**Lemma 3.2** *Let the conditions in Theorem 1.1 hold. Then the solution of system (1.3) satisfies*

$$\|u(\cdot, t)\|_{L^\infty} \leq C \text{ for all } t \in (0, T_{max}), \tag{3.10}$$

where the constant  $C > 0$  independent of  $t$ .

**Proof** First, we claim that if  $\|u(\cdot, t)\|_{L^p} \leq M_0 (p \geq 1)$ , then it holds that

$$\|\nabla v(\cdot, t)\|_{L^r} \leq c_1, \text{ for all } t \in (0, T_{max}), \tag{3.11}$$

with

$$r \in \begin{cases} [1, \frac{np}{n-p}), & \text{if } p < n, \\ [1, \infty), & \text{if } p = n, \\ [1, \infty], & \text{if } p > n. \end{cases} \tag{3.12}$$

In fact, from the second equation of system (1.3), we know that  $v$  solves the following problem:

$$v_t = D\Delta v - v + g(u, v) \text{ in } \Omega, \quad \frac{\partial v}{\partial \nu} = 0, \tag{3.13}$$

where  $g(u, v) := v - uF(v) + f(v)$ . Noting the properties of  $F(v), f(v)$  and the fact that  $0 \leq v(x, t) \leq K_0$  in (2.3), one has

$$\|g(u, v)\|_{L^p} \leq c_2(\|u\|_{L^p} + 1) \leq c_2(M_0 + 1) := c_3. \tag{3.14}$$

Then applying the results of Ref. [19, Lemma 1] (see also Ref. [39, Lemma 1.2]) to the problem (3.13) with (3.14), we obtain (3.11) with (3.12).

Using  $u^{p-1}$  with  $p \geq 2$  as a test function for the first equation in (1.3), and integrating the resulting equation by parts, we obtain

$$\begin{aligned} & \frac{1}{p} \frac{d}{dt} \int_{\Omega} u^p dx + (p-1) \int_{\Omega} d(v) u^{p-2} |\nabla u|^2 dx + \alpha \int_{\Omega} u^{p+1} dx + \theta \int_{\Omega} u^p dx \\ & = (p-1) \int_{\Omega} \chi(v) u^{p-1} \nabla u \cdot \nabla v dx + \gamma \int_{\Omega} F(v) u^p dx. \end{aligned} \tag{3.15}$$

Using the facts  $0 \leq \chi(v) \leq c_4$ ,  $d(v) \geq d(K_0) > 0$ ,  $0 < F(v) \leq F(K_0)$ ,  $\alpha \geq 0$  and applying the Young’s inequality, from (3.15) we obtain

$$\begin{aligned} & \frac{1}{p} \frac{d}{dt} \int_{\Omega} u^p dx + (p-1)d(K_0) \int_{\Omega} u^{p-2} |\nabla u|^2 dx + \theta \int_{\Omega} u^p dx \\ & \leq c_4(p-1) \int_{\Omega} u^{p-1} |\nabla u| |\nabla v| dx + \gamma F(K_0) \int_{\Omega} u^p dx \\ & \leq \frac{(p-1)d(K_0)}{2} \int_{\Omega} u^{p-2} |\nabla u|^2 dx + \frac{c_4^2(p-1)}{2d(K_0)} \int_{\Omega} u^p |\nabla v|^2 dx + \gamma F(K_0) \int_{\Omega} u^p dx, \end{aligned}$$

which along with the fact

$$\frac{p(p-1)d(K_0)}{2} \int_{\Omega} u^{p-2} |\nabla u|^2 dx = \frac{2(p-1)d(K_0)}{p} \int_{\Omega} |\nabla u^{\frac{p}{2}}|^2 dx,$$

gives

$$\begin{aligned} & \frac{d}{dt} \int_{\Omega} u^p dx + p\theta \int_{\Omega} u^p dx + \frac{2(p-1)d(K_0)}{p} \int_{\Omega} |\nabla u^{\frac{p}{2}}|^2 dx \\ & \leq \frac{c_4^2 p(p-1)}{2d(K_0)} \int_{\Omega} u^p |\nabla v|^2 dx + \gamma F(K_0) \int_{\Omega} u^p dx, \end{aligned} \tag{3.16}$$

for all  $t \in (0, T_{max})$  and for all  $p \geq 2$ . From Lemma 3.1, one has  $\|u(\cdot, t)\|_{L^2} \leq c_5$  and hence  $\|\nabla v(\cdot, t)\|_{L^4} \leq c_6$  by noting (3.11). Then using the Hölder inequality and the Gagliardo–Nirenberg inequality with the fact  $\|u^{\frac{p}{2}}(\cdot, t)\|_{L^{\frac{4}{p}}} = \|u(\cdot, t)\|_{L^2}^{\frac{p}{2}} \leq c_5^{\frac{p}{2}}$ , one has for all  $t \in (0, T_{max})$

$$\begin{aligned}
 \frac{c_4^2 p(p-1)}{2d(K_0)} \int_{\Omega} u^p |\nabla v|^2 dx &\leq \frac{c_4^2 p(p-1)}{2d(K_0)} \left( \int_{\Omega} u^{2p} dx \right)^{\frac{1}{2}} \left( \int_{\Omega} |\nabla v|^4 dx \right)^{\frac{1}{2}} \\
 &\leq \frac{c_4^2 c_6^2 p(p-1)}{2d(K_0)} \|u^{\frac{p}{2}}\|_{L^4}^2 \\
 &\leq c_7 \left( \|\nabla u^{\frac{p}{2}}\|_{L^2}^{2(1-\frac{1}{p})} \|u^{\frac{p}{2}}\|_{L^{\frac{4}{p}}}^{\frac{2}{p}} + \|u^{\frac{p}{2}}\|_{L^{\frac{4}{p}}}^2 \right) \tag{3.17} \\
 &\leq c_7 c_5 \|\nabla u^{\frac{p}{2}}\|_{L^2}^{2(1-\frac{1}{p})} + c_7 c_5^p \\
 &\leq \frac{(p-1)d(K_0)}{p} \|\nabla u^{\frac{p}{2}}\|_{L^2}^2 + \frac{d(K_0)}{p} \left( \frac{c_7 c_5}{d(K_0)} \right)^p + c_7 c_5^p,
 \end{aligned}$$

and

$$\begin{aligned}
 \gamma F(K_0) \int_{\Omega} u^p dx &= \gamma F(K_0) \|u^{\frac{p}{2}}\|_{L^2}^2 \\
 &\leq c_8 \left( \|\nabla u^{\frac{p}{2}}\|_{L^2}^{2(1-\frac{2}{p})} \|u^{\frac{p}{2}}\|_{L^{\frac{4}{p}}}^{\frac{4}{p}} + \|u^{\frac{p}{2}}\|_{L^{\frac{4}{p}}}^2 \right) \tag{3.18} \\
 &\leq c_8 c_5^2 \|\nabla u^{\frac{p}{2}}\|_{L^2}^{2(1-\frac{2}{p})} + c_8 c_5^p \\
 &\leq \frac{(p-1)d(K_0)}{p} \|\nabla u^{\frac{p}{2}}\|_{L^2}^2 + \frac{d(K_0)}{p} \left( \frac{c_8 c_5^2}{d(K_0)} \right)^p + c_8 c_5^p.
 \end{aligned}$$

Substituting (3.17) and (3.18) into (3.16), and letting  $c_9 := \frac{d(K_0)}{p} \left( \frac{c_7 c_5}{d(K_0)} \right)^p + \frac{d(K_0)}{p} \left( \frac{c_8 c_5^2}{d(K_0)} \right)^p + (c_7 + c_8)c_5^p$ , we obtain

$$\frac{d}{dt} \int_{\Omega} u^p dx + p\theta \int_{\Omega} u^p dx \leq c_9,$$

which, combined with the Gronwall’s inequality, yields

$$\|u(\cdot, t)\|_{L^p}^p \leq e^{-p\theta t} \|u_0\|_{L^p}^p + \frac{c_9}{p\theta} (1 - e^{-p\theta t}) \leq \|u_0\|_{L^p}^p + \frac{c_9}{p\theta}. \tag{3.19}$$

Then choosing  $p = 4$  in (3.19) and using (3.11) again, one can find a constant  $c_{10} > 0$  independent of  $p$  such that  $\|\nabla v(\cdot, t)\|_{L^\infty} \leq c_7$ . Then using the Moser iteration procedure (cf. Ref. [1]), one has (3.10). Hence, we complete the proof of Lemma 3.2. □

**Proof of Theorem 1.1** Theorem 1.1 is a consequence of Lemmas 2.1 and 3.2. □

### 4 Globally asymptotic stability of solutions

Based on some ideas in Ref. [12], we shall prove the global stability results in Theorem 1.2 in this section by the method of Lyapunov functionals with the help of LaSalle’s invariant principle under the hypotheses (H1)–(H4). Here, we employ the same Lyapunov functionals as

in Ref. [12] and hence will skip many similar computations. To proceed, we first derive some regularity results for the solution  $(u, v)$  by using some ideas in Ref. [38].

**Lemma 4.1** *Let  $(u, v)$  be the non-negative global classical solution of system (1.3) obtained in Theorem 1.1. Then there exist  $\sigma \in (0, 1)$  and  $C > 0$  such that*

$$\|v\|_{C^{2+\sigma, 1+\frac{\sigma}{2}}(\bar{\Omega} \times [t, t+1])} \leq C, \text{ for all } t > 1. \tag{4.1}$$

**Proof** From Theorem 1.1, we can find three positive constants  $c_1, c_2, c_3$  such that

$$0 < u(x, t) \leq c_1, 0 < v(x, t) \leq c_2 \text{ and } |\nabla v(x, t)| \leq c_3 \text{ for all } x \in \Omega \text{ and } t > 0.$$

The first equation of system (1.3) can be rewritten as

$$u_t = \nabla \cdot A(x, t, \nabla u) + B(x, t) \text{ for all } x \in \Omega \text{ and } t > 0,$$

where

$$A(x, t, \nabla u) := d(v)\nabla u - \chi(v)u\nabla v,$$

and

$$B(x, t) := (\gamma F(v) - \theta - \alpha u)u.$$

By the assumptions in (H1)–(H3) and using the Young’s inequality, then for all  $x \in \Omega$  and  $t > 0$ , we obtain that

$$\begin{aligned} A(x, t, \nabla u) \cdot \nabla u &= d(v)|\nabla u|^2 - \chi(v)u\nabla v \cdot \nabla u \\ &\geq d(v)|\nabla u|^2 - |\chi(v)u|\nabla v|\nabla u| \\ &\geq \frac{d(v)}{2}|\nabla u|^2 - \frac{|\chi(v)|^2}{2d(v)}u^2|\nabla v|^2 \\ &\geq \frac{d(c_2)}{2}|\nabla u|^2 - c_4, \end{aligned} \tag{4.2}$$

and

$$|A(x, t, \nabla u)| \leq d(0)|\nabla u| + c_5, \tag{4.3}$$

as well as

$$|B(x, t)| \leq c_6. \tag{4.4}$$

Then (4.2)–(4.4) allow us to apply the Hölder regularity for quasilinear parabolic equations [32, Theorem 1.3 and Remark 1.4] to obtain  $\|u\|_{C^{\sigma, \frac{\sigma}{2}}(\bar{\Omega} \times [t, t+1])} \leq c_7$  for all  $t > 1$ . Moreover, applying the standard parabolic Schauder theory [20] to the second equation of (1.3), one has (4.1). Then the proof of Lemma 4.1 is completed.  $\square$

With the results in Lemma 4.1 in hand, we next derive the following results.

**Lemma 4.2** *Suppose the conditions in Lemma 4.1 hold. Then we can find a constant  $C > 0$  such that*

$$\|\nabla u\|_{L^4} \leq C \text{ for all } t > 1. \tag{4.5}$$

**Proof** From the first equation of system (1.3), we have

$$\begin{aligned} \frac{1}{4} \frac{d}{dt} \int_{\Omega} |\nabla u|^4 dx &= \int_{\Omega} |\nabla u|^2 \nabla u \cdot \nabla u_t dx \\ &= \int_{\Omega} |\nabla u|^2 \nabla u \cdot \nabla (\nabla \cdot (d(v)\nabla u)) dx - \int_{\Omega} |\nabla u|^2 \nabla u \cdot \nabla (\nabla \cdot (\chi(v)u\nabla v)) dx \\ &\quad + \int_{\Omega} \nabla(\gamma F(v)u - \theta u - \alpha u^2) \cdot \nabla u |\nabla u|^2 dx \\ &=: I_1 + I_2 + I_3. \end{aligned} \tag{4.6}$$

With the integration by parts, the term  $I_1$  becomes

$$\begin{aligned} I_1 &= - \int_{\Omega} (|\nabla u|^2 \Delta u) \nabla \cdot (d(v)\nabla u) dx - \int_{\Omega} (\nabla |\nabla u|^2 \cdot \nabla u) \nabla \cdot (d(v)\nabla u) dx \\ &= \int_{\Omega} d(v) |\nabla u|^2 (\nabla \Delta u \cdot \nabla u) dx - \int_{\Omega} d'(v) (\nabla |\nabla u|^2 \cdot \nabla u) (\nabla u \cdot \nabla v) dx, \end{aligned}$$

which, combined with the fact  $\nabla \Delta u \cdot \nabla u = \frac{1}{2} \Delta |\nabla u|^2 - |D^2 u|^2$  where  $D^2 u$  denotes the Hessian matrix of  $u$ , gives

$$\begin{aligned} I_1 &= \frac{1}{2} \int_{\Omega} d(v) |\nabla u|^2 \Delta |\nabla u|^2 dx - \int_{\Omega} d(v) |\nabla u|^2 |D^2 u|^2 dx - \int_{\Omega} d'(v) (\nabla |\nabla u|^2 \cdot \nabla u) (\nabla u \cdot \nabla v) dx \\ &= \frac{1}{2} \int_{\partial\Omega} d(v) |\nabla u|^2 \frac{\partial |\nabla u|^2}{\partial \nu} dS - \frac{1}{2} \int_{\Omega} d'(v) |\nabla u|^2 \nabla v \cdot \nabla |\nabla u|^2 dx - \frac{1}{2} \int_{\Omega} d(v) |\nabla |\nabla u|^2|^2 dx \\ &\quad - \int_{\Omega} d(v) |\nabla u|^2 |D^2 u|^2 dx - \int_{\Omega} d'(v) (\nabla |\nabla u|^2 \cdot \nabla u) (\nabla u \cdot \nabla v) dx \\ &\leq \frac{1}{2} \int_{\partial\Omega} d(v) |\nabla u|^2 \frac{\partial |\nabla u|^2}{\partial \nu} dS - \frac{1}{2} \int_{\Omega} d(v) |\nabla |\nabla u|^2|^2 dx - \int_{\Omega} d(v) |\nabla u|^2 |D^2 u|^2 dx \\ &\quad + \frac{3}{2} \int_{\Omega} |d'(v)| |\nabla |\nabla u|^2| |\nabla u|^2 |\nabla v| dx. \end{aligned} \tag{4.7}$$

With the facts  $\|u(\cdot, t)\|_{L^\infty} + \|v(\cdot, t)\|_{W^{1,\infty}} \leq c_1$  and (4.1), we have

$$\begin{aligned} \nabla \cdot (\chi(v)u\nabla v) &= \chi'(v)u|\nabla v|^2 + \chi(v)\nabla u \cdot \nabla v + \chi(v)u\Delta v \\ &\leq c_2(1 + |\nabla u|) \text{ for all } t > 1, \end{aligned}$$

which updates  $I_2$  as

$$\begin{aligned} I_2 &= \int_{\Omega} \nabla |\nabla u|^2 \cdot \nabla u \nabla \cdot (\chi(v)u\nabla v) dx + \int_{\Omega} |\nabla u|^2 \Delta u \nabla \cdot (\chi(v)u\nabla v) dx \\ &\leq c_2 \int_{\Omega} |\nabla u| |\nabla |\nabla u|^2| (1 + |\nabla u|) dx + c_2 \int_{\Omega} |\nabla u|^2 |\Delta u| (1 + |\nabla u|) dx. \end{aligned} \tag{4.8}$$

Moreover, the term  $I_3$  can be estimated as follows:

$$\begin{aligned}
 I_3 &= \int_{\Omega} \nabla(\gamma F(v)u - \theta u - \alpha u^2) \cdot \nabla u |\nabla u|^2 dx \\
 &= \int_{\Omega} \gamma F'(v)u \nabla v \cdot \nabla u |\nabla u|^2 dx + \gamma \int_{\Omega} F(v) |\nabla u|^4 dx - \theta \int_{\Omega} |\nabla u|^4 dx - 2\alpha \int_{\Omega} u |\nabla u|^4 dx \quad (4.9) \\
 &\leq c_3 \int_{\Omega} |\nabla u|^4 dx + c_4.
 \end{aligned}$$

Substituting (4.7)–(4.9) into (4.6), and using the fact  $0 < d(K_0) \leq d(v) \leq d(0)$ , we have

$$\begin{aligned}
 &\frac{1}{4} \frac{d}{dt} \int_{\Omega} |\nabla u|^4 dx + \frac{d(K_0)}{2} \int_{\Omega} |\nabla |\nabla u|^2|^2 dx + d(K_0) \int_{\Omega} |\nabla u|^2 |D^2 u|^2 dx \\
 &\leq \frac{1}{2} \int_{\partial\Omega} d(v) |\nabla u|^2 \frac{\partial |\nabla u|^2}{\partial \nu} dS + \frac{3}{2} \int_{\Omega} |d'(v)| |\nabla |\nabla u|^2| |\nabla u|^2 |\nabla v| dx \\
 &\quad + c_2 \int_{\Omega} |\nabla u| |\nabla |\nabla u|^2| (1 + |\nabla u|) dx + c_2 \int_{\Omega} |\nabla u|^2 |\Delta u| (1 + |\nabla u|) dx \quad (4.10) \\
 &\quad + c_3 \int_{\Omega} |\nabla u|^4 dx + c_4 \\
 &\triangleq J_1 + J_2 + J_3 + J_4 + c_3 \int_{\Omega} |\nabla u|^4 dx + c_4.
 \end{aligned}$$

With the inequality  $\frac{\partial |\nabla u|^2}{\partial \nu} \leq 2\lambda' |\nabla u|^2$  on  $\partial\Omega$  (see Ref. [28, Lemma 4.2]) we derive

$$\begin{aligned}
 J_1 &\leq \frac{d(0)}{2} \left| \int_{\partial\Omega} |\nabla u|^2 \frac{\partial |\nabla u|^2}{\partial \nu} dS \right| \\
 &\leq \lambda' d(0) \| |\nabla u|^2 \|_{L^2(\partial\Omega)}^2 \leq \frac{d(K_0)}{8} \int_{\Omega} |\nabla |\nabla u|^2|^2 dx + c_5 \int_{\Omega} |\nabla u|^4 dx, \quad (4.11)
 \end{aligned}$$

where we have used the following trace inequality (cf. Ref. [36, Remark 52.9]):

$$\|\varphi\|_{L^2(\partial\Omega)} \leq \varepsilon \|\nabla \varphi\|_{L^2(\Omega)} + C_{\varepsilon} \|\varphi\|_{L^2(\Omega)}, \text{ for any } \varepsilon > 0.$$

By the boundedness of  $\|\nabla v\|_{L^\infty}$  and Young’s inequality, one derives

$$\begin{aligned}
 J_2 + J_3 &\leq c_6 \int_{\Omega} |\nabla |\nabla u|^2| |\nabla u| (1 + |\nabla u|) dx \\
 &\leq \frac{d(K_0)}{8} \int_{\Omega} |\nabla |\nabla u|^2|^2 dx + c_7 \int_{\Omega} |\nabla u|^4 dx + c_7. \quad (4.12)
 \end{aligned}$$

Since  $|\Delta u| \leq \sqrt{2} |D^2 u|$ , one can estimate  $J_4$  as follows:

$$\begin{aligned}
 J_4 &\leq c_2 \sqrt{2} \int_{\Omega} |\nabla u|^2 |D^2 u| dx + c_2 \sqrt{2} \int_{\Omega} |\nabla u|^3 |D^2 u| dx \\
 &\leq \frac{d(K_0)}{2} \int_{\Omega} |\nabla u|^2 |D^2 u|^2 dx + \frac{2c_2^2}{d(K_0)} \int_{\Omega} (|\nabla u|^2 + |\nabla u|^4) dx \quad (4.13) \\
 &\leq \frac{d(K_0)}{2} \int_{\Omega} |\nabla u|^2 |D^2 u|^2 dx + \frac{4c_2^2}{d(K_0)} \int_{\Omega} |\nabla u|^4 dx + \frac{c_2^2 |\Omega|}{2d(K_0)}.
 \end{aligned}$$

Substituting (4.11)–(4.13) into (4.10), we end up with

$$\begin{aligned} & \frac{d}{dt} \int_{\Omega} |\nabla u|^4 dx + d(K_0) \int_{\Omega} |\nabla |\nabla u|^2|^2 dx + 2d(K_0) \int_{\Omega} |\nabla u|^2 |D^2 u|^2 dx \\ & \leq c_8 \int_{\Omega} |\nabla u|^4 dx + c_9, \end{aligned} \tag{4.14}$$

where  $c_8 := 4 \left( c_3 + c_5 + c_7 + \frac{4c_2^2}{d(K_0)} \right)$  and  $c_9 := 4 \left( c_4 + c_7 + \frac{c_2^2 |\Omega|}{2d(K_0)} \right)$ . On the other hand, integrating by parts, noting  $\|u(\cdot, t)\|_{L^\infty} \leq c_{10}$  and using the Young’s inequality, one has

$$\begin{aligned} (c_8 + 2) \int_{\Omega} |\nabla u|^4 dx &= c_{11} \int_{\Omega} |\nabla u|^2 \nabla u \cdot \nabla u dx \\ &= -c_{11} \int_{\Omega} u \nabla |\nabla u|^2 \cdot \nabla u dx - c_{11} \int_{\Omega} u |\nabla u|^2 \Delta u dx \\ &\leq c_{10} c_{11} \int_{\Omega} |\nabla |\nabla u|^2| |\nabla u| dx + c_{10} c_{11} \sqrt{2} \int_{\Omega} |\nabla u|^2 |D^2 u| dx \\ &\leq d(K_0) \int_{\Omega} |\nabla |\nabla u|^2|^2 dx + 2d(K_0) \int_{\Omega} |\nabla u|^2 |D^2 u|^2 dx + \frac{c_{10}^2 c_{11}^2}{2} \int_{\Omega} |\nabla u|^2 dx \\ &\leq d(K_0) \int_{\Omega} |\nabla |\nabla u|^2|^2 dx + 2d(K_0) \int_{\Omega} |\nabla u|^2 |D^2 u|^2 dx \\ &\quad + \int_{\Omega} |\nabla u|^4 dx + \frac{|\Omega| c_{10}^4 c_{11}^4}{16}, \end{aligned}$$

which updates (4.14) to

$$\frac{d}{dt} \int_{\Omega} |\nabla u|^4 dx + \int_{\Omega} |\nabla u|^4 dx \leq c_9 + \frac{c_{10}^4 c_{11}^4 |\Omega|}{16}. \tag{4.15}$$

Applying the Gronwall’s inequality to (4.15) along with the fact  $\|\nabla u(\cdot, 1)\|_{L^\infty} \leq c_{12}$ , one obtains (4.5). Thus the proof of Lemma 4.2 is finished.  $\square$

Next, we state a basic result which will be used later.

**Lemma 4.3** ([12]) *Let  $F$  satisfy the conditions in (H2) and define a function for some constant  $\omega_* > 0$ :*

$$\zeta(v) = \int_{\omega_*}^v \frac{F(\sigma) - F(\omega_*)}{F(\sigma)} d\sigma.$$

*Then  $\zeta(v)$  is a convex function such that  $\zeta(v) \geq 0$ . If  $(u, v)$  is a solution of (1.3) satisfying  $v \rightarrow \omega_*$  as  $t \rightarrow \infty$ , then there is a constant  $T_0 > 0$  such that for all  $t \geq T_0$  it holds that*

$$\frac{F'(\omega_*)}{4F(\omega_*)} (v - \omega_*)^2 \leq \zeta(v) = \int_{\omega_*}^v \frac{F(\sigma) - F(\omega_*)}{F(\sigma)} d\sigma \leq \frac{F'(\omega_*)}{F(\omega_*)} (v - \omega_*)^2.$$



**4.1 Global stability of the prey-only steady state**

In this subsection, we shall prove that  $(u, v)$  converges to  $(0, K)$  in  $L^\infty$  as  $t \rightarrow \infty$  when  $\gamma F(K) \leq \theta$  and further show that the convergence rate is exponential if  $\gamma F(K) < \theta$  and algebraic if  $\gamma F(K) = \theta$  and  $\alpha > 0$ .

**Lemma 4.4** *Let the assumptions in Theorem 1.2 hold. If  $\gamma F(K) \leq \theta$ , then the solution  $(u, v)$  of (1.3) satisfies*

$$\|u\|_{L^\infty} + \|v - K\|_{L^\infty} \rightarrow 0 \text{ as } t \rightarrow \infty,$$

where the convergence rate is exponential if  $\gamma F(K) < \theta$  and algebraic if  $\gamma F(K) = \theta$  and  $\alpha > 0$ .

**Proof** We start with the following functional:

$$V_1(u(t), v(t)) =: V_1(t) = \frac{1}{\gamma} \int_{\Omega} u dx + \int_{\Omega} \int_K^v \frac{F(\sigma) - F(K)}{F(\sigma)} d\sigma dx, \tag{4.16}$$

and show  $\frac{d}{dt} V_1(u, v) = \frac{d}{dt} V_1(t) \leq 0$  as well as  $\frac{d}{dt} V_1(u, v) = 0$  iff  $(u, v) = (0, K)$ . Indeed differentiating the functional (4.16) with respect to  $t$  and using the equations in (1.3), one has

$$\begin{aligned} \frac{d}{dt} V_1(t) &= \frac{1}{\gamma} \int_{\Omega} u_t dx + \int_{\Omega} \frac{F(v) - F(K)}{F(v)} v_t dx \\ &= \frac{1}{\gamma} \int_{\Omega} u(\gamma F(v) - \theta - \alpha u) dx + \int_{\Omega} \left(1 - \frac{F(K)}{F(v)}\right) (D\Delta v - uF(v) + f(v)) dx. \end{aligned} \tag{4.17}$$

For the second term on the right-hand side of (4.17), we use the integration by parts with some calculations and cancelations to have

$$\begin{aligned} \frac{d}{dt} V_1(t) &= \frac{1}{\gamma} \int_{\Omega} u(\gamma F(K) - \theta - \alpha u) dx - DF(K) \int_{\Omega} F'(v) \left| \frac{\nabla v}{F(v)} \right|^2 dx \\ &\quad + \int_{\Omega} \frac{f(v)}{F(v)} (F(v) - F(K)) dx. \end{aligned}$$

The rest of the proof only depends on the assumption (H2)–(H4) and hence we can follow the exact procedures in the proof of Ref. [12, Lemma 4.2] along with the LaSalle’s invariant principle (cf. Refs. [18, 21]) (or the compact method together with the Lyapunov functional as in Refs. [43, 45]) to prove that  $(0, K)$  is globally asymptotically stable if  $\gamma F(K) \leq \theta$  and the following convergence:

$$\begin{cases} \|u\|_{L^1} + \|v - K\|_{L^2}^2 \leq c_1 e^{-c_1 t}, & \text{if } \gamma F(K) < \theta, \\ \|u\|_{L^1} + \|v - K\|_{L^2}^2 \leq c_2 (1 + t)^{-1}, & \text{if } \gamma F(K) = \theta, \alpha > 0 \end{cases} \tag{4.18}$$

hold for  $t > t_0$  with some  $t_0 > 1$ . Furthermore, the Gagliardo–Nirenberg inequality and the result  $\|\nabla u\|_{L^4} \leq c_3$  for  $t > t_0$  (cf. Lemma 4.2) entail that

$$\|u\|_{L^\infty} \leq c_4 \|\nabla u\|_{L^4}^{\frac{4}{3}} \|u\|_{L^1}^{\frac{1}{3}} + c_4 \|u\|_{L^1} \leq c_4 c_3^{\frac{4}{3}} \|u\|_{L^1}^{\frac{1}{3}} + c_4 \|u\|_{L^1}, \tag{4.19}$$

which together with (4.18) yields the decay rate of  $u$ . Similarly, with  $\|\nabla v\|_{L^4} \leq c_5$  (cf. (3.11)), we obtain

$$\begin{aligned} \|v - K\|_{L^\infty} &\leq c_6 \|\nabla v\|_{L^4}^{\frac{2}{3}} \|v - K\|_{L^2}^{\frac{1}{3}} + c_6 \|v - K\|_{L^2} \\ &\leq c_6 c_5^{\frac{2}{3}} \|v - K\|_{L^2}^{\frac{1}{3}} + c_6 \|v - K\|_{L^2}, \end{aligned} \tag{4.20}$$

which combined with (4.18) gives the decay rate of  $v$ . This completes the proof of Lemma 4.4.  $\square$

### 4.2 Global stability of the coexistence steady state

Now we turn to the case  $\gamma F(K) > \theta$  and prove the homogeneous coexistence steady state  $(u_*, v_*)$  is globally asymptotically stable under certain conditions. We shall prove our result based on the following Lyapunov functional as in Ref. [12]:

$$V_2(u(t), v(t)) =: V_2(t) = \frac{1}{\gamma} \int_{\Omega} \left( u - u_* - u_* \ln \frac{u}{u_*} \right) dx + \int_{\Omega} \int_{v_*}^v \frac{F(\sigma) - F(v_*)}{F(\sigma)} d\sigma dx.$$

**Lemma 4.5** *Assume the assumptions in Theorem 1.2 hold. If  $\gamma F(K) > \theta$  and*

$$D \geq \max_{0 \leq v \leq K_0} \frac{u_* |F(v)|^2 |\chi(v)|^2}{4\gamma F(v_*) F'(v) d(v)}$$

where  $u_*$  and  $v_*$  are determined by (1.5) and independent of  $D$ , then the solution  $(u, v)$  of (1.3) satisfies

$$\|u - u_*\|_{L^\infty} + \|v - v_*\|_{L^\infty} \rightarrow 0 \text{ as } t \rightarrow \infty$$

where the convergence is exponential when  $\alpha > 0$ .

**Proof** First by the same argument as the proof of Ref. [12, Lemma 4.3], we have that  $V_2(t) \geq 0$  for all  $u, v \geq 0$ . Next we differentiate  $V_2(t)$  with respect to  $t$  and use the equations of (1.3) to obtain that

$$\begin{aligned} \frac{d}{dt} V_2(t) &= \frac{1}{\gamma} \int_{\Omega} \left( 1 - \frac{u_*}{u} \right) u_t dx + \int_{\Omega} \frac{F(v) - F(v_*)}{F(v)} v_t dx \\ &= \underbrace{-\frac{u_*}{\gamma} \int_{\Omega} \frac{d(v) |\nabla u|^2}{u^2} dx - DF(v_*) \int_{\Omega} F'(v) \left| \frac{\nabla v}{F(v)} \right|^2 dx + \frac{u_*}{\gamma} \int_{\Omega} \frac{\chi(v) \nabla u \cdot \nabla v}{u} dx}_{I_1} \\ &\quad + \underbrace{\frac{1}{\gamma} \int_{\Omega} \left( 1 - \frac{u_*}{u} \right) (\gamma u F(v) - \theta u - \alpha u^2) dx + \int_{\Omega} (F(v) - F(v_*)) \left( \frac{f(v)}{F(v)} - u \right) dx}_{I_2}. \end{aligned}$$

$I_1$  can be rewritten as

$$I_1 = - \int_{\Omega} \Theta^T A \Theta, \quad \Theta = \begin{bmatrix} \nabla u \\ \nabla v \end{bmatrix}, \quad A = \begin{bmatrix} \frac{u_* d(v)}{\gamma u^2} & -\frac{\chi(v) u_*}{2\gamma u} \\ -\frac{\chi(v) u_*}{2\gamma u} & \frac{DF(v_*) F'(v)}{|F(v)|^2} \end{bmatrix}.$$

where  $\Theta^T$  denotes the transpose of  $\Theta$ . Then, it can be easily checked with Sylvesters criterion that the matrix  $A$  is non-negative definite (and hence  $I_1 \leq 0$ ) if and only if

$$\frac{DF(v_*)F'(v)u_*d(v)}{\gamma|F(v)|^2u^2} \geq \frac{|\chi(v)|^2u_*^2}{4\gamma^2u^2} \text{ or } D \geq \frac{u_*|F(v)|^2|\chi(v)|^2}{4\gamma F(v_*)F'(v)d(v)},$$

where  $u_*$  and  $v_*$  do not depend on  $D$ , see (1.5). Note that the above  $I_2$  is exactly the same as the  $I_2$  in the proof of Ref. [12, Lemma 4.3]. Hence, we can follow the same procedure for the proof of Ref. [12, Lemma 4.3] to show the homogeneous coexistence state  $(u_*, v_*)$  is globally asymptotically stable and satisfies

$$\|u - u_*\|_{L^2} + \|v - v_*\|_{L^2} \leq c_1 e^{-c_2 t} \text{ for } t > t_0, \tag{4.21}$$

with some  $t_0 > 1$  for  $\alpha > 0$ . Using the Gagliardo–Nirenberg inequality and the boundedness of  $\|\nabla u\|_{L^4}$  and  $\|\nabla v\|_{L^\infty}$  (see Lemmas 4.1 and 4.2), we use a similar procedure as for (4.19)–(4.20) to finally obtain the exponential decay rate in  $L^\infty$ -norm from (4.21).  $\square$

### 4.3 Proof of Theorem 1.2

Theorem 1.2 is directly obtained from Lemmas 4.4 and 4.5.

## 5 Applications and spatio-temporal patterns

The first purpose of this section is to apply our general results obtained in Theorems 1.1 and 1.2 to two most widely used predator–prey interactions: Lotka–Volterra type (i.e.  $F(v) = v$ ) and Rosenzweig–MacArthur type (i.e.  $F(v) = \frac{v}{\lambda+v}$  and  $\alpha = 0$ ) [33]. Note that these results only give the global existence of solutions (by Theorem 1.1) and global stability of constant steady states (by Theorem 1.2), the distribution of the predator and the prey in space and the time-asymptotic dynamics of the population outside the parameter regimes found in Theorem 1.2 are unclear, but indeed they are more interesting from the application point of view in ecology though hard to study analytically. Hence, the second purpose of this section is to numerically exploit the spatio-temporal patterns generated by the Lotka–Volterra and Rosenzweig–MacArthur predator–prey systems, which not only display the distribution patterns of the predator and the prey in space or evolution of population in time but also provide useful sources for the future research.

### 5.1 Application of our results

In this subsection, we shall give some examples to illustrate the applications of Theorems 1.1 and 1.2. The first example is the Lotka–Volterra predator–prey system [25] with prey-taxis

$$\begin{cases} u_t = \nabla \cdot (d(v)\nabla u) - \nabla \cdot (u\chi(v)\nabla v) + \gamma uv - \theta u - \alpha u^2, & x \in \Omega, \ t > 0, \\ v_t = D\Delta v - uv + \mu v(1 - \frac{v}{K}), & x \in \Omega, \ t > 0, \\ \frac{\partial u}{\partial \nu} = \frac{\partial v}{\partial \nu} = 0, & x \in \partial\Omega, \ t > 0, \\ u(x, 0) = u_0(x), \ v(x, 0) = v_0(x), & x \in \Omega. \end{cases} \tag{5.1}$$

Then, the application of the results in Theorems 1.1 and 1.2 yields the following results on the system (5.1).

**Proposition 5.1** *Let  $\Omega \subset \mathbb{R}^2$  be a bounded domain with smooth boundary and assume  $(u_0, v_0) \in [W^{1,p}(\Omega)]^2$  with  $u_0, v_0 \geq 0 (\neq 0)$  and  $p > 2$ . Assume  $\alpha > 0$ , then the problem (5.1) has a unique global classical solution  $(u, v) \in [C([0, \infty) \times \bar{\Omega}) \cap C^{2,1}((0, \infty) \times \bar{\Omega})]^2$  such that:*

- If  $\gamma K \leq \theta$ , then

$$\|u\|_{L^\infty} + \|v - K\|_{L^\infty} \rightarrow 0 \text{ as } t \rightarrow \infty,$$

where the convergence is exponential if  $\gamma K < \theta$  and algebraic if  $\gamma K = \theta$ .

- If  $\gamma K > \theta$  and

$$\frac{4D\gamma K(\mu\alpha + \theta)}{\mu(\gamma K - \theta)K_0^2} \geq M_0,$$

with  $M_0 = \max_{0 \leq v \leq K_0} \frac{|\chi(v)|^2}{d(v)}$ , then

$$\|u - u^*\|_{L^\infty} + \|v - v^*\|_{L^\infty} \rightarrow 0 \text{ exponentially as } t \rightarrow \infty,$$

where

$$(u_*, v_*) = \left( \frac{\mu(\gamma K - \theta)}{\gamma K + \mu\alpha}, \frac{K(\mu\alpha + \theta)}{\gamma K + \mu\alpha} \right). \tag{5.2}$$

The second example is the Rosenzweig–MacArthur type predator–prey interaction. From Theorem 1.1 with  $\alpha = 0$ , we only have the results for the special case  $\chi(v) = -d'(v)$  (density suppressed motility) which leads to the following system:

$$\begin{cases} u_t = \Delta(d(v)u) + \frac{\gamma uv}{\lambda + v} - \theta u, & x \in \Omega, \quad t > 0, \\ v_t = D\Delta v - \frac{uv}{\lambda + v} + \mu v \left(1 - \frac{v}{K}\right), & x \in \Omega, \quad t > 0, \\ \frac{\partial u}{\partial \nu} = \frac{\partial v}{\partial \nu} = 0, & x \in \partial\Omega, \quad t > 0, \\ u(x, 0) = u_0(x), v(x, 0) = v_0(x), & x \in \Omega. \end{cases} \tag{5.3}$$

In this case, the hypothesis (H4) is satisfied by requiring  $\lambda > K$ . Then, the interpretation of our results of Theorems 1.1 and 1.2 into (5.3) yields the following results.

**Proposition 5.2** *Assume  $(u_0, v_0) \in [W^{1,p}(\Omega)]^2$  with  $u_0, v_0 \geq 0 (\neq 0)$  and  $p > 2$ . If  $\lambda > K$ , then the problem (5.3) has a unique global classical solution in  $\Omega \subset \mathbb{R}^2$  such that*

- If  $\frac{\gamma K}{\lambda + K} < \theta$ , then

$$\|u\|_{L^\infty} + \|v - K\|_{L^\infty} \rightarrow 0 \text{ exponentially as } t \rightarrow \infty.$$

- If  $\frac{\gamma K}{\lambda + K} > \theta$  and  $\frac{4D\theta\lambda}{K_0^2 u_*} \geq M_1$  with  $M_1 = \max_{0 \leq v \leq K_0} \frac{|d'(v)|^2}{d(v)}$ , it follows that

$$\|u - u^*\|_{L^\infty} + \|v - v^*\|_{L^\infty} \rightarrow 0 \text{ as } t \rightarrow \infty,$$

where

$$(u_*, v_*) = \left( \frac{\gamma\lambda\mu[(\gamma - \theta)K - \theta\lambda]}{(\gamma - \theta)^2K}, \frac{\theta\lambda}{\gamma - \theta} \right).$$

### 5.2 Linear instability analysis

In this section, we will study the possible pattern formation generated by the system (1.3) under the assumptions (H1)–(H3). We begin with the space-absent ordinary differential equation (ODE) system of (1.3):

$$\begin{cases} u_t = \gamma uF(v) - \theta u - \alpha u^2, \\ v_t = -uF(v) + f(v), \end{cases} \tag{5.4}$$

which has three possible equilibria:  $(0, 0)$ ,  $(0, K)$  and  $(u_*, v_*)$ . One can easily check that the steady state  $(0, 0)$  is linearly unstable and  $(0, K)$  is linearly stable for both Lotka–Volterra and Rosenzweig–MacArthur type predator–prey interactions. For the homogeneous coexistence steady state  $(u_*, v_*)$ , it is linearly stable for the Lotka–Volterra type interaction. While for the Rosenzweig–MacArthur type interaction, one can easily find that all the eigenvalues of the linearised system of (5.4) at  $(u_*, v_*)$  have negative real part (hence  $(u_*, v_*)$  is stable) if  $v_* > \frac{K-\lambda}{2}$ , and are complex with zero real part if  $v_* = \frac{K-\lambda}{2}$ , and are complex with positive real part (hence  $(u_*, v_*)$  is unstable) if  $0 < v_* < \frac{K-\lambda}{2}$ .

Next we proceed to consider the stability of equilibria  $(0, K)$  and  $(u_*, v_*)$  in the presence of spatial structure. To this end, we first linearise the system (1.3) at an equilibrium  $(u_s, v_s)$  and write the linearised system as

$$\begin{cases} \Phi_t = \mathcal{A}\Delta\Phi + \mathcal{B}\Phi, & x \in \Omega, t > 0, \\ (\nu \cdot \nabla)\Phi = 0, & x \in \partial\Omega, t > 0, \\ \Phi(x, 0) = (u_0 - u_s, v_0 - v_s)^T, & x \in \Omega, \end{cases} \tag{5.5}$$

where  $\mathcal{T}$  denotes the transpose and

$$\Phi = \begin{pmatrix} u - u_s \\ v - v_s \end{pmatrix}, \quad \mathcal{A} = \begin{pmatrix} d(v_s) & -u_s\chi(v_s) \\ 0 & D \end{pmatrix},$$

as well as

$$\mathcal{B} = \begin{pmatrix} -\alpha u_s & \gamma u_s F'(v_s) \\ -F(v_s) & -u_s F'(v_s) + f'(v_s) \end{pmatrix} = \begin{pmatrix} B_1 & B_2 \\ B_3 & B_4 \end{pmatrix}.$$

Let  $W_k(x)$  denote the eigenfunction of the following eigenvalue problem:

$$\Delta W_k(x) + k^2 W_k(x) = 0, \quad \frac{\partial W_k(x)}{\partial \nu} = 0,$$

where  $k$  is called the wavenumber. Since the system (5.5) is linear, the solution  $\Phi(x, t)$  has the form of

$$\Phi(x, t) = \sum_{k \geq 0} c_k e^{\rho t} W_k(x), \tag{5.6}$$

where the constants  $c_k$  are determined by the Fourier expansion of the initial conditions in terms of  $W_k(x)$  and  $\rho$  is the temporal eigenvalue. Substituting (5.6) into (5.5), one has

$$\rho W_k(x) = -k^2 A W_k(x) + B W_k(x),$$

which implies  $\rho$  is the eigenvalue of the following matrix:

$$\begin{aligned} M_k &= \begin{pmatrix} -d(v_s)k^2 - \alpha u_s & u_s \chi(v_s)k^2 + \gamma u_s F'(v_s) \\ -F(v_s) & -Dk^2 - u_s F'(v_s) + f'(v_s) \end{pmatrix}, \\ &= \begin{pmatrix} -d(v_s)k^2 + B_1 & u_s \chi(v_s)k^2 + B_2 \\ B_3 & -Dk^2 + B_4 \end{pmatrix}. \end{aligned}$$

Calculating the eigenvalue of matrix  $M_k$ , we get the eigenvalues  $\rho(k^2)$  as functions of the wavenumber  $k$  as the roots of

$$\rho^2 + a(D, k^2)\rho + b(D, k^2) = 0,$$

where

$$\begin{aligned} a(D, k^2) &= (d(v_s) + D)k^2 + (\alpha + F'(v_s))u_s - f'(v_s), \\ &= (d(v_s) + D)k^2 - (B_1 + B_4), \end{aligned}$$

and

$$b(D, k^2) = d(v_s)Dk^4 - (d(v_s)B_4 + u_s \chi(v_s)B_3 + B_1 D)k^2 + B_1 B_4 - B_2 B_3.$$

Then, it can be easily verified that the eigenvalue  $\rho$  for the prey-only steady state  $(0, K)$  has negative real part for both Lotka–Volterra and Rosenzweig–MacArthur type predator–prey interactions and hence  $(0, K)$  is linearly stable. Thus, the pattern (if any) can only arise from the homogeneous coexistence steady state  $(u_*, v_*)$ . In the following results, we first show that the Lotka–Volterra type predator–prey system (5.1) indeed has no pattern bifurcated from  $(u_*, v_*)$ .

**Lemma 5.3** *The homogeneous coexistence steady state  $(u_*, v_*)$  of system (5.1) is linearly stable if  $\gamma K > \theta$ .*

**Proof** Since  $F(v) = v$  and  $f(v) = \mu v(1 - v/K)$ , we can easily check that

$$B_1 = -\alpha u_* < 0, B_2 = \gamma u_* > 0, B_3 = -v_* < 0, B_4 = -\frac{\mu}{K} v_* < 0.$$

Hence,  $B_1 + B_4 < 0$  and  $B_1 B_4 - B_2 B_3 > 0$  as well as

$$d(v_*)B_4 + u_* \chi(v_*)B_3 + B_1 D < 0,$$

which imply  $a(D, k^2) > 0$  and  $b(D, k^2) > 0$ . Then, the corresponding characteristic equation only has eigenvalue with negative real part. Hence,  $(u_*, v_*)$  is linearly stable for all  $\gamma K > \theta$ .  $\square$

Therefore, we are left to consider the possibility of the patterns bifurcated from  $(u_*, v_*)$  for the Rosenzweig–MacArthur type predator–prey system (5.3) and hence hereafter  $\alpha = 0$ . In this case, the corresponding characteristic equation is

$$\rho^2 + a(D, k^2)\rho + b(D, k^2) = 0, \tag{5.7}$$

with

$$a(D, k^2) = (d(v_*) + D)k^2 - \beta_1, \quad b(D, k^2) = d(v_*)Dk^4 - \beta_2k^2 + \beta_3, \tag{5.8}$$

where

$$\begin{cases} \beta_1 = \frac{\mu v_*(K - \lambda - 2v_*)}{K(\lambda + v_*)}, \\ \beta_2 = \frac{\mu v_*(K - \lambda - 2v_*)d(v_*)}{K(\lambda + v_*)} - \frac{\mu v_*(K - v_*)\chi(v_*)}{K}, \\ \beta_3 = \frac{\lambda\theta\mu(K - v_*)}{K(\lambda + v_*)}. \end{cases} \tag{5.9}$$

If  $K > v_* > \frac{K-\lambda}{2}$ , one has  $K - \lambda - 2v_* < 0$  which yields  $a(D, k^2) > 0$  and  $b(D, k^2) > 0$  for all  $k \geq 0$ . Hence, all the eigenvalues have negative real parts and the homogeneous coexistence steady state  $(u_*, v_*)$  is linearly stable. This implies that the pattern formation is possible only when

$$0 < v_* \leq \frac{K - \lambda}{2}. \tag{5.10}$$

First, if  $v_* = \frac{K-\lambda}{2}$ , one has  $a(D, 0) = 0$  and  $b(D, 0) = \beta_3 > 0$ , which corresponds to the spatially homogeneous periodic solutions. Moreover, for any  $k > 0$ , one has  $a(D, k) > 0$  and  $b(D, k) > 0$ , which indicates that all the eigenvalues of (5.7) have negative real part and hence no spatially inhomogeneous patterns arise in this case. Next, we shall exploit whether the spatially inhomogeneous patterns may arise in the case  $0 < v_* < \frac{K-\lambda}{2}$ .

Define the set

$$\mathcal{H} = \{(D, \eta) \in \mathbb{R}_+^2 : a(D, \eta) = 0\},$$

as the Hopf bifurcation curve, and

$$\mathcal{S} = \{(D, \eta) \in \mathbb{R}_+^2 : b(D, \eta) = 0\},$$

as the steady-state bifurcation curve, where the linearised system around the homogeneous coexistence steady state has an eigenvalue with zero real part on the curves  $\mathcal{S}$  or  $\mathcal{H}$ . Furthermore, if we define

$$D_{\mathcal{H}}(\eta) = \frac{\mu v_*(K - \lambda - 2v_*)}{K(\lambda + v_*)\eta} - d(v_*), \tag{5.11}$$

and

$$D_{\mathcal{S}}(\eta) = \left( \frac{\mu v_*(K - \lambda - 2v_*)}{K(\lambda + v_*)} - \frac{\mu v_*(K - v_*)\chi(v_*)}{d(v_*)K} \right) \frac{1}{\eta} - \frac{\lambda\theta\mu(K - v_*)}{Kd(v_*)(\lambda + v_*)} \frac{1}{\eta^2}, \tag{5.12}$$

then, we have the following stability results for the coexistence steady state  $(u_*, v_*)$ .

**Lemma 5.4** Assume the parameters  $\theta, \gamma, \mu, \lambda$  and  $K > \lambda$  are fixed. Let  $D_{\mathcal{H}}(\eta)$  and  $D_{\mathcal{S}}(\eta)$  be defined in (5.11) and (5.12). Then, the homogeneous coexistence steady state is locally asymptotically stable provided the parameter  $D$  satisfies

$$D > \max_{\eta \geq 0} \{D_{\mathcal{H}}(\eta), D_{\mathcal{S}}(\eta)\}. \tag{5.13}$$

**Remark 5.1** In one dimension, say  $\Omega = [0, \ell]$ ,  $\eta = k^2 = (\frac{n\pi}{\ell})^2$  where  $n = 0, 1, 2, \dots$ , and  $\ell$  is the length of the domain. For fixed  $D > 0$ , the condition (5.13) can be satisfied if  $\ell$  is small enough. Hence, no pattern formation will develop if  $\ell$  is sufficient small.

Next we explore the possible (local) bifurcation by treating  $D$  as a bifurcation parameter. Under (5.10), we see that the sign of  $a(D, k^2)$  and  $b(D, k^2)$  could be generic and different type bifurcation may arise. In particular, the discriminant of (5.7)  $\Delta = |a(D, k^2)|^2 - 4b(D, k^2)$  has not determined sign. Therefore, there are two possibilities:  $\Delta < 0$  and  $\Delta \geq 0$ , where the former will lead to a Hopf bifurcation (periodic patterns) and the latter may lead to a steady-state bifurcation (aggregation patterns). Note the allowable wavenumbers  $k$  are discrete in a bounded domain, for instance, if  $\Omega = (0, \ell)$ , then  $k = \frac{n\pi}{\ell}$  ( $n = 0, 1, 2, \dots$ ). It can be easily verified that

$$\Delta = |a(D, k^2)|^2 - 4b(D, k^2) = (D - d(v_*))^2 k^4 - 2[(D + d(v_*))\beta_1 - 2\beta_2]k^2 + \beta_1^2 - 4\beta_3.$$

Hence, the Hopf bifurcation may occur (i.e.  $\Delta < 0$ ) if  $[(D + d(v_*))\beta_1 - 2\beta_2]^2 - (\beta_1^2 - 4\beta_3)(D - d(v_*))^2 > 0$  and there is allowable wavenumber  $k$  such that

$$k_1^- < k^2 < k_1^+,$$

where  $k_1^\pm = \frac{(D+d(v_*))\beta_1 - 2\beta_2 \pm \sqrt{[(D+d(v_*))\beta_1 - 2\beta_2]^2 - (\beta_1^2 - 4\beta_3)(D-d(v_*))^2}}{(D-d(v_*))^2}$  if  $D \neq d(v_*)$ .

If  $\Delta > 0$ , steady-state bifurcation may occur but the possibility can be generic. Indeed one can readily verify that the steady-state bifurcation will occur if there is allowable wavenumber  $k$  such that one of the following cases holds:

- (i)  $b(D, k^2) < 0$ ;
  - (ii)  $b(D, k^2) = 0, a(D, k^2) < 0$ ;
  - (iii)  $b(D, k^2) > 0, a(D, k^2) < 0$  and  $|a(D, k^2)|^2 - 4b(D, k^2) > 0$ .
- $$\tag{5.14}$$

The corresponding parameter regime guaranteeing each of (i), (ii) and (iii) can be found with easy calculations. For example, under (5.10), it follows that  $\beta_3 > 0$ . Hence, condition (i) is ensured if

$$\beta_2 > 0, \text{ namely } \frac{K - \lambda - 2v_*}{(\lambda + v_*)(K - v_*)} > \frac{\chi(v_*)}{d(v_*)}, \tag{5.15}$$

and the allowable wavenumber  $k$  satisfies

$$k_2^- < k^2 < k_2^+, \quad k_2^\pm = \frac{\beta_2 \pm \sqrt{\Lambda}}{2Dd(v_*)} \text{ with } \Lambda = \beta_2^2 - 4\beta_3Dd(v_*). \tag{5.16}$$

Conditions of ensuring (ii) or (iii) can be derived similarly and will not be detailed here since these conditions can be easily inspected when the parameter values and the motility function  $d(v), \chi(v)$  are specified, as shown in the next subsection.



### 5.3 Spatio-temporal patterns

In this subsection, we shall present some examples to illustrate the periodic and steady-state patterns. As discussed in the previous section, the Lotka–Volterra type predator–prey system (5.1) does not generate any spatially inhomogeneous patterns while the Rosenzweig–MacArthur type predator–prey system (5.3) with  $\chi(v) = -d'(v)$  may generate periodic or steady-state patterns in appropriate parameter regimes. Therefore, we only consider the Rosenzweig–MacArthur type predator–prey system (5.3) with  $\chi(v) = -d'(v)$ . We fix the value of the parameters in all simulations as follows:

$$K = 4, \gamma = 2, \theta = 1, \lambda = 1, \mu = 1. \quad (5.17)$$

Then, it can be checked from (5.2) that the coexistence steady state  $(u_*, v_*) = (3/2, 1)$ . Furthermore, it can be verified from (5.9) that

$$\beta_1 = \frac{1}{8}, \beta_3 = \frac{3}{8},$$

where the value of  $\beta_2$  depends on the specific form of  $d(v)$ . In this paper, we shall test three-motility function  $d(v)$  as follows:

$$d_1(v) = \frac{1}{1 + e^{2(v-1)}}, \quad d_2(v) = \frac{1}{1 + e^{\frac{1}{10}(v-1)}}, \quad d_3(v) = \frac{1}{9 + e^{2(v-1)}}. \quad (5.18)$$

Hence,  $d_1(v_*) = d_2(v_*) = 1/2$ ,  $d_3(v_*) = \frac{1}{10}$  and  $\chi_1(v_*) = -d'_1(v_*) = \frac{1}{2}$ ,  $\chi_2(v_*) = -d'_2(v_*) = \frac{1}{40}$ ,  $\chi_3(v_*) = -d'_3(v_*) = \frac{1}{50}$ . Next we shall numerically explore the possible patterns for different choices of  $d(v)$  given in (5.18).

**Case 1:**  $d(v) = d_1(v)$ . In this case, under the parameters chosen in (5.17), one can verify that

$$a(D, k^2) = \left(\frac{1}{2} + D\right) k^2 - \frac{1}{8}. \quad (5.19)$$

One also can verify from (5.8) that  $b(D, k^2) = \frac{D}{2}k^4 + \frac{5}{16}k^2 + \frac{3}{8} > 0$ ,  $\beta_2 = -\frac{5}{16}$  and

$$\Delta = |a(D, k^2)|^2 - 4b(D, k^2) = \left(\frac{1}{2} - D\right)^2 k^4 - \frac{1}{4} \left(\frac{11}{2} + D\right) k^2 - \frac{95}{64}.$$

This indicates that as long as  $D$  is close to  $1/2$ , then  $\Delta < 0$  and Hopf bifurcation will certainly arise. One is concerned whether the steady-state bifurcation will occur in this case. Indeed it can be readily checked  $a(D, k^2) < 0$  and  $|a(D, k^2)|^2 - 4b(D, k^2) > 0$  cannot be fulfilled simultaneously. Hence from (5.14), we know that the steady-state bifurcation is impossible in this case. However, the Hopf bifurcation will develop if  $D$  is suitably chosen so that  $\Delta < 0$  for some  $k$ . For simulation, we choose  $D = 1/10$  such that  $\Delta < 0$  and  $a(D, k^2) < 0$  with allowable wavenumber satisfying  $k^2 < \frac{5}{24}$  which, under the facts  $k^2 = \left(\frac{n\pi}{\ell}\right)^2$ ,  $n = 0, 1, 2, \dots$  with  $\ell = 8\pi$ , gives  $n = 0, 1, 2, 3$ . The numerical simulations of patterns are then shown in Figure 1(a) and (b) where we observe the spatially homogeneous time-periodic patterns. In principle, there will be three spatial modes arising from the homogeneous coexistence steady state  $(3/2, 1)$ , but we do not observe the spatial inhomogeneity. This implies from the plot in Figure 1(c) that as the solution amplitude become large as time increases, the nonlinearity will play a dominant role and the linearised dynamics is insufficient to explain the nonlinear behaviour.




FIGURE 1. Numerical simulation of spatially homogeneous time-periodic patterns generated by (5.3) with  $\chi(v) = -d'(v)$  in the interval  $[0, 8\pi]$ , where  $d(v) = d_1(v)$  given in (5.18) and parameter values are:  $K = 4, \gamma = 2, \theta = 1, \lambda = 1, \mu = 1, D = 1/10$ . The initial datum  $(u_0, v_0)$  is set as a small random perturbation of the homogeneous coexistence steady state  $(3/2, 1)$ . The simulation illustrates a spatially homogeneous time-periodic coexistence patterns for the predator and the prey.

**Case 2:**  $d(v) = d_2(v)$ . For this case,  $\beta_2 = \frac{7}{160} > 0$  and  $a(D, k^2)$  is still given by (5.19). Hence, the condition (5.15) is verified and

$$\Lambda = \beta_2^2 - 4\beta_3 D d(v_*) = \left(\frac{7}{160}\right)^2 - \frac{3}{4}D.$$

Clearly  $\Lambda > 0$  if  $D < \frac{49}{19200}$  and  $\Lambda \leq 0$  if  $D \geq \frac{49}{19200}$ , which indicates from (5.16) that the steady-state bifurcation will occur if  $0 < D < \frac{49}{19200}$ . This is confirmed by numerical simulations shown in Figure 2 where we take  $D = \frac{1}{4800}$  and observe the development of spatially inhomogeneous




FIGURE 2. Numerical simulation of spatio-temporal patterns generated by (5.3) with  $\chi(v) = -d'(v)$  in the interval  $[0, 4\pi]$ , where  $d(v) = d_2(v)$  given in (5.18) and parameter values are:  $K = 4, \gamma = 2, \theta = 1, \lambda = 1, \mu = 1, D = 1/4800$ . The initial datum  $(u_0, v_0)$  is set as a small random perturbation of the homogeneous coexistence steady state  $(3/2, 1)$ .

stationary patterns (see Figure 2(a) and (b)). Furthermore, both the predator and the prey reach a perfect inhomogeneous coexistence state in space (see Figure 2(c)) but remain oscillations in time (see Figure 2(d)). It has been proved that if  $d(v)$  is constant, the diffusive Rosenzweig–MacArthur predator–prey system (5.3) will not admit spatial patterns (cf. Refs. [47, 48]). The spatially inhomogeneous stationary patterns shown in Figure 2 imply that density-dependent nonlinear motility (i.e. function  $d(v)$ ), which leads to a cross-diffusion motion, is a trigger for pattern formation. This is a new observation although it is not justified in the paper. When  $d(v)$  is constant, the spatial patterns and time-periodic patterns have been obtained for prey-taxis systems with different predator–prey interactions or mobility coefficient  $\chi(v)$ , see Refs. [42, 44]. We also refer to Refs. [15, 27] for some other types of cross-diffusion which cause the emergence of spatial patterns.

**Case 3:**  $d(v) = d_3(v)$ . In this case, one has  $d(v_*) = \frac{1}{10}$  and  $\chi(v_*) = \frac{1}{50}$ . Furthermore,

$$a(D, k^2) = \left( \frac{1}{10} + D \right) k^2 - \frac{1}{8}, \quad b(D, k^2) = \frac{D}{10} k^4 + \frac{1}{400} k^2 + \frac{3}{8},$$




FIGURE 3. Numerical simulation of spatio-temporal patterns generated by (5.3) with  $\chi(v) = -d'(v)$  in the interval  $[0, 8\pi]$ , where  $d(v) = d_3(v)$  given in (5.18) and parameter values are:  $K = 4, \gamma = 2, \theta = 1, \lambda = 1, \mu = 1, D = 1/10$ . The initial datum  $(u_0, v_0)$  is set as a small random perturbation of the coexistence steady state  $(3/2, 1)$ .

and hence

$$\Delta = |a(D, k^2)|^2 - 4b(D, k^2) = \left(\frac{1}{10} - D\right)^2 k^4 - \frac{1}{4} \left(\frac{7}{50} + D\right) k^2 - \frac{95}{64}.$$

Choosing  $D = \frac{1}{10}$ , then  $\Delta < 0$  and  $a(D, k^2) < 0$  with allowable wavenumber  $k^2 < \frac{5}{8}$ . Hence, allowable wave modes are  $n = 0, 1, 2, 3, 4, 5, 6$  by noticing that  $k = \frac{n}{8}$  and Hopf bifurcation (with positive real part in the temporal eigenvalue) will arise. We show the numerical simulation in Figure 3, where we observe the development of chaotic spatio-temporal patterns, which are different from the patterns shown in Figures 1 and 2. They are not the periodic patterns either (see the lower panel of Figure 3) as we expect from the linear stability analysis, which indicates again that the dynamics between nonlinear and linearised systems are quite different. We also note that the simulations in Figures 1 and 3 demonstrate that the Hopf bifurcation arising from the time-periodic orbits can develop into spatially homogeneous time-periodic patterns (Figure 1) or chaotic spatio-temporal patterns (Figure 3). The difference in the simulations shown in Figures

1 and 3 lies in the choice of motility function  $d(v)$ . This observation hints us that the motility function  $d(v)$  of the predator plays an important role in determining the spatial distribution of the predator and the prey. In particular, the random motion ( $d(v)$  is constant) and non-random motion ( $d(v)$  is non-constant) will result in different patterns (i.e. spatial distribution of the predator and the prey). Hence, how does the motility function  $d(v)$  affects the dynamics of nonlinear predator–prey systems launches an interesting question for the future.

### Acknowledgements

The authors are grateful to the three referees and one editor for their valuable comments, which greatly improved the exposition of our paper. The research of H.Y. Jin was supported by the NSF of China (No. 11871226), Guangdong Basic and Applied Basic Research Foundation (No. 2020A1515010140, No. 2020B1515310015), Guangzhou Science and Technology Program (No. 202002030363) and the Fundamental Research Funds for the Central Universities. The research of Z.A. Wang was supported by the Hong Kong RGC GRF grant No. PolyU 153298/16P.

### Conflict of Interest

None.

### References

- [1] ALIKAKOS, N. D. (1979)  $L^p$  bounds of solutions of reaction-diffusion equations. *Comm. Part. Differ. Equations* **4**, 827–868.
- [2] AINSEBA, B. E., BENDAHMANE, M. & NOUSSAIR, A. (2008) A reaction-diffusion system modeling predator-prey with prey-taxis. *Nonlinear Anal. Real World Appl.* **9**(5), 2086–2105.
- [3] AMANN, H. (1989) Dynamic theory of quasilinear parabolic equations III. Global existence. *Math. Z.* **202**, 219–250.
- [4] AMANN, H. (1990) Dynamic theory of quasilinear parabolic equations II. Reaction-diffusion systems. *Differ. Integral Equ.* **3**(1), 13–75.
- [5] AMANN, H. (1992) Nonhomogeneous linear and quasilinear elliptic and parabolic boundary value problems. In: *Function Spaces, Differential Operators and Nonlinear Analysis (Friedrichroda, 1992)*, Vol. 133. Teubner-Texte Math., Teubner, Stuttgart, pp. 9–126.
- [6] CHAKRABORTY, A., SINGH, M., LUCY, D. & RIDLAND, P. (2007) Predator-prey model with prey-taxis and diffusion. *Math. Comp. Mod.* **46**, 482–498.
- [7] CHESSON, P. & MURDOCH, W. (1986) Aggregation of risk: relationships among host-parasitoid models. *Am. Nat.* **127**, 696–715.
- [8] COSNER, C. (2014) Reaction-diffusion-advection models for the effects and evolution of dispersal. *Discrete Contin. Dyn. Syst.* **34**, 1701–1745.
- [9] FU, X., TANG, L.-H., LIU, C., HUANG, J.-D., HWA, T. & LENZ, P. (2012) Stripe formation in bacterial system with density-suppressed motility. *Phys. Rev. Lett.* **108**, 198102.
- [10] GRÜNBAUM, D. (1998) Using spatially explicit models to characterize foraging performance in heterogeneous landscapes. *Am. Nat.* **151**, 97–115.
- [11] HOLLING, C. S. (1965) The functional response of predators to prey density and its role in mimicry and population regulation. *Mem. Entom. Soc. Can.* **45**, 1–60.
- [12] JIN, H. Y. & WANG, Z. A. (2017) Global stability of prey-taxis systems. *J. Differ. Equations* **262**, 1257–1290.
- [13] JIN, H. Y., KIM, Y. J. & WANG, Z. A. (2018) Boundedness, stabilization and pattern formation driven by density-suppressed motility. *SIAM J. Appl. Math.* **78**, 1632–1657.

- [14] JÜNGEL, A. (2010) Diffusive and nondiffusive population models. In: *Mathematical Modeling of Collective Behavior in Socio-Economic and Life Sciences*, Model. Simul. Sci. Eng. Technol., Boston, MA: Birkhäuser Boston, Inc., pp. 397–425.
- [15] JÜNGEL, A., KUEHN, C. & TRUSSARDI, L. (2017) A meeting point of entropy and bifurcations in cross-diffusion herding. *European J. Appl. Math.* **28**(2), 317–356.
- [16] KAREIVA, P. & ODELL, G. T. (1987) Swarms of predators exhibit “preytaxis” if individual predators use area-restricted search. *Am. Nat.* **130**, 233–270.
- [17] KELLER, E. F. & SEGEL, L. A. (1971) Model for chemotaxis. *J. Theor. Biol.* **30**(2), 225–234.
- [18] KELLEY, W. G. & PETERSON, A. C. (2010) *The Theory of Differential Equations - Classical and Qualitative*, Springer.
- [19] KOWALCZYK, R. & SZYMAŃSKA, Z. (2008) On the global existence of solutions to an aggregation model. *J. Math. Anal. Appl.* **343**, 379–398.
- [20] LADYZHENSKAYA, O., SOLONNIKOV, V. & URALCEVA, N. (1968) *Linear and Quasilinear Equations of Parabolic Type*. Providence, RI: AMS.
- [21] LASALLE, J. P. (1960) Some extensions of Lyapunov’s second method. *IRE Trans. Circ. Theo.* **CT-7**, 520–527 (1960).
- [22] LEE, J. M., HILLEN, T. & LEWIS, M. A. (2008) Continuous traveling waves for prey-taxis. *Bull. Math. Biol.* **70**, 654–676.
- [23] LEE, J. M., HILLEN, T. & LEWIS, M. A. (2009) Pattern formation in prey-taxis systems. *J. Biol. Dyn.* **3**(6), 551–573.
- [24] LIU, C., FU, X., LIU, L., REN, X., CHAU, C. K. L., LI, S., ZENG, H., CHEN, G., TANG, L., LENZ, P., CUI, X., HUANG, W., HWA, T. & HUANG, J. (2011) Sequential establishment of stripe patterns in an expanding cell population. *Science* **334**, 238–241.
- [25] LOTKA, A. J. (1925) *Elements of Physical Biology*. Baltimore: Williams and Wilkins Co.
- [26] MA, M., PENG, R. & WANG, Z. (2020) Stationary and non-stationary patterns of the density-suppressed motility model. *Phys. D* **402**, 132259, 13 pages.
- [27] MIMURA, M. & KAWASAKI, K. (1980) Spatial segregation in competitive interaction-diffusion equations. *J. Math. Biol.* **9**(1), 49–64.
- [28] MIZOGUCHI, N. & SOUPLLET, P. (2014) Nondegeneracy of blow-up points for the parabolic Keller-Segel system. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **31**, 851–875.
- [29] MURDOCH, W. W., BRIGGS, C. J. & NISBERT, R. M. (2003) *Consumer-Resource Dynamics (Monographs in Population Biology-36)*. Princeton University Press.
- [30] MURDOCH, W., CHESSON, J. & CHESSON, P. (1985) Biological control in theory and practice. *Am. Nat.* **125**, 344–366.
- [31] OKUBO, A. & LEVIN, S. A. (2001) *Diffusion and Ecological Problems: Modern Perspective*. Interdisciplinary Applied Mathematics, Vol. 14. 2nd ed. Berlin: Springer.
- [32] PORZIO, M. M. & VESPRI, V. (1993) Hölder estimates for local solutions of some doubly nonlinear degenerate parabolic equations. *J. Differ. Equations* **103**(1), 146–178.
- [33] ROSENZWEIG, M. L. & MACARTHUR, R. H. (1963) Graphical representation and stability conditions of predator-prey interactions. *Am. Nat.* **97**, 209–223.
- [34] SAPOUKHINA, N., TYUTYUNOV, Y. & ARDITI, R. (2003) The role of prey taxis in biological control: a spatial theoretical model. *Am. Nat.* **162**, 61–76.
- [35] SMITH-ROBERGE, J., IRON, D. & KOLOKOLNIKOV, T. (2019) Pattern formation in bacterial colonies with density-dependent diffusion. *European J. Appl. Math.* **30**(1), 196–218.
- [36] SOUPLLET, P. & QUITTNER, P. (2007) *Superlinear Parabolic Problems: Blow-up, Global Existence and Steady States*. Basel, Boston, Berlin: Birkhäuser Advanced Texts.
- [37] TAO, Y. S. (2010) Global existence of classical solutions to a predator-prey model with nonlinear prey-taxis. *Nonlinear Anal. Real World Appl.* **11**(3), 2056–2064.
- [38] TAO, Y. S. & WINKLER, M. (2011) A chemotaxis-haptotaxis model: the roles of nonlinear diffusion and logistic source. *SIAM J. Math. Anal.* **43**(2), 685–704.
- [39] TAO, Y. S. & WINKLER, M. (2012) Boundedness in a quasilinear parabolic-parabolic Keller-Segel system with subcritical sensitivity. *J. Differ. Equations* **252**(1), 692–715.

- [40] TURCHIN, P. (2003) *Complex Population Dynamics: A Theoretical/Empirical Synthesis*. Monographs in Population Biology-35. Princeton University Press.
- [41] VOLTERRA, V. (1926) Fluctuations in the abundance of a species considered mathematically. *Nature* **118**, 558–560.
- [42] WANG, K., WANG, Q. & YU, F. (2017) Stationary and time periodic patterns of two predator and one-prey systems with prey-taxis. *Discrete. Contin. Dyn. Syst.* **37**(1), 505–543.
- [43] WANG, M. X. (2018) Note on the Lyapunov functional method. *Appl. Math. Lett.* **75**, 102–107.
- [44] WANG, Q., SONG, Y. & SHAO, L. J. (2017) Nonconstant positive steady states and pattern formation of 1D prey-taxis systems. *J. Nonlinear Sci.* **27**(1), 71–97.
- [45] WANG, J. P. & WANG, M. X. (2018) Boundedness and global stability of the two-predator and one-prey models with nonlinear prey-taxis. *Z. Angew. Math. Phys.* **69**(3), 63, 24 pages.
- [46] WU, S., SHI, J. P. & WU, B. (2016) Global existence of solutions and uniform persistence of a diffusive predator-prey model with prey-taxis. *J. Differ. Equations* **260**(7), 5847–5874.
- [47] WU, S., WANG, J. & SHI, J. (2018) Dynamics and pattern formation of a diffusive predator-prey model with predator-taxis. *Math. Models Method Appl. Sci.* **28**(11), 2275–2312.
- [48] YI, F., WEI, J. & SHI, J. (2009) Bifurcation and spatiotemporal patterns in a homogeneous diffusive predator-prey system. *J. Differ. Equations* **246**, 1944–1977.