

Global classical solutions for a class of reaction-diffusion system with density-suppressed motility

Wenbin Lyu ^{*} Zhi-An Wang [†]

Abstract. This paper is concerned with a class of reaction-diffusion system with density-suppressed motility

$$\begin{cases} u_t = \Delta(\gamma(v)u) + \alpha u F(w), & x \in \Omega, \quad t > 0, \\ v_t = D\Delta v + u - v, & x \in \Omega, \quad t > 0, \\ w_t = \Delta w - u F(w), & x \in \Omega, \quad t > 0, \end{cases}$$

under homogeneous Neumann boundary conditions in a smooth bounded domain $\Omega \subset \mathbb{R}^n$ ($n \leq 2$), where $\alpha > 0$ and $D > 0$ are constants. The random motility function γ satisfies

$$\gamma \in C^3((0, +\infty)), \quad \gamma > 0, \quad \gamma' < 0 \quad \text{on} \quad (0, +\infty) \quad \text{and} \quad \lim_{v \rightarrow +\infty} \gamma(v) = 0.$$

The intake rate function F satisfies

$$F \in C^1([0, +\infty)), \quad F(0) = 0 \quad \text{and} \quad F > 0 \quad \text{on} \quad (0, +\infty).$$

We show that the above system admits a unique global classical solution for all non-negative initial data

$$u_0 \in C^0(\bar{\Omega}), \quad v_0 \in W^{1,\infty}(\Omega), \quad w_0 \in W^{1,\infty}(\Omega).$$

Moreover, if there exist $k > 0$ and $\bar{v} > 0$ such that

$$\inf_{v > \bar{v}} v^k \gamma(v) > 0,$$

then the global solution is bounded uniformly in time.

Keywords: Reaction-diffusion system, Density-suppressed motility, Global existence, Boundedness.

2020 Mathematics Subject Classification.: 35A01, 35B45, 35K51, 35Q92.

1 Introduction and main results

To explain the strip pattern formation observed in the experiment of [21] induced by the “self-trapping mechanism”, the following three-component reaction-diffusion system with density-dependent motility was proposed in [21]

$$\begin{cases} u_t = \Delta(\gamma(v)u) + \frac{\alpha w^2 u}{w^2 + \lambda}, & x \in \Omega, \quad t > 0, \\ v_t = D\Delta v + u - v, & x \in \Omega, \quad t > 0, \\ w_t = \Delta w - \frac{w^2 u}{w^2 + \lambda}, & x \in \Omega, \quad t > 0, \end{cases} \quad (1.1)$$

^{*}School of Mathematical Sciences, Shanxi University, Taiyuan 030006, P.R. China; lvwenbin@sxu.edu.cn.

[†]Department of Applied Mathematics, The Hong Kong Polytechnic University, Hong Kong; mawza@polyu.edu.hk.

where $u(x, t), v(x, t), w(x, t)$ denote the bacterial cell density, concentration of acyl-homoserine lactone (AHL) and nutrient density, respectively; $\alpha, \lambda, D > 0$ are constants and Ω is a bounded domain in \mathbb{R}^n . The first equation of (1.1) describes the random motion of bacterial cells with AHL-density dependent motility coefficient $\gamma(v)$, and cell growth due to the nutrient intake. The second equation of (1.1) describes the diffusion, production and turnover of AHL, while the third equation gives the dynamics of the nutrient with diffusion and consumption. Simultaneously a simplified two-component system was discussed in the supplemental material of [21] and formally analyzed in [7]:

$$\begin{cases} u_t = \Delta(\gamma(v)u) + \mu u(1 - u), & x \in \Omega, t > 0, \\ v_t = D\Delta v + u - v, & x \in \Omega, t > 0, \end{cases} \quad (1.2)$$

where the decay of bacterial cells at high density was used to approximate the nutrient depletion effect. A striking feature of systems (1.1) and (1.2) is that the cell diffusion rate depends on a motility function $\gamma(v)$ satisfying $\gamma'(v) < 0$, which takes into account the repressive effect of AHL concentration on the cell motility (cf. [21]). The density-suppressed motility mechanism has also been used to model other biological processes, preytaxis [16, 14] and chemotaxis [17, 36]. From the expansion

$$\Delta(\gamma(v)u) = \nabla \cdot (\gamma(v)\nabla u + u\gamma'(v)\nabla v) = \gamma(v)\Delta u + 2\gamma'(v)\nabla v \cdot \nabla u + u\gamma''(v)|\nabla v|^2 + u\gamma'(v)\Delta v,$$

we see that the nonlinear diffusion rate function $\gamma(v)$ not only contributes a cross-diffusion structure but also renders a possible diffusion degeneracy (i.e., $\gamma(v) \rightarrow 0$ as $v \rightarrow +\infty$). Therefore many conventional methods are inapplicable and the analysis of (1.1) or (1.2) is very delicate. The progresses were not made to the system (1.2) until recently with homogeneous Neumann boundary conditions in a smooth bounded domain $\Omega \subset \mathbb{R}^n$. The existing results on (1.2) can be classified into two cases: $\mu > 0$ and $\mu = 0$, to be recalled below.

When $\mu > 0$, the global existence and asymptotic behavior of solutions was first established in [11] under certain conditions on $\gamma(v)$ in two dimensions, followed by a series of works [35, 20, 10, 15]. For small $\mu > 0$, the existence/nonexistence of nonconstant steady states of (1.2) was rigorously established in [28, 37] in appropriate parameter regimes. Some other works with generalized logistic source or indirect production of chemical signals can be found in [24, 25, 26, 27]. When $\mu = 0$, the global well-posedness of solutions is more delicate. If $\gamma(v)$ decays algebraically in v , the solution may exist globally in two or higher dimensions [38, 6, 1]. While if $\gamma(v)$ decays exponentially, the solution may blow-up in two dimensions with a critical mass [13, 9, 10]. The global weak solution large initial data was studied in [6, 5]. Except the studies on the bounded domain with zero Neumann boundary conditions, there are some results obtained in the whole space \mathbb{R} : when $\gamma(v)$ is a piecewise constant function, the dynamics of discontinuity interface was studied in [30] and discontinuous traveling wave solutions of (1.2) with $\mu > 0$ were constructed in [23]; the existence of smooth traveling wave solutions of (1.2) with $\mu > 0$ and a particular motility function $\gamma(v) = 1/(1 + v)^m (m > 0)$ was recently shown in [19].

Compared to the abundant results recently obtained for the system (1.2) as recalled above, the progress made to the three-component system (1.1) is very limited. The purpose of this paper is to explore the global well-posedness of the following system

$$\begin{cases} u_t = \Delta(\gamma(v)u) + \alpha u F(w), & x \in \Omega, \quad t > 0, \\ v_t = D\Delta v + u - v, & x \in \Omega, \quad t > 0, \\ w_t = \Delta w - u F(w), & x \in \Omega, \quad t > 0, \\ \frac{\partial u}{\partial \nu} = \frac{\partial v}{\partial \nu} = \frac{\partial w}{\partial \nu} = 0, & x \in \partial\Omega, \quad t > 0, \\ (u, v, w)(x, 0) = (u_0, v_0, w_0)(x), & x \in \Omega, \end{cases} \quad (1.3)$$

with constants $\alpha > 0$ and $D > 0$, where the system (1.1) is a special case of (1.3) with $F(w) = \frac{w^2}{w^2 + \lambda}$. By postulating that

$$\gamma(v) \in C^3([0, +\infty)) \text{ and } 0 < \gamma_1 \leq \gamma(v) \leq \gamma_2, |\gamma'(v)| < \eta \text{ on } [0, +\infty) \quad (1.4)$$

where γ_1, γ_2 are positive constants, and

$$F \in C^1([0, +\infty)), F(0) = 0 \text{ and } F(w) > 0, F'(w) > 0 \text{ on } (0, +\infty),$$

a recent work [12] showed that the problem (1.3) admits a global classical solution (u, v, w) which asymptotically converges to $(u_*, u_*, 0)$ in L^∞ with $u_* = \frac{1}{|\bar{\Omega}|} (\|u_0\|_{L^1} + \alpha \|w_0\|_{L^1})$ if $D > 0$ is suitably large. The main approaches employed in [12] to establish the global classical solutions with uniform-in-time bounds are based on the method of energy estimates and Moser iteration by fully capturing the diffusive dissipation of u with the assumption that $\gamma(v)$ has a positive lower bound. The assumption (1.4) bypasses the possible diffusion degeneracy/singularity and rules out a large class of functions such as $\gamma(v) = \frac{c_0}{v^k} (c_0, k > 0)$ and $\gamma(v) = e^{-\chi v} (\chi > 0)$ widely studied in the existing works as recalled above. The goal of this paper is to remove this essential restriction imposed in (1.4) and establish the global well-posedness of solutions to (1.3). Roughly speaking, under the following structural assumptions on $\gamma(v)$ and $F(v)$:

$$\gamma(v) \in C^3([0, +\infty)), \gamma(v) > 0, \gamma'(v) < 0 \text{ on } (0, +\infty), \text{ and } \lim_{v \rightarrow +\infty} \gamma(v) = 0, \quad (1.5)$$

and

$$F \in C^1([0, +\infty)), F(0) = 0 \text{ and } F > 0 \text{ on } (0, +\infty), \quad (1.6)$$

then for any initial data (u_0, v_0, w_0) satisfying

$$\begin{aligned} u_0 &\in C^0(\bar{\Omega}), v_0 \in W^{1,\infty}(\Omega), w_0 \in W^{1,\infty}(\Omega), \\ u_0 &\geq 0, v_0 \geq c_0 > 0, w_0 \geq 0 \text{ and } u_0 \not\equiv 0, \end{aligned} \quad (1.7)$$

with some constant $c_0 > 0$, we show the problem (1.3) admits a unique global classical solution in two dimensions. Moreover if there exist $k > 0$ and $\bar{v} > 0$ such that

$$\inf_{v > \bar{v}} v^k \gamma(v) > 0 \quad (1.8)$$

the solution is uniformly bounded in time.

Our main results are precisely stated as follows.

Theorem 1.1. *Let $\Omega \subset \mathbb{R}^n (n \leq 2)$ be a bounded domain with smooth boundary. Assume that the conditions (1.5) and (1.6) hold. Then for any initial data (u_0, v_0, w_0) satisfying the condition (1.7), there exists a triple (u, v, w) of non-negative functions*

$$(u, v, w) \in [C^0(\bar{\Omega} \times [0, +\infty)) \cap C^{2,1}(\bar{\Omega} \times (0, +\infty))]^3$$

which solves (1.3) in the classical sense. Moreover, if the motility function γ satisfies the condition (1.8), then the global solution is uniformly bounded in time, that is there exists a constant $C > 0$ such that

$$\|u(\cdot, t)\|_{L^\infty} + \|v(\cdot, t)\|_{W^{1,\infty}} + \|w(\cdot, t)\|_{W^{1,\infty}} \leq C \text{ for all } t > 0.$$

The key of proving Theorem 1.1 is to derive that v has a positive lower bound to rule out the diffusion singularity and has an upper bound to exclude the diffusion degeneracy (see section 3.3.) The positive lower bound of v can be obtained easily by showing that $\int_{\Omega} u dx$ has a positive lower bound along with a nice result of [8]. The crucial step is to show that v has an upper bound. Inspired by an idea from the work [10], we construct an auxiliary problem and use the maximum principle for the inverse operator $(I - D\Delta)^{-1}$ to derive an upper bound of v through the constructed auxiliary problem.

The rest of this paper is organized as follows. Section 2 is devoted to the local existence of solutions and extensibility of (1.3). With some important inequalities which will be used frequently, we derive a priori estimates of solutions for the system (1.3) in section 3. Finally, we prove Theorem 1.1 in section 4.

2 Preliminaries

In this section, we present some basic results and facts, including local existence and extensibility criterion of classical solutions as well as some frequently used well-known inequalities.

The existence of local solutions and extensibility criterion for the system (1.3) can be obtained by Amann's theorem (cf. [3]) or fixed point theorem (cf. [11]). Below, we only state the local existence result without proof.

Lemma 2.1 (Local existence). *Let $\Omega \subset \mathbb{R}^n$ be a bounded domain with smooth boundary. If the initial data satisfy the condition (1.7), then there exist a constant $T_{max} \in (0, \infty]$ and a triple (u, v, w) of non-negative functions*

$$(u, v, w) \in [C^0(\bar{\Omega} \times [0, T_{max})) \cap C^{2,1}(\bar{\Omega} \times (0, T_{max}))]^3,$$

which solves (1.3) in the classical sense in $\Omega \times (0, T_{max})$. Moreover, if $T_{max} < +\infty$, then

$$\limsup_{t \nearrow T_{max}} (\|u(\cdot, t)\|_{L^\infty} + \|v(\cdot, t)\|_{W^{1,\infty}} + \|w(\cdot, t)\|_{W^{1,\infty}}) = \infty.$$

Next, we recall some well-known results which will be used later frequently. The first one is an ODE inequality [31].

Lemma 2.2. *Let $T_{max} > 0$, $\tau \in (0, T_{max})$, $a > 0$ and $b > 0$. Suppose that $y : [0, T_{max}) \rightarrow [0, \infty)$ is absolutely continuous and satisfies*

$$y'(t) + ay(t) \leq h(t) \quad \text{for all } t \in (0, T_{max})$$

with some nonnegative function $h \in L^1_{loc}([0, T_{max}))$ fulfilling

$$\int_t^{t+\tau} h(s) ds \leq b \quad \text{for all } t \in [0, T_{max} - \tau).$$

Then it follows that

$$y(t) \leq \max \left\{ y(0) + b, \frac{b}{a\tau} + 2b \right\} \quad \text{for all } t \in [0, T_{max}).$$

Below is an uniform Grönwall inequality [34] which can help us to derive the uniform-in-time estimates of solutions.

Lemma 2.3. *Let $T_{max} > 0$, $\tau \in (0, T_{max})$. Suppose that a, b, y are three positive locally integrable functions on $(0, T_{max})$ such that y' is locally integrable on $(0, T_{max})$ and the following inequalities are satisfied:*

$$y'(t) \leq a(t)y(t) + b(t) \quad \text{for all } t \in (0, T_{max})$$

as well as

$$\int_t^{t+\tau} a \leq a_1, \quad \int_t^{t+\tau} b \leq a_2, \quad \int_t^{t+\tau} y \leq a_3 \quad \text{for all } t \in [0, T_{max} - \tau),$$

where $a_i (i = 1, 2, 3)$ are positive constants. Then

$$y(t) \leq \left(\frac{a_3}{\tau} + a_2 \right) e^{a_1} \quad \text{for all } t \in [\tau, T_{max}).$$

The third one is about the regularity of solutions to the linear parabolic equation and the proof can be found in [18].

Lemma 2.4. *Assume that $\Omega \subset \mathbb{R}^n$ is a bounded domain with smooth boundary. Suppose that $y(x, t) \in C^{2,1}(\bar{\Omega} \times (0, T_{max}))$ is the solution of*

$$\begin{cases} y_t = \Delta y - y + \phi(x, t), & x \in \Omega, t \in (0, T_{max}), \\ \frac{\partial y}{\partial \nu} = 0, & x \in \partial\Omega, t \in (0, T_{max}), \\ y(x, t) = y_0(x) \in C^0(\bar{\Omega}), \end{cases}$$

where $\phi(x, t) \in L^\infty((0, T_{max}); L^p(\Omega))$. Then there exists a constant $C > 0$ such that

$$\|y(\cdot, t)\|_{W^{1,q}} \leq C \quad \text{for all } t \in (0, T_{max})$$

with

$$q \in \begin{cases} [1, \frac{np}{n-p}), & \text{if } p \leq n, \\ [1, \infty], & \text{if } p > n. \end{cases}$$

3 A priori estimates

This section is devoted to deriving a priori estimates of solutions for the system (1.3), so that the global existence of solutions can be obtained by the extensibility criterion. We will proceed in several steps. In the following, we shall use $C_i (i = 1, 2, \dots)$ to denote a generic positive constant which may vary in the context.

3.1 The boundedness of u in $L^1(\Omega)$

A basic property of solutions is the uniform-in-time L^1 boundedness of u due to the special structure of the system (1.3).

Lemma 3.1. *Let (u, v, w) be a solution of (1.3). Then there exist constants $c, C > 0$ such that*

$$c \leq \int_{\Omega} u \leq C \quad \text{for all } t \in (0, T_{max}). \quad (3.1)$$

Proof. Integrating the first equation of (1.3) over Ω with the boundary conditions, we have $\frac{d}{dt} \int_{\Omega} u \geq 0$ which implies

$$\int_{\Omega} u \geq \int_{\Omega} u_0.$$

We multiply the third equation of (1.3) by α and add the resulting equation to the first equation of (1.3). Then integrating the result over Ω by parts along with the boundary conditions, we get

$$\frac{d}{dt} \left(\int_{\Omega} u + \alpha \int_{\Omega} w \right) = 0$$

which yields

$$\int_{\Omega} u + \alpha \int_{\Omega} w = \int_{\Omega} u_0 + \alpha \int_{\Omega} w_0.$$

Then, the non-negativity of u and w yields (3.1). \square

3.2 The upper bound of w

The following lemma concerns the upper bound of w which is an immediate consequence of the maximum principle (see [12]).

Lemma 3.2. *Let (u, v, w) be a solution of (1.3). We can find a constant $C > 0$ such that*

$$w \leq C \text{ for any } (x, t) \in \Omega \times (0, T_{max}).$$

3.3 The lower and upper bounds of v

The following lemma is vital for us to rule out the possible singularity of $\gamma(v)$ near $v = 0$. The mass inequality (3.1) plays a key role in the proof of this lemma. The proof can be found in [8].

Lemma 3.3. *Let (u, v, w) be a solution of (1.3). Then there exists a constant $C > 0$ fulfilling*

$$v \geq C \text{ for any } (x, t) \in \Omega \times (0, T_{max}).$$

Motivated from the paper [9, 10], next, we derive the upper bound of v , which is a key to preclude the degeneracy of diffusion.

Let us introduce an auxiliary function g which satisfies the following equation

$$\begin{cases} -D\Delta g + g = u, & x \in \Omega, t \in [0, T_{max}), \\ \frac{\partial g}{\partial \nu} = 0, & x \in \partial\Omega, t \in [0, T_{max}), \\ g(x, 0) = g_0(x) \geq 0, & x \in \Omega, \end{cases} \quad (3.2)$$

where u is the solution of (1.3) obtained in Lemma 2.1. Then, g is non-negative since $u \geq 0$ and has the following basic properties. The first property states that the L^q norm of g can be controlled by the L^1 norm of u (cf. [4]).

Lemma 3.4. *Let $u \in C(\overline{\Omega})$ be a non-negative function such that $\int_{\Omega} u > 0$. If $g \in C^2(\overline{\Omega})$ is a solution of the system (3.2), then for any q satisfying $1 \leq q < \frac{n}{(n-2)_+}$ there exists a constant $C > 0$ such that*

$$\|g\|_{L^q} \leq C \|u\|_{L^1}.$$

The second property tells us that g satisfies a simple inequality.

Lemma 3.5. *Let (u, v, w) be a solution of (1.3) and g satisfies (3.2). Then for all $(x, t) \in \Omega \times (0, T_{max})$, we have*

$$g_t + \frac{1}{D}\gamma(v)u = \frac{1}{D}(I - D\Delta)^{-1}[\gamma(v)u] + \alpha(I - D\Delta)^{-1}[uF(w)]. \quad (3.3)$$

Moreover, there exists a constant $C > 0$ such that

$$g_t \leq Cg \quad \text{for any } (x, t) \in \Omega \times (0, T_{max}). \quad (3.4)$$

Proof. The first equation of (1.3) can be rewritten as

$$u_t = -\frac{1}{D}(I - D\Delta)[\gamma(v)u] + \alpha uF(w) + \frac{1}{D}\gamma(v)u.$$

Taking the operator $(I - D\Delta)^{-1}$ on both side of the above equation and noticing the definition of g , we can get (3.3) directly.

Now we prove (3.4). According to the non-increasing property of γ and Lemma 3.3, there exists a constant $C_1 > 0$ such that

$$\gamma(v) \leq C_1.$$

Noticing Lemma 2.1, Lemma 3.2 and the smoothing property of F , we get a constant $C_2 > 0$ such that

$$F(w) \leq C_2.$$

Owing to the nonnegativity of u , it holds that

$$\gamma(v)u \geq 0.$$

Recall (3.2). Then by the comparison principle for elliptic equations, we have

$$\frac{1}{D}(I - D\Delta)^{-1}[\gamma(v)u] + \alpha(I - D\Delta)^{-1}[uF(w)] \leq \left(\frac{C_1}{D} + \alpha C_2\right)g,$$

which, combined with (3.3), implies that

$$g_t \leq \left(\frac{C_1}{D} + \alpha C_2\right)g.$$

This finishes the proof. □

With the help of Lemma 3.5 and the standard comparison principle for parabolic equations, we shall derive the upper bound of v .

Lemma 3.6. *Let (u, v, w) be a solution of (1.3) and g satisfies (3.2). Then there is a constant $C > 0$ such that*

$$v \leq C(g + 1) \quad \text{for any } (x, t) \in \Omega \times (0, T_{max}).$$

Moreover, if $T_{max} < +\infty$, there exists $C_0 > 0$ such that

$$v \leq C_0 \quad \text{for any } (x, t) \in \Omega \times (0, T_{max}).$$

Proof. With the hypothesis (1.5), we can choose a constant $C_1 \geq 0$ large enough such that

$$0 < \gamma(C_1) < D.$$

Let

$$\Gamma(s) := \frac{1}{D} \int_{C_1}^s \gamma(x) dx \quad \text{for all } s \geq 0,$$

which gives

$$\begin{aligned} \gamma(v)u &= \gamma(v)(v_t - D\Delta v + v) \\ &= D\Gamma_t(v) - D^2\Delta\Gamma(v) + D\gamma'(v)|\nabla v|^2 + \gamma(v)v. \end{aligned}$$

This, combined with Lemma 3.5, implies

$$\begin{aligned} v_t - D\Delta v + v &= -D\Delta g + g \\ &= g_t - D\Delta g + g + \frac{1}{D}\gamma(v)u \\ &\quad - \frac{1}{D}(I - D\Delta)^{-1}[\gamma(v)u] - \alpha(I - D\Delta)^{-1}[uF(w)] \\ &= (g + \Gamma(v))_t - D\Delta(g + \Gamma(v)) + (g + \Gamma(v)) + \gamma'(v)|\nabla v|^2 + \frac{1}{D}\gamma(v)v - \Gamma(v) \\ &\quad - \frac{1}{D}(I - D\Delta)^{-1}[\gamma(v)u] - \alpha(I - D\Delta)^{-1}[uF(w)]. \end{aligned} \tag{3.5}$$

Now, we estimate the terms on the right hand side of (3.5). In view of the monotone decreasing property of γ , Lemma 3.3 and the definition of Γ , we see that there exists a constant $C_2 > 0$ such that

$$\begin{aligned} \frac{1}{D}\gamma(v)v - \Gamma(v) &= \frac{1}{D}\gamma(v)v + \frac{1}{D} \int_v^{C_1} \gamma(x) dx \\ &\leq \frac{1}{D} [\gamma(v)v + \gamma(v)(C_1 - v)] \\ &= \frac{C_1}{D}\gamma(v) \leq \frac{C_2}{D} \quad \text{for } 0 \leq v \leq C_1 \end{aligned}$$

or otherwise

$$\begin{aligned} \frac{1}{D}\gamma(v)v - \Gamma(v) &= \frac{1}{D}\gamma(v)v - \frac{1}{D} \int_{C_1}^v \gamma(x) dx \\ &\leq \frac{1}{D} [\gamma(v)v - \gamma(v)(v - C_1)] \\ &= \frac{C_1}{D}\gamma(v) \leq \frac{C_2}{D} \quad \text{for } v \geq C_1. \end{aligned}$$

Due to the non-negativity of $-\gamma'(v)$, $\gamma(v)u$ as well as $uF(w)$ and the comparison principle for elliptic equations, we get from (3.5)

$$v_t - D\Delta v + v \leq (g + \Gamma(v))_t - D\Delta(g + \Gamma(v)) + (g + \Gamma(v)) + \frac{C_2}{D}.$$

Noticing for the initial data, we can choose a constant $C_3 > 0$ large enough such that $\frac{C_2}{D} \leq C_3$ and

$$v_0 \leq g_0 + \Gamma(v_0) + C_3.$$

Hence, the comparison principle for parabolic equations gives that

$$v \leq g + \Gamma(v) + C_3,$$

which along with the fact

$$\Gamma(v) \leq \frac{\gamma(C_1)}{D}v,$$

implies

$$v \leq \frac{1}{1 - \frac{\gamma(C_1)}{D}}(g + C_3).$$

With the aid of Lemma 3.5, if $T_{max} < +\infty$, then there exists a constant $C_4 > 0$ such that

$$v \leq C_4.$$

Hence we complete the proof of this lemma. \square

Note the upper bound derived in Lemma 3.6 may depend on T_{max} , see (3.4). The following lemma asserts the upper bound of v which is independent of T_{max} under additional condition (1.8).

Lemma 3.7. *Let (u, v, w) be a solution of (1.3). If the motility function γ satisfies the condition (1.8), then there exists a constant $C > 0$ independent of T_{max} such that*

$$v \leq C \text{ for any } (x, t) \in \Omega \times (0, T_{max}).$$

Proof. We can rewrite the first equation of (1.3) as

$$((I - D\Delta)g)_t + \frac{1}{D}(I - D\Delta)(\gamma(v)u) = \frac{1}{D}\gamma(v)u + \alpha uF(w).$$

Multiplying the above equation by $g = (I - D\Delta)^{-1}u$ and integrating the result over Ω , we have

$$\frac{1}{2} \frac{d}{dt} \left(\int_{\Omega} g^2 + D \int_{\Omega} |\nabla g|^2 \right) + \frac{1}{D} \int_{\Omega} \gamma(v)u^2 = \frac{1}{D} \int_{\Omega} \gamma(v)ug + \alpha \int_{\Omega} uF(w)g. \quad (3.6)$$

In view of the assumption (1.5) and Lemma 3.3, we get $C_1 > 0$ fulfilling

$$\gamma(v) \leq C_1. \quad (3.7)$$

Noticing Lemma 2.1, Lemma 3.2 and the smoothing property of F , we get a constant $C_2 > 0$ such that

$$F(w) \leq C_2. \quad (3.8)$$

Substituting (3.7) and (3.8) into (3.6), we obtain from Lemma 3.1 that

$$\frac{1}{2} \frac{d}{dt} \left(\int_{\Omega} g^2 + D \int_{\Omega} |\nabla g|^2 \right) + \frac{1}{D} \int_{\Omega} \gamma(v)u^2 \leq \left(\frac{C_1}{D} + \alpha C_2 \right) C_3 \|g\|_{L^\infty}, \quad (3.9)$$

holds for some constant $C_3 > 0$. Moreover, it follows from (3.2) that

$$D \int_{\Omega} |\nabla g|^2 + \int_{\Omega} g^2 = \int_{\Omega} ug \leq C_3 \|g\|_{L^\infty}$$

which, added to (3.9) yields

$$\begin{aligned} & \frac{d}{dt} \left(\int_{\Omega} g^2 + D \int_{\Omega} |\nabla g|^2 \right) + \left(\int_{\Omega} g^2 + D \int_{\Omega} |\nabla g|^2 \right) + \frac{2}{D} \int_{\Omega} \gamma(v) u^2 \\ & \leq 2 \left(\frac{C_1}{D} + \alpha C_2 + 1 \right) C_3 \|g\|_{L^\infty}. \end{aligned} \quad (3.10)$$

Now we estimate the right hand side of the above inequality. For any $\max\{\frac{n}{2}, 1\} < p < 2$, thanks to the Sobolev embedding theorem, the standard elliptic estimate and Hölder's inequality, we can find constants $C_4, C_5, C_6 > 0$ such that

$$\begin{aligned} \|g\|_{L^\infty} & \leq C_4 \|g\|_{W^{2,p}} \leq C_5 \|u\|_{L^p} \\ & \leq \frac{1}{2D} \frac{1}{\left(\frac{C_1}{D} + \alpha C_2 + 1\right) C_3} \int_{\Omega} \gamma(v) u^2 + C_6 \left(\int_{\Omega} \gamma^{-\frac{p}{2-p}}(v) \right)^{\frac{2-p}{p}}. \end{aligned}$$

In view of the assumption (1.8), there exist $k > 0, \bar{v} > 0$ and $C_7 > 0$ such that

$$v^k \gamma(v) \geq C_7 \quad \text{for all } v > \bar{v}$$

i.e.

$$\gamma^{-1}(v) \leq C_7^{-1} v^k \quad \text{for all } v > \bar{v}.$$

Noticing the non-increasing property of γ , we get

$$\gamma^{-1}(v) \leq \gamma^{-1}(\bar{v}) \quad \text{for all } 0 \leq v \leq \bar{v}.$$

Therefore, it holds that

$$\gamma^{-1}(v) \leq \gamma^{-1}(\bar{v}) + C_7^{-1} v^k \quad \text{for all } v \geq 0.$$

Hence, using Lemma 3.6 and Lemma 3.4, there exist constants $C_8, C_9, C_{10} > 0$ such that

$$\begin{aligned} \int_{\Omega} \gamma^{-\frac{p}{2-p}}(v) & \leq \int_{\Omega} \left(\gamma^{-1}(\bar{v}) + C_7^{-1} v^k \right)^{\frac{p}{2-p}} \\ & \leq \int_{\Omega} \left(\gamma^{-1}(\bar{v}) + C_7^{-1} (C_8(g+1))^k \right)^{\frac{p}{2-p}} \\ & \leq C_9 \int_{\Omega} g^{\frac{pk}{2-p}} dx + C_9 \\ & \leq C_{10} \end{aligned} \quad (3.11)$$

which implies that

$$\|g\|_{L^\infty} \leq \frac{1}{2D} \frac{1}{\left(\frac{C_1}{D} + \alpha C_2 + 1\right) C_3} \int_{\Omega} \gamma(v) u^2 + C_6 C_{10}^{\frac{2-p}{p}}. \quad (3.12)$$

Combining (3.10), (3.11) with (3.12), we get

$$\begin{aligned} & \frac{d}{dt} \left(\int_{\Omega} g^2 + D \int_{\Omega} |\nabla g|^2 \right) + \left(\int_{\Omega} g^2 + D \int_{\Omega} |\nabla g|^2 \right) + \frac{1}{D} \int_{\Omega} \gamma(v) u^2 \\ & \leq 2 \left(\frac{C_1}{D} + \alpha C_2 + 1 \right) C_3 C_6 C_{10}^{\frac{2-p}{p}} \end{aligned}$$

which along with Grönwall's inequality yields a constant $C_{11} > 0$ such that

$$\int_{\Omega} g^2 + D \int_{\Omega} |\nabla g|^2 \leq C_{11}$$

and

$$\int_t^{t+\tau} \int_{\Omega} \gamma(v) u^2 \leq C_{11} \quad \text{for all } t \in (0, T_{max} - \tau) \quad (3.13)$$

with $\tau = \min\{1, \frac{1}{2}T_{max}\}$. Due to (3.12) and (3.13), the following inequality

$$\int_t^{t+\tau} \int_{\Omega} g \leq |\Omega| \int_t^{t+\tau} \|g\|_{L^\infty} \leq C_{12} \quad \text{for all } t \in (0, T_{max} - \tau), \quad (3.14)$$

holds for some constant $C_{12} > 0$. According to Lemma 3.5, we can find a constant $C_{13} > 0$ such that

$$g_t \leq C_{13}g \quad \text{for all } t \in (0, T_{max}).$$

Using Lemma 2.3 with (3.14) and the definition of τ , we get a constant $C_{14} > 0$ so that

$$g \leq C_{14} = \frac{C_{12}}{|\Omega|\tau} e^{C_{13}} \quad \text{for any } (x, t) \in \Omega \times (\tau, T_{max})$$

which, along with Lemma 3.5 applied to any $(x, t) \in \Omega \times [0, \tau]$, asserts that

$$g \leq C_{15} \quad \text{for any } (x, t) \in \Omega \times [0, T_{max})$$

holds for some constant $C_{15} > 0$. This completes the proof by using Lemma 3.6. \square

3.4 L^∞ -estimates of u

Once we get the positive lower and upper bounds of v in the previous section, then the diffusion motility function $\gamma(v)$ is neither degenerate nor singular and the estimate of L^∞ -norm of u essentially can be derived by the procedures as shown in [12]. For completeness, we briefly demonstrate the main steps below.

3.4.1 The space-time L^2 -bound of u

In this subsection, we aim to derive the bound of u in space-time L^2 -norm by the classical duality-based arguments (cf. [12, 22, 33]). For convenience, we introduce some notations here. Let A_0 denote the self-adjoint realization of $-\Delta$ defined in the Hilbert space

$$L^2_{\perp}(\Omega) = \left\{ \phi \in L^2(\Omega) \mid \int_{\Omega} \phi = 0 \right\},$$

with its domain

$$D(A_0) = \left\{ \phi \in W^{2,2}(\Omega) \cap L^2_{\perp}(\Omega) \mid \frac{\partial \phi}{\partial \nu} = 0 \text{ on } \partial\Omega \right\}.$$

Then A_0 is self-adjoint and possesses bound self-adjoint fractional powers $A_0^{-\beta}$ for any $\beta > 0$ (cf. [29]).

Now the classical duality-based arguments lead to the boundedness of u in space-time L^2 .

Lemma 3.8. *Let (u, v, w) be a solution of (1.3). Then there exists a constant $C > 0$ such that*

$$\int_t^{t+\tau} \int_{\Omega} u^2 \leq C \quad \text{for all } t \in [0, T_{max} - \tau)$$

with $\tau = \min\{1, \frac{1}{2}T_{max}\}$.

Proof. According to Lemma 3.3, Lemma 3.6 and (1.5), we can find constants $C_1, C_2 > 0$ such that

$$C_1 \leq \gamma(v) \leq C_2.$$

Multiplying the third equation of (1.3) by α and adding the result equation to the first equation of (1.3), we get

$$(u + \alpha w)_t = \Delta(\gamma(v)u + \alpha w)$$

which can be rewritten as

$$(u + \alpha w - \bar{u} - \alpha \bar{w})_t = -A_0(\gamma(v)u + \alpha w - \overline{\gamma(v)u} - \alpha \bar{w}). \quad (3.15)$$

In view of (3.15) and the fact $\int_{\Omega}(u + \alpha w - \bar{u} - \alpha \bar{w}) = 0$, integrating by parts, we obtain

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{\Omega} |A_0^{-\frac{1}{2}}(u + \alpha w - \bar{u} - \alpha \bar{w})|^2 \\ &= \int_{\Omega} A_0^{-\frac{1}{2}}(u + \alpha w - \bar{u} - \alpha \bar{w}) \cdot A_0^{-\frac{1}{2}}(u + \alpha w - \bar{u} - \alpha \bar{w})_t \\ &= \int_{\Omega} A_0^{-1}(u + \alpha w - \bar{u} - \alpha \bar{w}) \cdot (u + \alpha w - \bar{u} - \alpha \bar{w})_t \\ &= - \int_{\Omega} A_0^{-1}(u + \alpha w - \bar{u} - \alpha \bar{w}) \cdot A_0(\gamma(v)u + \alpha w - \overline{\gamma(v)u} - \alpha \bar{w}) \\ &= - \int_{\Omega} (u + \alpha w - \bar{u} - \alpha \bar{w}) \cdot (\gamma(v)u + \alpha w - \overline{\gamma(v)u} - \alpha \bar{w}) \\ &= - \int_{\Omega} \gamma(v)(u - \bar{u})^2 - \bar{u} \int_{\Omega} \gamma(v)(u - \bar{u}) - \alpha \int_{\Omega} (1 + \gamma(v))(u - \bar{u})(w - \bar{w}) \\ &\quad - \alpha \bar{u} \int_{\Omega} \gamma(v)(w - \bar{w}) - \alpha^2 \int_{\Omega} (w - \bar{w})^2 \\ &\leq -C_1 \int_{\Omega} (u - \bar{u})^2 + C_2 |\Omega| \bar{u}^2 + 2\alpha(1 + C_2) |\Omega| \bar{u} \cdot \bar{w} + C_2 |\Omega| \alpha \bar{u} \cdot \bar{w} - \alpha^2 \int_{\Omega} (w - \bar{w})^2 \end{aligned}$$

which yields a constant $C_3 > 0$ such that

$$\frac{d}{dt} \int_{\Omega} |A_0^{-\frac{1}{2}}(u + \alpha w - \bar{u} - \alpha \bar{w})|^2 + 2C_1 \int_{\Omega} (u - \bar{u})^2 + 2\alpha^2 \int_{\Omega} (w - \bar{w})^2 \leq C_3. \quad (3.16)$$

By the Poincaré inequality and the fact

$$\int_{\Omega} A_0^{-\frac{1}{2}}(u + \alpha w - \bar{u} - \alpha \bar{w}) = 0,$$

we can find a constant $C_4 > 0$ such that

$$\int_{\Omega} \left| A_0^{-\frac{1}{2}}(u + \alpha w - \bar{u} - \alpha \bar{w}) \right|^2$$

$$\begin{aligned}
&\leq C_4 \int_{\Omega} \left| \nabla A_0^{-\frac{1}{2}}(u + \alpha w - \bar{u} - \alpha \bar{w}) \right|^2 \\
&= C_4 \int_{\Omega} |u + \alpha w - \bar{u} - \alpha \bar{w}|^2 \\
&\leq 2C_4 \int_{\Omega} (u - \bar{u})^2 + 2C_4 \alpha^2 \int_{\Omega} (w - \bar{w})^2 \\
&\leq 2C_4 \int_{\Omega} (u - \bar{u})^2 + 2C_4 \alpha^2 |\Omega| \|w_0\|_{L^\infty}^2
\end{aligned}$$

which combined with (3.16) implies there exists a constant $C_5 > 0$ such that

$$\begin{aligned}
\frac{d}{dt} \int_{\Omega} |A_0^{-\frac{1}{2}}(u + \alpha w - \bar{u} - \alpha \bar{w})|^2 + \frac{C_1}{2C_4} \int_{\Omega} \left| A_0^{-\frac{1}{2}}(u + \alpha w - \bar{u} - \alpha \bar{w}) \right|^2 \\
+ C_1 \int_{\Omega} (u - \bar{u})^2 + 2\alpha^2 \int_{\Omega} (w - \bar{w})^2 \leq C_5.
\end{aligned} \tag{3.17}$$

An application of Grönwall's inequality gives a constant $C_6 > 0$ such that

$$\int_{\Omega} |A_0^{-\frac{1}{2}}(u + \alpha w - \bar{u} - \alpha \bar{w})|^2 \leq C_6.$$

Integrating (3.17) over $(t, t + \tau)$, we get

$$\int_t^{t+\tau} \int_{\Omega} (u - \bar{u})^2 \leq C_7$$

for some constant $C_7 > 0$, which implies

$$\int_t^{t+\tau} \int_{\Omega} u^2 = \int_t^{t+\tau} \int_{\Omega} (u - \bar{u})^2 + \int_t^{t+\tau} \int_{\Omega} \bar{u}^2 \leq C_7 + \bar{u}^2 |\Omega|.$$

Hence, we complete the proof of the lemma. \square

3.4.2 L^2 -estimate of u

Lemma 3.9. *Let (u, v, w) be a solution of (1.3). Then there exists a constant $C > 0$ such that*

$$\int_{\Omega} |\nabla v|^2 \leq C \quad \text{for all } t \in (0, T_{max})$$

and

$$\int_t^{t+\tau} \int_{\Omega} |\Delta v|^2 \leq C \quad \text{for all } t \in [0, T_{max} - \tau).$$

Proof. By simple computations, we have

$$\begin{aligned}
\frac{1}{2} \frac{d}{dt} \int_{\Omega} |\nabla v|^2 &= \int_{\Omega} \nabla v \cdot \nabla v_t \\
&= \int_{\Omega} \nabla v \cdot \nabla (D\Delta v - v + u) \\
&= -D \int_{\Omega} |\Delta v|^2 - \int_{\Omega} |\nabla v|^2 - \int_{\Omega} u \Delta v
\end{aligned}$$

$$\leq -\frac{D}{2} \int_{\Omega} |\Delta v|^2 - \int_{\Omega} |\nabla v|^2 + \frac{1}{2D} \int_{\Omega} u^2$$

which leads to

$$\frac{d}{dt} \int_{\Omega} |\nabla v|^2 + D \int_{\Omega} |\Delta v|^2 + 2 \int_{\Omega} |\nabla v|^2 \leq \frac{1}{D} \int_{\Omega} u^2.$$

An application of the Grönwall inequality along with Lemma 2.2 and Lemma 3.8 gives a constant $C_1 > 0$ such that

$$\int_{\Omega} |\nabla v|^2 + \int_t^{t+\tau} \int_{\Omega} |\Delta v|^2 \leq C_1.$$

Therefore, we finish the proof of this lemma. \square

Lemma 3.10. *Let (u, v, w) be a solution of (1.3). There exist constants $c, C > 0$ such that for any $p \geq 2$, we have*

$$\begin{aligned} & \frac{d}{dt} \int_{\Omega} u^p + cp(p-1) \int_{\Omega} u^{p-2} |\nabla u|^2 \\ & \leq Cp(p-1) \int_{\Omega} u^p |\nabla v|^2 + Cp(p-1) \int_{\Omega} u^p \text{ for all } t \in (0, T_{max}). \end{aligned}$$

Proof. According to Lemma 2.1, Lemma 3.2 and the hypothesis on F , we can find a constant $C_1 > 0$ such that

$$F(w) \leq C_1.$$

Noticing Lemma 3.3, Lemma 3.6 and the smoothing property of γ , there exist constants $C_2, C_3 > 0$ such that

$$\gamma(v) \geq C_2 \tag{3.18}$$

and

$$\frac{|\gamma'(v)|^2}{\gamma(v)} \leq C_3. \tag{3.19}$$

Using u^{p-1} with $p \geq 2$ as a test function for the first equation in (1.3), integrating the resulting equation by parts and using Young's inequality, we obtain

$$\begin{aligned} \frac{1}{p} \frac{d}{dt} \int_{\Omega} u^p &= \int_{\Omega} u^{p-1} \Delta(\gamma(v)u) + \alpha \int_{\Omega} u^p F(w) \\ &\leq -(p-1) \int_{\Omega} \gamma(v) u^{p-2} |\nabla u|^2 + (p-1) \int_{\Omega} \gamma'(v) u^{p-1} \nabla u \cdot \nabla v + C_1 \alpha \int_{\Omega} u^p \\ &\leq -\frac{p-1}{2} \int_{\Omega} \gamma(v) u^{p-2} |\nabla u|^2 + \frac{p-1}{2} \int_{\Omega} \frac{|\gamma'(v)|^2}{\gamma(v)} u^p |\nabla v|^2 + C_1 \alpha \int_{\Omega} u^p \end{aligned}$$

which, combined with (3.18) and (3.19), yields that

$$\begin{aligned} & \frac{d}{dt} \int_{\Omega} u^p + \frac{p(p-1)}{2} C_2 \int_{\Omega} u^{p-2} |\nabla u|^2 \\ & \leq \frac{p(p-1)}{2} C_3 \int_{\Omega} u^p |\nabla v|^2 + C_1 \alpha p \int_{\Omega} u^p. \end{aligned}$$

This finishes the proof of this lemma. \square

Now the uniform-time boundedness of u in $L^2(\Omega)$ can be established.

Lemma 3.11. *Let (u, v, w) be a solution of (1.3). Then there is a constant $C > 0$ such that*

$$\int_{\Omega} u^2 \leq C \quad \text{for all } t \in (0, T_{max}).$$

Proof. Taking $p = 2$ in Lemma 3.10, we get the following estimate

$$\frac{d}{dt} \int_{\Omega} u^2 + C_1 \int_{\Omega} |\nabla u|^2 \leq C_2 \int_{\Omega} u^2 |\nabla v|^2 + C_2 \int_{\Omega} u^2 \quad (3.20)$$

for some constants $C_1, C_2 > 0$. Using Lemma 3.9, the Gagliardo-Nirenberg inequality (see [11, Lemma 2.5]) and Young's inequality, we can find constants $C_3, C_4, C_5 > 0$ such that

$$\begin{aligned} C_2 \int_{\Omega} u^2 |\nabla v|^2 &\leq C_2 \|u\|_{L^4}^2 \|\nabla v\|_{L^4}^2 \\ &\leq C_3 \left(\|\nabla u\|_{L^2}^{\frac{n}{4}} \|u\|_{L^2}^{\frac{4-n}{4}} + \|u\|_{L^2} \right)^2 \left(\|\Delta v\|_{L^2}^{\frac{n}{4}} \|\nabla v\|_{L^2}^{\frac{4-n}{4}} + \|\nabla v\|_{L^2} \right)^2 \\ &\leq 4C_3 \left(\|\nabla u\|_{L^2}^{\frac{n}{2}} \|u\|_{L^2}^{\frac{4-n}{2}} \|\Delta v\|_{L^2}^{\frac{n}{2}} \|\nabla v\|_{L^2}^{\frac{4-n}{2}} + \|\nabla u\|_{L^2}^{\frac{n}{2}} \|u\|_{L^2}^{\frac{4-n}{2}} \|\nabla v\|_{L^2}^2 \right. \\ &\quad \left. + \|u\|_{L^2}^2 \|\Delta v\|_{L^2}^{\frac{n}{2}} \|\nabla v\|_{L^2}^{\frac{4-n}{2}} + \|u\|_{L^2}^2 \|\nabla v\|_{L^2}^2 \right) \\ &\leq C_4 \left(\|\nabla u\|_{L^2}^{\frac{n}{2}} \|u\|_{L^2}^{\frac{4-n}{2}} \|\Delta v\|_{L^2}^{\frac{n}{2}} + \|\nabla u\|_{L^2}^{\frac{n}{2}} \|u\|_{L^2}^{\frac{4-n}{2}} \right. \\ &\quad \left. + \|u\|_{L^2}^2 \|\Delta v\|_{L^2}^{\frac{n}{2}} + \|u\|_{L^2}^2 \right) \\ &\leq \frac{C_1}{2} \|\nabla u\|_{L^2}^2 + C_5 \left(1 + \|\Delta v\|_{L^2}^{\frac{2n}{4-n}} + \|\Delta v\|_{L^2}^{\frac{n}{2}} \right) \|u\|_{L^2}^2 \\ &\leq \frac{C_1}{2} \|\nabla u\|_{L^2}^2 + C_5 (1 + \|\Delta v\|_{L^2}^2) \|u\|_{L^2}^2 \end{aligned}$$

where in the last inequality we have used the fact $n \leq 2$ so that $\frac{2n}{4-n} \leq 2$. This along with (3.20) implies there exists a constant $C_6 > 0$ such that

$$\frac{d}{dt} \int_{\Omega} u^2 + \frac{C_1}{2} \int_{\Omega} |\nabla u|^2 \leq C_6 (1 + \|\Delta v\|_{L^2}^2) \int_{\Omega} u^2.$$

An application of Lemma 3.9 and Lemma 2.3 gives the desired result. \square

3.4.3 L^∞ -estimate of u

Lemma 3.12. *Let (u, v, w) be a solution of (1.3). For any $p \geq 1$, there exists $C > 0$ such that*

$$\int_{\Omega} |\nabla v|^p \leq C \quad \text{for all } t \in (0, T_{max}).$$

Moreover, if $n = 1$, then we can find $C > 0$ such that

$$\|\nabla v\|_{L^\infty} \leq C \quad \text{for all } t \in (0, T_{max}).$$

Proof. Applying Lemma 2.4 and Lemma 3.11, the desired result is obtained. \square

Combining Lemma 3.10 and Lemma 3.12, we get the following result.

Lemma 3.13. *Let (u, v, w) be a solution of (1.3). There exists a constant $C > 0$ such that for any $p \geq 2$, we have*

$$\frac{d}{dt} \int_{\Omega} u^p + p(p-1) \int_{\Omega} u^p \leq Cp(p-1)(1+p)^{\frac{6n}{4-n}} \left(\int_{\Omega} u^{\frac{p}{2}} \right)^2 \quad \text{for all } t \in (0, T_{max}).$$

Proof. From Lemma 3.10, we can find constants $C_1, C_2 > 0$ such that

$$\frac{d}{dt} \int_{\Omega} u^p + C_1 \frac{p-1}{p} \int_{\Omega} |\nabla u^{\frac{p}{2}}|^2 \leq C_2 p(p-1) \int_{\Omega} u^p |\nabla v|^2 + C_2 p(p-1) \int_{\Omega} u^p. \quad (3.21)$$

By means of Lemma 3.12 and Young's inequality, there exists a constant $C_3 > 0$ such that

$$C_2 p(p-1) \int_{\Omega} u^p |\nabla v|^2 \leq C_2 p(p-1) \left(\int_{\Omega} u^{2p} \right)^{\frac{1}{2}} \left(\int_{\Omega} |\nabla v|^4 \right)^{\frac{1}{2}} \leq C_3 p(p-1) \left(\int_{\Omega} u^{2p} \right)^{\frac{1}{2}}$$

and

$$(C_2 + 1)p(p-1) \int_{\Omega} u^p \leq C_3 p(p-1) \left(\int_{\Omega} u^{2p} \right)^{\frac{1}{2}},$$

which combined with (3.21) gives that

$$\frac{d}{dt} \int_{\Omega} u^p + C_1 \frac{p-1}{p} \int_{\Omega} |\nabla u^{\frac{p}{2}}|^2 + p(p-1) \int_{\Omega} u^p \leq 2C_3 p(p-1) \left(\int_{\Omega} u^{2p} \right)^{\frac{1}{2}}. \quad (3.22)$$

The Gagliardo-Nirenberg inequality and Young's inequality with ε provide constants $C_4, C_5 > 0$ such that

$$\begin{aligned} 2C_3 \left(\int_{\Omega} u^{2p} \right)^{\frac{1}{2}} &= 2C_3 \|u^{\frac{p}{2}}\|_{L^4}^2 \\ &\leq C_4 \left(\|\nabla u^{\frac{p}{2}}\|_{L^2}^{\frac{3n}{2(n+2)}} \|u^{\frac{p}{2}}\|_{L^1}^{\frac{4-n}{2(n+2)}} + \|u^{\frac{p}{2}}\|_{L^1} \right)^2 \\ &\leq 2C_4 \left(\|\nabla u^{\frac{p}{2}}\|_{L^2}^{\frac{3n}{n+2}} \|u^{\frac{p}{2}}\|_{L^1}^{\frac{4-n}{n+2}} + \|u^{\frac{p}{2}}\|_{L^1}^2 \right) \\ &\leq C_1 \frac{1}{2p^2} \|\nabla u^{\frac{p}{2}}\|_{L^2}^2 + C_5 \left(1 + p^{\frac{6n}{4-n}} \right) \|u^{\frac{p}{2}}\|_{L^1}^2. \end{aligned} \quad (3.23)$$

Substituting (3.23) into (3.22) and noting $1 + p^{\frac{6n}{4-n}} \leq (1+p)^{\frac{6n}{4-n}}$, we obtain

$$\frac{d}{dt} \int_{\Omega} u^p + p(p-1) \int_{\Omega} u^p \leq C_5 p(p-1)(1+p)^{\frac{6n}{4-n}} \left(\int_{\Omega} u^{\frac{p}{2}} \right)^2.$$

Then we get the desired result. \square

Now we can obtain the uniform-in-time boundedness of u in $L^\infty(\Omega)$.

Lemma 3.14. *Let (u, v, w) be a solution of (1.3). There exists a constant $C > 0$ such that*

$$\|u\|_{L^\infty} \leq C \quad \text{for all } t \in (0, T_{max}).$$

Proof. According to Lemma 3.13, there exists a constant $C_1 > 0$ such that for any $p \geq 2$

$$\frac{d}{dt} \int_{\Omega} u^p + p(p-1) \int_{\Omega} u^p \leq C_1 p(p-1) (1+p)^{\frac{6n}{4-n}} \left(\int_{\Omega} u^{\frac{p}{2}} \right)^2$$

which gives

$$\frac{d}{dt} \left[e^{p(p-1)t} \int_{\Omega} u^p \right] \leq C_1 e^{p(p-1)t} p(p-1)(1+p)^6 \left(\int_{\Omega} u^{\frac{p}{2}} \right)^2. \quad (3.24)$$

Integrating (3.24) over the time interval $[0, t]$ for $0 < t < T_{max}$, we get

$$\int_{\Omega} u^p \leq \int_{\Omega} u_0^p + C_1(1+p)^6 \sup_{0 \leq t \leq T_{max}} \left(\int_{\Omega} u^{\frac{p}{2}} \right)^2.$$

Then, employing a standard Moser iteration (cf. [2]) or the similar argument as in [32], the desired result can be obtained. \square

4 Proof of Theorem 1.1

Proof of Theorem 1.1. Theorem 1.1 is a consequence of Lemma 3.14, Lemma 2.4 and the extensibility criterion Lemma 2.1. \square

Acknowledgement. The authors thank the referee for nice comments improving the exposition of the manuscript. The research of Z.A. Wang was supported by the Hong Kong RGC GRF grant No. 15303019 (Project ID P0030816) and an internal grant No. UAH0 (Project ID P0031504) from the Hong Kong Polytechnic University.

References

- [1] J. Ahn and C.W. Yoon. Global well-posedness and stability of constant equilibria in parabolic-elliptic chemotaxis systems without gradient sensing. *Nonlinearity*, 32(4):1327–1351, 2019.
- [2] N.D. Alikakos. L^p bounds of solutions of reaction-diffusion equations. *Comm. Partial Differential Equations*, 4(8):827–868, 1979.
- [3] H. Amann. Nonhomogeneous linear and quasilinear elliptic and parabolic boundary value problems. In *Function spaces, differential operators and nonlinear analysis (Friedrichroda, 1992)*, volume 133 of *Teubner-Texte Math.*, pages 9–126. Teubner, Stuttgart, 1993.
- [4] H. Brézis and W.A. Strauss. Semi-linear second-order elliptic equations in L^1 . *J. Math. Soc. Japan*, 25:565–590, 1973.
- [5] M. Burger, P. Lanrençot and A. Trescases, Delayed blow-up for chemotaxis models with local sensing, arXiv:2005.02734v2, 2020.
- [6] L. Desvillettes, Y.J. Kim, A. Trescases, and C.W. Yoon. A logarithmic chemotaxis model featuring global existence and aggregation. *Nonlinear Anal. Real World Appl.*, 50:562–582, 2019.
- [7] X.F. Fu, L.H. Tang, C.L. Liu, J.D. Huang, T. Hwa, and P. Lenz. Stripe formation in bacterial systems with density-suppressed motility. *Physical Review Letters*, 108(19):1981–1988, 2012.
- [8] K. Fujie. Boundedness in a fully parabolic chemotaxis system with singular sensitivity. *J. Math. Anal. Appl.*, 424(1):675–684, 2015.

- [9] K. Fujie and J. Jiang. Comparison methods for a KellerSegel-type model of pattern formations with density-suppressed motilities. *arXiv:2001.00144v1*.
- [10] K. Fujie and J. Jiang. Global existence for a kinetic model of pattern formation with density-suppressed motilities. *J. Differential Equations*, 269(6):5338–5378, 2020.
- [11] H.Y. Jin, Y.J. Kim, and Z.A. Wang. Boundedness, stabilization, and pattern formation driven by density-suppressed motility. *SIAM J. Appl. Math.*, 78(3):1632–1657, 2018.
- [12] H.Y. Jin, S.J. Shi, and Z.A. Wang. Boundedness and asymptotics of a reaction-diffusion system with density-dependent motility. *J. Differential Equations*, 269(9):6758–6793, 2020.
- [13] H.Y. Jin and Z.A. Wang. Critical mass on the Keller-Segel system with signal-dependent motility. *Proc. Amer. Math. Soc.*, 148(11):4855–4873, 2020.
- [14] H.Y. Jin and Z.A. Wang. Global dynamics and spatio-temporal patterns of predator-prey systems with density-dependent motion. *European J. Appl. Math.*, Doi:10.1017/S0956792520000248, 2020.
- [15] H.Y. Jin and Z.A. Wang. The Keller-Segel system with logistic growth and signal-dependent motility. *Discrete Contin. Dyn. Syst. Ser. B*, 26(6):3023-3041, 2021.
- [16] P. Kareiva and G.T. Odell, Swarms of predators exhibit “preytaxis” if individual predators use area-restricted search. *Am. Nat.*, 130:233-270, 1987.
- [17] E.F. Keller and L.A. Segel, Models for chemotaxis. *J. Theor. Biol.*, 30:225-234, 1971.
- [18] R. Kowalczyk and Z. Szymańska. On the global existence of solutions to an aggregation model. *J. Math. Anal. Appl.*, 343(1):379–398, 2008.
- [19] J. Li and Z.A. Wang. Traveling wave solutions to the density-suppressed motility model. *Preprint*, 2020.
- [20] Z.R. Liu and J. Xu. Large time behavior of solutions for density-suppressed motility system in higher dimensions. *J. Math. Anal. Appl.*, 475(2):1596–1613, 2019.
- [21] C. Liu et al. Sequential establishment of stripe patterns in an expanding cell population. *Science*, 334:238-241, 2011.
- [22] Y. Lou and M. Winkler. Global existence and uniform boundedness of smooth solutions to a cross-diffusion system with equal diffusion rates. *Comm. Partial Differential Equations*, 40(10):1905–1941, 2015.
- [23] R. Lui and H. Ninomiya, Traveling wave solutions for a bacteria system with density-suppressed motility, *Disc. Cont. Dyn. Syst.-B*, 24: 931-940, 2018.
- [24] W.B. Lv. Global existence for a class of chemotaxis-consumption systems with signal-dependent motility and generalized logistic source. *Nonlinear Anal. Real World Appl.*, 56:103160, 2020.
- [25] W.B. Lv and Q. Wang. A chemotaxis system with signal-dependent motility, indirect signal production and generalized logistic source: Global existence and asymptotic stabilization. *J. Math. Anal. Appl.*, 488(2):124108, 2020.
- [26] W.B. Lv and Q.Y. Wang. Global existence for a class of chemotaxis systems with signal-dependent motility, indirect signal production and generalized logistic source. *Z. Angew. Math. Phys.*, 71(2):Paper No. 53, 2020.
- [27] W.B. Lv and Q.Y. Wang. An n -dimensional chemotaxis system with signal-dependent motility and generalized logistic source: global existence and asymptotic stabilization. *Proc. Roy. Soc. Edinburgh Sect. A*, Doi: 10.1017/prm.2020.38, 2020.

- [28] M.J. Ma, R. Peng, and Z.A. Wang. Stationary and non-stationary patterns of the density-suppressed motility model. *Phys. D*, 402:132259, 13, 2020.
- [29] M. Schechter. Self-adjoint realizations in another Hilbert space. *Amer. J. Math.*, 106(1):43–65, 1984.
- [30] J. Smith-Roberge, D. Iron and T. Kolokolnikov, Pattern formation in bacterial colonies with density-dependent diffusion. *Eur. J. Appl. Math.*, 30:196-218, 2019.
- [31] C. Stinner, C. Surulescu, and M. Winkler. Global weak solutions in a PDE-ODE system modeling multiscale cancer cell invasion. *SIAM J. Math. Anal.*, 46(3):1969–2007, 2014.
- [32] Y.S. Tao and Z.A. Wang. Competing effects of attraction vs. repulsion in chemotaxis. *Math. Models Methods Appl. Sci.*, 23(1):1–36, 2013.
- [33] Y.S. Tao and M. Winkler. Effects of signal-dependent motilities in a Keller-Segel-type reaction-diffusion system. *Math. Models Methods Appl. Sci.*, 27(9):1645–1683, 2017.
- [34] R. Temam. *Infinite-dimensional dynamical systems in mechanics and physics*, volume 68 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1988.
- [35] J.P. Wang and M.X. Wang. Boundedness in the higher-dimensional Keller-Segel model with signal-dependent motility and logistic growth. *J. Math. Phys.*, 60(1):011507, 14, 2019.
- [36] Z.A. Wang. On the parabolic-elliptic Keller-Segel system with signal-dependent motilities: a paradigm for global boundedness and steady states. *Preprint*, 2020.
- [37] Z.A. Wang and X. Xu Steady states and pattern formation of the density-suppressed motility model. *IMA. J. Appl. Math.*, doi:10.1093/imamat/hxab006, 2021.
- [38] C. Yoon and Y.J. Kim. Global existence and aggregation in a Keller-Segel model with Fokker-Planck diffusion. *Acta Appl. Math.*, 149:101–123, 2017.