



# Uniqueness and convergence on equilibria of the Keller–Segel system with subcritical mass

Jun Wang<sup>a</sup>, Zhian Wang<sup>b</sup>, and Wen Yang<sup>c</sup>

<sup>a</sup>Faculty of Science, Jiangsu University, Zhenjiang, Jiangsu, People's Republic of China; <sup>b</sup>Department of Applied Mathematics, Hong Kong Polytechnic University, Hung Hom, Kowloon, Hong Kong;

<sup>c</sup>Wuhan Institute of Physics and Mathematics Chinese Academy of Sciences, Wuhan, People's Republic of China.

## ABSTRACT

This paper is concerned with the uniqueness of solutions to the following nonlocal semi-linear elliptic equation

$$\Delta u - \beta u + \lambda \frac{e^u}{\int_{\Omega} e^u} = 0 \text{ in } \Omega, \quad (*)$$

where  $\Omega$  is a bounded domain in  $\mathbb{R}^2$  and  $\beta, \lambda$  are positive parameters. The above equation arises as the stationary problem of the well-known classical Keller–Segel model describing chemotaxis. For Eq. (\*) with Neumann boundary condition, we establish an integral inequality and prove that the solution of Eq. (\*) is unique if  $0 < \lambda \leq 8\pi$  and  $u$  satisfies some symmetric properties. While for Eq. (\*) with Dirichlet boundary condition, the same uniqueness result is obtained without symmetric condition by a different approach inspired by some recent works (Gui and Moradifam, 2018, *Invent. Math.* 214(3):1169–1204; Gui and Moradifam, Proc. Am. Math. Soc. 146(3):1231–1124). As an application of the uniqueness results, we prove that the radially symmetric solution of the classical Keller–Segel system with subcritical mass subject to Neumann boundary conditions will converge to the unique constant equilibrium as time tends to infinity if  $\Omega$  is a disc in two dimensions. As far as we know, this is the first result that asserts the exact asymptotic behavior of solutions to the classical Keller–Segel system with subcritical mass in two dimensions.

## ARTICLE HISTORY

Received 9 April 2018

Accepted 20 January 2019

## KEYWORDS

Keller–Segel system; mean field equation; subcritical-mass; uniqueness

## 1. Introduction

This paper is concerned with a system of partial differential equations modeling chemotaxis which refers to the active motion of biological species toward higher concentrations of chemical substances that they emit themselves. The classical chemotaxis model is well-known as the Keller–Segel (KS) system [1, 2], reading as

$$\begin{cases} v_t = \nabla \cdot (\nabla v - v \nabla u) & \text{in } \Omega, \\ u_t = \Delta u - \beta u + v & \text{in } \Omega, \\ \partial_\nu v = \partial_\nu u = 0 & \text{on } \partial\Omega, \\ v(x, 0) = v_0(x), \quad u(x, 0) = u_0(x) & \text{in } \Omega, \end{cases} \tag{1.1}$$

where  $\Omega$  is smooth bounded domain in  $\mathbb{R}^N (N \geq 2)$ ,  $v(x, t)$  and  $u(x, t)$  denote the cell density and chemical concentration, respectively;  $\beta$  is a positive constant accounting for the chemical death rate,  $\nu$  is the unit outward normal vector at the boundary  $\partial\Omega$ .

The underlying equations in system (1.1) were proposed by Keller and Segel in 1970 [1, 2] to describe the aggregation phase of cellular slime molds *Dictyostelium discoideum* in response to the chemical substance cyclic adenosine monophosphate (cAMP) that they secreted. An immediate result derived from Eq. (1.1) is the mass conservation for  $v(x, t)$  by integrating the first equation

$$\lambda =: \int_{\Omega} v(x, t) dx = \int_{\Omega} v_0(x) dx.$$

Among other things, the most striking feature of the KS system (1.1) lies in the existence of critical space dimension and critical mass. Roughly speaking, in one-dimensional space ( $N=1$ ), the KS system (1.1) admits globally bounded classical solutions [3, 4]. In the two-dimensional radially symmetric domain (disc), global bounded classical solutions exist (cf. [5]) if  $\lambda < 8\pi$  (subcritical mass), whereas solutions may blow up in finite or infinite time (cf. [6, 7]) if  $\lambda > 8\pi$  (super-critical mass), where the critical mass  $8\pi$  becomes  $4\pi$  if  $\Omega$  is a general domain without symmetry. In three dimensions, the solution may blow up in finite time for any mass  $\lambda > 0$  (cf. [8]). Therefore  $N=2$  is a borderline space dimension where a critical mass exists, and as such the KS model (1.1) and its variants (extensions) have attracted extensive attentions and a vast number of fruitful results have been obtained (cf. review papers [9–11] and book [12]). However, as far as we know, the long-time behavior of solutions to the KS system (1.1) with subcritical mass in two dimensions still remains unknown. It is the purpose of this paper to explore this open question. Precisely we shall show that the radial solution of (1.1) in a disc with subcritical mass will converge to the unique constant equilibrium as time tends to infinity. Since it has been shown in [13] that the globally bounded solution (if it exists) of (1.1) converges to the steady states in  $L^\infty$ -norm, our question boils down to prove the uniqueness of constant equilibrium for the stationary problem of (1.1) in  $\Omega \subset \mathbb{R}^2$

$$\begin{cases} \nabla \cdot (\nabla v - v \nabla u) = 0 & \text{in } \Omega, \\ \Delta u - \beta u + v = 0 & \text{in } \Omega, \\ \partial_\nu v = \partial_\nu u = 0 & \text{on } \partial\Omega, \end{cases} \tag{1.2}$$

in the case of subcritical mass  $\int_{\Omega} v(x) dx < 8\pi$ . To study the stationary system (1.2), we note that the first equation can be written as

$$\nabla \cdot (v \nabla (\log v - u)) = 0.$$

Testing the above equation against  $\log v - u$ , then an integration by parts shows that any solution of (1.2) verifies the equation

$$\int_{\Omega} v |\nabla(\log v - u)|^2 dx = 0,$$

so that  $v = Ce^u$  for some positive constant  $C$ . Hence, all the solutions  $(v, u)$  of (1.2) would satisfy the relation  $v = Ce^u$ . Denoting

$$\lambda = \int_{\Omega} v(x) dx,$$

we have  $C = \frac{\lambda}{\int_{\Omega} e^u dx}$ . Therefore, the stationary problem (1.2) is equivalent to the following nonlocal semi-linear Neumann problem

$$\begin{cases} \Delta u - \beta u + \lambda \frac{e^u}{\int_{\Omega} e^u dx} = 0 & \text{in } \Omega, \\ \partial_{\nu} u = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.3)$$

and  $v = \frac{e^u}{\int_{\Omega} e^u dx}$ . Integrating the first equation of (1.3), one immediately obtains that

$$\int_{\Omega} u(x) dx = \frac{\lambda}{\beta}.$$

Then by a (new) shifted variable

$$U(x) = u(x) - \bar{u}, \quad \bar{u} = \frac{1}{|\Omega|} \int_{\Omega} u(x) dx = \frac{\lambda}{\beta|\Omega|}, \quad (1.4)$$

the problem (1.3) can be transformed as the following

$$\begin{cases} \Delta U - \beta U + \lambda \left( \frac{e^U}{\int_{\Omega} e^U dx} - \frac{1}{|\Omega|} \right) = 0 & \text{in } \Omega, \\ \partial_{\nu} U = 0 & \text{on } \partial\Omega. \end{cases} \quad (1.5)$$

From (1.4) we know that the mean of  $U$  is zero, namely

$$\int_{\Omega} U(x) dx = 0. \quad (1.6)$$

When  $\beta = 0$ , the nonlinear differential equation in (1.5) is closely related to the Gaussian curvature problem on a surface (see [14]). In Onsager's vortex theory, the asymptotic limit of the Gibbs measure yields to similar problems (see [15–17]). Moreover the first equation of (1.5) on a torus arises in the Chern–Simons gauge theory [18] and has been investigated among others by Struwe and Tarantello [19]. Chen and Lin [20] and Machioldi [21] have independently derived the Leray–Schauder Topological degree for (1.5) on the Riemann surface without boundary. By assuming  $\beta > \frac{2}{|\Omega|} - \lambda_1$  and  $\lambda > 4\pi$ , where  $\lambda_1$  is referred to the first eigenvalue of the Neumann eigenvalue problem, Wang and Wei [22] and Horstmann [23] have independently shown the existence of non-constant solutions for (1.5). Very recently, Battaglia obtains the existence of non-constant solutions to (1.5) with  $(\lambda, \beta)$  in a wider range, see [24, Theorem 1.1] for the details.

Since the Neumann problem (1.5)–(1.6) admits a trivial solution  $U = 0$ , it is natural to ask whether there is any other solution. When  $\beta = 0$ , a simple application of the

maximum principle will show that  $U \equiv 0$  is the unique solution to problem (1.5)–(1.6) provided that  $\lambda \leq 0$  though it is not of interest in applications. For  $\lambda > 0$ , the first equation of (1.5) (without Neumann boundary condition) on a standard sphere admits only the constant solution whenever  $\lambda < 8\pi$  (cf. [16, 25, 26]), while the uniqueness result also holds for some flat torus provided  $\lambda \leq 8\pi$  (cf. [27]). When  $\Omega$  is the unit disc, Horstmann and Lucia [28] proved that  $U \equiv 0$  is the unique solution for Neumann problem (1.5)–(1.6) if  $\lambda \leq \frac{32}{\pi}$ . The uniqueness result also holds when the solution is constant on the boundary (i.e.  $\text{osc}_{\partial\Omega}(u) \equiv 0$ ) and  $\lambda \leq 8\pi$ . As a consequence of their results, we may expect to conclude that  $U \equiv 0$  is the unique radially symmetric solution to the Neumann problem (1.5)–(1.6) if  $\lambda \leq 8\pi$  and  $\Omega$  is the unit disc.

We remark that the above results for the Neumann problem (1.5) with  $\beta = 0$  cannot be converted to the Neumann problem (1.2) since the transformation (1.4) no longer works for  $\beta = 0$ . Indeed, if  $\beta = 0$ , the integration of the second equation of (1.2) along with the Neumann boundary condition yields that  $\int_{\Omega} v dx = 0$ , which along with the fact  $v \geq 0$  indicates that (1.2) with  $\beta = 0$  has no solution if  $\lambda > 0$  or  $u \equiv 0$  if  $\lambda = 0$ . Hence the solution for the case  $\beta = 0$  is clear but not of interest. To the best of our knowledge, there are very few results available for the case  $\beta > 0$  for which the shifted problem (1.5) is equivalent to (1.3) under the shifting (1.4). In this paper, we shall investigate the uniqueness of solutions to (1.3) with  $\beta > 0$  which is equivalent to (1.2) with  $v = \frac{e^u}{\int_{\Omega} e^u dx}$ . Hereafter, we shall assume  $\beta > 0$  unless otherwise stated. When  $\Omega$  is the unit disc, Senba and Suzuki [29, Theorem 4] have obtained the existence of nontrivial radial solutions of (1.3) for  $\lambda > 8\pi$ . In this paper, we shall prove that the constant equilibrium  $u = \frac{\lambda}{\beta\pi}$  is the only radially symmetric solution to the Neumann problem (1.3) if  $\Omega$  is the unit disc and  $\lambda \leq 8\pi$ . Using the maximum principle, one can get  $u > 0$  in  $\Omega$  and we leave the proof in the Appendix (see Lemma 6.1). Therefore our study will be focused on the positive solutions of (1.3).

Our first result concerning the Neumann problem (1.3) is the following:

**Theorem 1.1.** *Let  $\Omega$  be a non-empty bounded open set in  $\mathbb{R}^2$ . Then for  $\lambda \leq 8\pi$  the constant  $u = \frac{\lambda}{\beta|\Omega|}$  is the unique solution to the problem (1.3) with  $\text{osc}_{\partial\Omega}(u) \equiv 0$ .*

**Remark 1.1.** When  $\Omega$  is the unit disc, we can obtain from Theorem 1.1 that  $u = \frac{\lambda}{\beta\pi}$  is the only radially symmetric solution to the problem (1.3) provided  $\lambda \leq 8\pi$ .

Inspired by the result of Theorem 1.1, one may ask what happens if the condition  $\text{osc}_{\partial\Omega} u \equiv 0$  is replaced by some symmetric condition, i.e., the solutions which are invariant under the group of isometries of a unit disc. To state the results, letting  $G$  be an isometry group of the unit disc  $B$ , we introduce the following classes of functions:

$$H^G := \{w \in H^1(B) : w = w \circ g, \forall g \in G\}.$$

and

$$H^\circ G := \left\{ w \in H^G : \int_{\Omega} w = 0 \right\},$$

where  $B$  denotes the unit disc. For any  $\theta \in (0, 2\pi)$  and  $C \neq 0$  we use the notations

$$R_\theta := \text{rotation of angle } \theta,$$

$$D_C := \text{reflection with respect to } \vec{OC} \text{ (} O \text{ stands for the origin point),}$$

and  $\langle g \rangle$  stands for the subgroup generated by an isometry  $g$ . We list the following examples which are used usually:

- (a) When  $G = SO(2) = \left\{ \begin{pmatrix} \cos \theta & \sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}, \theta \in [0, 2\pi) \right\}$ , the space  $H^G$  consists of the radial  $H^1(B)$ -functions.
- (b)  $H^{\langle R_\pi \rangle}$  consists of the  $H^1(B)$ -functions satisfying  $w(x) = w(-x)$ .
- (c)  $H^{\langle R_{2\pi/m} \rangle}$  consists of the  $H^1(B)$ -functions satisfying  $w(x) = w(R_{2\pi/m} \circ x)$ , where

$$R_{2\pi/m} \circ x = \begin{pmatrix} x_1 & x_2 \end{pmatrix} \cdot \begin{pmatrix} \cos(2\pi/m) & \sin(2\pi/m) \\ \sin(2\pi/m) & \cos(2\pi/m) \end{pmatrix}$$

for any  $x = (x_1, x_2) \in B$ .

- (d) Let  $D_m : \langle R_{2\pi/m}, D_C \rangle$  be a group generated by the rotation  $R_{2\pi/m}$  and a reflection  $D_C$ . The group with  $2m$  elements is called the  $m$ th *dihedral* group. For  $m \geq 3$ , it is the symmetry group of regular  $m$ -polygon.

For solutions of (1.3) which are invariant under rotations, we have:

**Theorem 1.2.** *Let  $G = \langle R_{2\pi/m} \rangle$ ,  $m \in \mathbb{N}$  with  $m \geq 2$ . If a non-constant function  $u \in H^G$  solves the problem (1.3) in  $B$ , then*

$$\lambda > \Lambda_m = \begin{cases} \frac{64}{\pi}, & \text{if } m = 2, \\ 8\pi, & \text{if } m \geq 3. \end{cases} \tag{1.7}$$

To prove Theorems 1.1 and 1.2, we first consider a more general problem

$$\begin{cases} \Delta u - g(u) + f(u) = 0 & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ \partial_\nu u = 0 & \text{on } \partial\Omega, \end{cases} \tag{1.8}$$

where  $f, g$  satisfy the following conditions:

$$f, g \in C^2(\mathbb{R}), \quad f(t), f'(t) > 0, \quad g(t), g'(t) \geq 0 \text{ for } t \geq 0. \tag{1.9}$$

For Eq. (1.8) with  $f, g$  satisfying (1.9), we derive an integral inequality

$$\frac{A_g}{2} \int_{-\infty}^{\infty} f'(t) \mu(t) (|\Omega| - \mu(t)) dt \geq \int_{-\infty}^{\infty} f(t) I_\Omega^2(\mu(t)) dt, \tag{1.10}$$

where

$$A_g = \frac{1}{|\Omega|} \int_\Omega g(u) dx, \quad \mu(t) = |\{x \in \Omega : u(x) > t\}|$$

and  $I_\Omega(s)$  is referred to the “isoperimetric profile” of  $\Omega$  with volume  $s$ , the detailed definition of the “isoperimetric profile” will be given in Section 2. In [28, 30], the authors study the problem (1.8) with  $g(u)$  replaced by a constant  $A = \frac{1}{|\Omega|} \int_\Omega f(u) dx$ . In their approach, a similar inequality of (1.10) is obtained with  $A_g$  replaced by  $A$ . Different from their problem, we assume that  $g(u)$  is a non-constant function with some increasing property. To derive the inequality (1.10), we have to estimate the integration of  $g(u)$  with respect to the level set of  $u$ . With a simple manipulation, see Lemma 2.2, we manage to estimate the integration of  $g(u)$  in terms of the mean value  $A_g$ . This is the key

point which helps us to generalize all the results to  $g(u)$  – a nontrivial function. Another new ingredient in our proof is that we find to study the augmented functions  $\Psi$  and  $\tilde{\Psi}$  (see the definition of  $\Psi$  and  $\tilde{\Psi}$  in (3.2)) together to show that the jump of each discontinuous point is positive, see (3.3). It is different from the simpler case of constant function  $g(u)$ , where  $\Psi$  and  $\tilde{\Psi}$  can be studied separately (cf. [28, 30]).

We shall apply our uniqueness results to exploit the asymptotic behavior of solutions to the KS system (1.1). Specifically we show that if  $\lambda < \Lambda_m$  any rotationally invariant solution to (1.1) is uniformly bounded, exists globally and converges to the unique constant equilibrium.

**Theorem 1.3.** *Consider the problem (1.1) in the unit disc  $B \subset \mathbb{R}^2$ . Let  $(v, u)$  be a  $C^2$ -solution to (1.1) belonging to  $H^G \times \dot{H}^G$  with  $G = \langle \mathbb{R}_{2\pi/m} \rangle$  ( $m \geq 2$ ). If  $\lambda < \Lambda_m$  with  $\Lambda_m$  defined in (1.7), then the solution of (1.1) is globally defined and*

$$\lim_{t \rightarrow \infty} (v, u)(\cdot, t) = \left( \frac{\lambda}{\pi}, \frac{\lambda}{\beta\pi} \right) \text{ in } C^1(\bar{B}). \quad (1.11)$$

*In particular, if  $\lambda < 8\pi$ , any radial solution  $(v, u)$  of (1.1) will satisfy (1.11).*

We remark that the convergence of solutions (1.1) to the constant equilibrium as time tends to infinity in the critical case  $\lambda = 8\pi$  remains unknown. Indeed the asymptotic behavior of solutions to the classical KS system in the case of critical mass is a rather complicated issue and the answer was only partially given in the whole space  $\mathbb{R}^2$  for the case  $\beta = 0$  (cf. [31, 32]). Although the asymptotics of solutions for critical mass in a bounded domain still remains unknown in this paper, our results in Theorem 1.1 and Theorem 1.2 will shed lights on this problem for further pursues in the future.

The last problem to be considered in this paper is the uniqueness of solutions for the following Dirichlet problem:

$$\begin{cases} \Delta u - \beta u + \lambda \frac{e^u}{\int_{\Omega} e^u dx} = 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases} \quad (1.12)$$

The above problem is the steady state problem of the KS system with mixed zero-flux and Dirichlet boundary conditions

$$\begin{cases} v_t = \nabla \cdot (\nabla v - v \nabla u) & \text{in } \Omega, \\ u_t = \Delta u - \beta u + v & \text{in } \Omega, \\ \partial_\nu v - v \partial_\nu u = u = 0 & \text{on } \partial\Omega, \\ u(x, 0) = u_0(x), \quad v(x, 0) = v_0(x) & \text{in } \Omega. \end{cases}$$

When  $\beta = 0$ , Suzuki [33] proved that if  $\Omega$  is simply-connected, then the problem (1.12) has a unique solution for  $0 < \lambda < 8\pi$ . The uniqueness result for  $\lambda = 8\pi$  is obtained by Chang et al. [34]. Later Bartolucci and Lin [35] extended the result to multiple-connected domains. Recently, based on the Bol's inequalities and equi-measurable symmetric rearrangement, Gui and Moradifam [36] developed a new tool named "Sphere Covering Inequality". This inequality and its generalizations are applied to establish the best constant in a Moser–Trudinger type inequalities, some symmetry and uniqueness results for the mean field equations and Onsager vortex (cf. [36–40] for

details). Based on their results in [36, 39], we shall derive the uniqueness of (1.12) in the subcritical mass cases.

**Theorem 1.4.** *Let  $\Omega$  be an open bounded and simply-connected set in  $\mathbb{R}^2$ . If  $0 \leq \lambda < 8\pi$ , then there exists a unique solution of the problem (1.12) with  $\beta > 0$ . While if  $\lambda = 8\pi$ , Eq. (1.12) has at most one solution.*

**Remark 1.2.** We remark that the the Dirichlet problem (1.12) no longer has a constant solution provided that  $\lambda > 0$ . In other words, the unique solution of the problem (1.12) for  $0 \leq \lambda < 8\pi$  must be non-trivial, which differs from the Neumann problem (1.3). Furthermore, we can show that the degree of Eq. (1.12) is 0 for  $\lambda \in (8\pi, 16\pi)$ , see Theorem 5.2. As a consequence, the Dirichlet problem (1.12) has no solution or at least two solutions if  $\lambda \in (8\pi, 16\pi)$ . On the other hand, we note that the existence of solutions to (1.12) with  $\lambda = 8\pi$  is still unknown and it may depend on the topology of  $\Omega$  (cf. [35] for  $\beta = 0$ ). In this sense,  $\lambda = 8\pi$  is a threshold for the uniqueness of solutions for the Dirichlet problem (1.12).

The paper is organized as follows. In Section 2, we derive a differential inequality which involves the distribution of  $u$ , the function  $\int_{\{u>t\}} f(u) dx$ , the average of function  $g(u)$  and the isoperimetric profile of the domain. Based on this result, in Section 3 we derive an integral inequality which is the key to the proof of Theorems 1.1 and 1.2. Section 4 is devoted to the proof of our results on the Neumann problem (1.3). While in Section 5 we study the uniqueness of solutions to the Dirichlet problem. In the Appendix, we give a proof for the positivity of solutions to (1.3) in  $\Omega$ .

## 2. A differential inequality

In the present section, we shall establish a differential inequality, which plays an important role in our discussion. Given two functions  $f, g$  which satisfy (1.9). We set

$$\mathcal{C}(f, g) = \{t | f(t) = g(t), t \geq 0\}, \quad (2.1)$$

and consider the following nonlinear problem:

$$\begin{cases} \Delta u - g(u) + f(u) = 0 & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ \partial_\nu u = 0 & \text{on } \partial\Omega. \end{cases} \quad (2.2)$$


To proceed with our argument, we make the following preparation. Denote by  $\mathcal{H}^s$  the  $s$ -dimensional Hausdorff measure in  $\Omega$ . Given  $\omega \subset \Omega$ , its perimeter relative to  $\Omega$  is defined as

$$\mathcal{P}(\omega, \Omega) := \mathcal{H}^1(\partial\omega \cap \Omega),$$

and its area  $\mathcal{H}^2(\omega)$  will be denoted by  $|\omega|$ . (See Figure 1 for an illustration of  $\partial\omega \cap \Omega$ .)

**Definition 2.1.** Let  $\mathcal{O}_\Omega$  be the class of open subsets  $\omega \subset \Omega$  satisfying

$$\omega = \omega' \cap \Omega, \quad \omega' \in \mathbb{R}^2 \text{ of class } C^1.$$



**Figure 1.** For the figure on the left, the part on the left of the dashed curve is  $\omega$  and the dashed curve represents  $\partial\omega \cap \Omega$ , for the figure on the right, the part enclosed by the dashed closed curve is  $\omega$  and the dashed closed curve represents  $\partial\omega \cap \Omega$ .

The “isoperimetric profile” of  $\Omega$  is the function  $I_\Omega : [0, |\Omega|] \rightarrow (0, \infty)$  defined as

$$I_\Omega(s) := \inf \{ \mathcal{H}^1(\partial\omega \cap \Omega) : \omega \in \mathcal{O}_\Omega, \mathcal{H}^2(\omega) = s \}, \quad \forall s \in (0, |\Omega|],$$

and we set  $I_\Omega(0) = 0$ .

**Remark 2.1.** For each  $\omega \in \mathcal{O}_\Omega$  the boundary of  $\omega$  in  $\Omega$  is a 1-submanifold of class  $C^1$ . We remark that only  $\partial\omega \cap \Omega$  is taken into consideration in the definition of the isoperimetric profile. We mention two properties of the isoperimetric profile that will be used in the following:

$$I_\Omega(s) = I_\Omega(|\Omega| - s), \quad \forall s \in [0, |\Omega|]; \tag{2.3}$$

$$I_\Omega(s) = 0 \iff s = 0 \text{ or } s = |\Omega|. \tag{2.4}$$

The symmetric property (2.3) readily follows from the definition of isoperimetric profile, while for (2.4) we refer to [41].

We will need the following lemma.

**Lemma 2.1.** *Assume that (1.9) holds. Then for any non-constant solution  $u$  of problem (2.1), it holds that*

$$\mathcal{H}^2(\{x \in \Omega | u(x) = t\}) = 0$$

whenever  $t \in \mathbb{R} \setminus \mathcal{C}(f, g)$ .

*Proof.* Given a fixed  $t \in \mathbb{R} \setminus \mathcal{C}(f, g)$ , we denote

$$E(t) := \{x \in \Omega | u(x) = t\}$$

and divide  $E(t) = E_1(t) \cup E_2(t)$  with

$$E_1(t) := \{x \in \Omega | u(x) = t, \nabla u \neq 0\},$$

and

$$E_2(t) := \{x \in \Omega | u(x) = t, \nabla u(x) = 0\}.$$

Using the implicit function theorem, we get the set  $E_1(t)$  is locally a one-dimensional manifold. Then we can deduce that  $E_1(t)$  is at most countable union of sets of measure zero. Hence  $\mathcal{H}^2(E_1(t)) = 0$  for all  $t \in \mathbb{R}$ .



For the set  $E_2(t)$ , using the Eq. (2.2) we have

$$\Delta u \neq 0, \text{ whenever } u(x) \notin \mathcal{C}(f, g).$$

Therefore if  $t \notin \mathcal{C}(f, g)$ , then we have

$$\begin{aligned} E_2(t) &\subseteq \{x \in \Omega : \nabla u(x) = 0, u(x) \notin \mathcal{C}(f, g)\} \\ &\subseteq \bigcup_{i=1}^2 \{x \in \Omega : \partial_i u(x) = 0, \nabla(\partial_i u)(x) \neq 0\}. \end{aligned}$$

Thus, for  $t \notin \mathcal{C}(f, g)$ , the set  $E_2(t)$  is contained in a finite union of 1-submanifolds. So we conclude that  $\mathcal{H}^2(E_2(t)) = 0$  for all  $t \in \mathbb{R} \setminus \mathcal{C}(f, g)$ .

From the above discussion we get  $\mathcal{H}^2(E(t)) = 0$  for all  $t \in \mathbb{R} \setminus \mathcal{C}(f, g)$ , which finishes the proof. □

For any solution  $u$  of (2.1), we introduce the following notations:

$$F(t) := \int_{\{u > t\}} f(u) dx, \quad G(t) := \int_{\{u > t\}} g(u) dx, \quad \mu(t) := |\{u > t\}| = \mathcal{H}^2(\{u > t\}),$$

and

$$\tilde{F}(t) := \int_{\{u < t\}} f(u) dx, \quad \tilde{G}(t) := \int_{\{u < t\}} g(u) dx, \quad \tilde{\mu}(t) := |\{u < t\}| = \mathcal{H}^2(\{u < t\}).$$

By Lemma 2.1, the functions  $F, \tilde{F}, G, \tilde{G}, \mu$ , and  $\tilde{\mu}$  are continuous on  $\mathbb{R} \setminus \mathcal{C}(f, g)$ . Furthermore, if the set  $\mathcal{C}(f, g)$  is finite, then the above functions are monotone and therefore differentiable a.e.  $t \in \mathbb{R}$ .

We set

$$A_g = \frac{1}{|\Omega|} \int_{\Omega} g(u) dx.$$

Before stating the main result of this section, we give the following lemma.

**Lemma 2.2.** Suppose that  $h$  is a non-decreasing function. For any solution  $u$  of (2.2), it holds that

$$A_h |\{u > t\}| \leq \int_{\{u > t\}} h(u) dx \text{ and } A_h |\{u < t\}| \geq \int_{\{u < t\}} h(u) dx, \quad \forall t \in \left[ \min_{\Omega} u, \max_{\Omega} u \right].$$

*Proof.* We only give the proof of the first inequality, the other one can be proved similarly. Using  $h$  as non-decreasing, we immediately get

$$\frac{\int_{\{u > t\}} h(u) dx}{|\{u > t\}|} \geq \frac{\int_{\{u \leq t\}} h(u) dx}{|\{u \leq t\}|}.$$

As a consequence

$$\frac{\int_{\{u > t\}} h(u) dx}{|\{u > t\}|} \geq \frac{\int_{\{u > t\}} h(u) dx + \int_{\{u \leq t\}} h(u) dx}{|\{u > t\}| + |\{u \leq t\}|} = A_h,$$

which implies

$$A_h|\{u > t\}| \leq \int_{\{u > t\}} h(u)dx.$$

It proves the result. □

Now we establish the main result in this section.

**Proposition 2.3.** *Let  $f, g$  be a function satisfying (1.9). Assume  $\mathcal{C}(f, g)$  is finite. Then any non-constant solution  $u$  of (2.2) satisfies the following inequalities:*

$$\frac{d}{dt} \left( A_g f \mu^2 - F^2 \right) \geq A_g f'(t) \mu^2(t) + 2f(t) I_\Omega^2(\mu(t)), \quad \forall t \in \mathbb{R} \setminus \mathcal{D}, \tag{2.5}$$

$$\frac{d}{dt} \left( A_g f \tilde{\mu}^2 - \tilde{F}^2 \right) \geq A_g f'(t) \tilde{\mu}^2(t) + 2f(t) I_\Omega^2(\tilde{\mu}(t)), \quad \forall t \in \mathbb{R} \setminus \mathcal{D}, \tag{2.6}$$

where  $\mathcal{D} := \{u(x) : x \in \Omega, \nabla u(x) = 0\}$  and  $I_\Omega$  stands for the isoperimetric profile of  $\Omega$ .

*Proof.* At first, we notice that Sard’s Theorem ensures that the set of critical value  $\mathcal{D}$  associate to  $u$  has Lebesgue measure zero in  $\mathbb{R}$ .

Let us prove (2.5) first. By Lemma 2.1, the functions  $F, G$  and  $\mu$  are continuous on  $\mathbb{R} \setminus \mathcal{C}(f, g)$ . Therefore, by using the co-area formula, we obtain

$$\mu'(t) = - \int_{\{u=t\}} \frac{1}{|\nabla u|} d\mathcal{H}^1, \quad \forall t \in \mathbb{R} \setminus \mathcal{D}, \tag{2.7}$$

$$F'(t) = - \int_{\{u=t\}} \frac{f(u)}{|\nabla u|} d\mathcal{H}^1 = f(t)\mu'(t), \quad \forall t \in \mathbb{R} \setminus \mathcal{D}, \tag{2.8}$$

$$G'(t) = - \int_{\{u=t\}} \frac{g(u)}{|\nabla u|} d\mathcal{H}^1 = g(t)\mu'(t), \quad \forall t \in \mathbb{R} \setminus \mathcal{D}. \tag{2.9}$$

Second, by integrating Eq. (2.2) on the set  $\{u > t\}$  and using the Stoke’s Theorem, we obtain

$$\int_{\partial\{u > t\}} |\nabla u| d\mathcal{H}^1 = F(t) - G(t), \quad \forall t \in \mathbb{R} \setminus \mathcal{D}. \tag{2.10}$$

Since

$$\begin{aligned} \partial\{u > t\} &= (\partial\{u > t\} \cap \Omega) \cup (\partial\{u > t\} \cap \partial\Omega) \\ &= (\{u = t\} \cap \Omega) \cup (\partial\{u > t\} \cap \partial\Omega), \end{aligned}$$

and furthermore  $u$  satisfies the Neumann boundary condition on  $\partial\{u > t\} \cap \partial\Omega$ , the left hand side of (2.10) equals to  $\int_{\{u=t\} \cap \Omega} |\nabla u| d\mathcal{H}^1$ . Based on this observation, we have

$$F(t) = G(t) + \int_{\{u=t\} \cap \Omega} |\nabla u| d\mathcal{H}^1, \quad \forall t \in \mathbb{R} \setminus \mathcal{D}. \tag{2.11}$$

Using (2.11) and the assumption  $f \geq 0$  we derive:

$$\begin{aligned}
 -F'(t)F(t) &= \left( G(t) + \int_{\{u=t\} \cap \Omega} |\nabla u| d\mathcal{H}^1 \right) \int_{\{u=t\}} \frac{f(t)}{|\nabla u|} d\mathcal{H}^1 \\
 &= f(t) \int_{\{u=t\} \cap \Omega} |\nabla u| d\mathcal{H}^1 \int_{\{u=t\}} \frac{1}{|\nabla u|} d\mathcal{H}^1 + \left( \int_{\{u=t\}} \frac{f(t)}{|\nabla u|} d\mathcal{H}^1 \right) \int_{\{u>t\}} g(u) dx \\
 &\geq f(t) \left( \int_{\{u=t\} \cap \Omega} d\mathcal{H}^1 \right)^2 - f(t) A_g \mu(t) \mu'(t) \\
 &= f(t) [\mathcal{H}^1(\{u = t\} \cap \Omega)]^2 - f(t) A_g \mu(t) \mu'(t),
 \end{aligned} \tag{2.12}$$

where we have used the Schwarz inequality and Lemma 2.2. Recall the definition of isoperimetric profile of  $\Omega$ , we have

$$\int_{\{u=t\} \cap \Omega} d\mathcal{H}^1 = \mathcal{P}(\{u > t\}, \Omega) \geq I_\Omega(\mu(t)). \tag{2.13}$$

Hence, (2.12) and (2.13) yield

$$\frac{1}{2} (A_g f \mu^2 - F^2)'(t) \geq f(t) I_\Omega^2(\mu(t)) + \frac{A_g}{2} f'(t) \mu^2(t),$$

where we have used

$$f \mu \mu' = f \frac{(\mu^2)'}{2} = \frac{1}{2} ((f \mu^2)' - f' \mu^2).$$

Following almost the same argument, we can derive (2.6) by replacing (2.7)–(2.9) and (2.11) with

$$\tilde{\mu}'(t) = \int_{\{u=t\}} \frac{1}{|\nabla u|} d\mathcal{H}^1, \quad \tilde{F}'(t) = f(t) \tilde{\mu}'(t), \quad \tilde{G}'(t) = g(t) \tilde{\mu}'(t),$$

and

$$\tilde{F}(t) = \tilde{G}(t) - \int_{\{u=t\}} |\nabla u| d\mathcal{H}^1.$$

□

### 3. An integral inequality

In this section, we shall derive an important integral inequality, which plays a key role in the proof of our uniqueness result.

**Proposition 3.1.** *Assume (1.9) holds and the set  $\mathcal{C}(f, g)$  is finite. The any solution  $u$  of (2.2) satisfies*

$$\frac{A_g}{2} \int_{-\infty}^{\infty} f'(t) \mu(t) (|\Omega| - \mu(t)) dt \geq \int_{-\infty}^{\infty} f(t) I_\Omega^2(\mu(t)) dt. \tag{3.1}$$

*Proof.* For any solution  $u$  of (2.2), we set

$$t_0 := \min_{\Omega} u, \quad t_1 := \max_{\Omega} u, \quad F_{\Omega} = \int_{\Omega} f(u) dx.$$

The result of Proposition 3.1 will follow by integrating the differential inequalities (2.5)–(2.6) on the interval  $(t_0, t_1)$  as shown below.

Let us consider the functions

$$\Psi := A_g f \mu^2 - F^2 \quad \text{and} \quad \tilde{\Psi} := A_g f \tilde{\mu}^2 - \tilde{F}^2. \quad (3.2)$$

By Lemma 2.1, the functions  $\Psi$  and  $\tilde{\Psi}$  are continuous on any interval  $[a, b] \subset \mathbb{R} \setminus \mathcal{C}(f, g)$ . At the point  $t \in \mathcal{C}(f, g)$ , these functions may be discontinuous, and hence we have to treat it separately.

For each  $a \in \mathcal{C}(f, g)$ , we set  $\Gamma_a := \{u = a\}$ , and claim

$$\begin{aligned} & \Psi(a^+) + \tilde{\Psi}(a^+) - \Psi(a^-) - \tilde{\Psi}(a^-) \\ &= 2f(a)|\Gamma_a| \left[ (F(a) - \mu(a)A_g) + (\tilde{\mu}(a)A_g - \tilde{F}(a)) \right] \\ &\geq 0, \end{aligned} \quad (3.3)$$

where  $\Psi(a^{\pm}) := \lim_{\varepsilon \rightarrow 0} \Psi(a \pm \varepsilon)$ ,  $\tilde{\Psi}(a^{\pm}) := \lim_{\varepsilon \rightarrow 0} \tilde{\Psi}(a \pm \varepsilon)$ .

Indeed, according to the definition of  $\Psi$  and  $\tilde{\Psi}$ , we have

$$\begin{aligned} \Psi(a^+) - \Psi(a^-) &= A_g f(a) \left[ \mu(a)^2 - (\mu(a) + |\Gamma_a|)^2 \right] \\ &\quad + (F(a) + f(a)|\Gamma_a|)^2 - F(a)^2, \end{aligned} \quad (3.4)$$

and

$$\begin{aligned} \tilde{\Psi}(a^+) - \tilde{\Psi}(a^-) &= A_g f(a) \left[ (\tilde{\mu}(a) + |\Gamma_a|)^2 - \tilde{\mu}(a)^2 \right] \\ &\quad + \tilde{F}(a)^2 - (\tilde{F}(a) + f(a)|\Gamma_a|)^2. \end{aligned} \quad (3.5)$$

Adding (3.4) and (3.5) together, we have

$$\begin{aligned} & \Psi(a^+) + \tilde{\Psi}(a^+) - \Psi(a^-) - \tilde{\Psi}(a^-) \\ &= 2f(a)|\Gamma_a| \left[ (F(a) - \mu(a)A_g) + (\tilde{\mu}(a)A_g - \tilde{F}(a)) \right]. \end{aligned} \quad (3.6)$$

It is easy to see that  $A_g = A_f$  from the Eq. (2.2) and the Neumann boundary condition. Using Lemma 2.2, we get

$$\frac{F(a)}{\mu(a)} \geq \frac{\int_{\Omega} f(u) dx}{|\Omega|} = A_f = A_g \quad \text{and} \quad \frac{\tilde{F}(a)}{\tilde{\mu}(a)} \leq \frac{\int_{\Omega} f(u) dx}{|\Omega|} = A_f = A_g,$$

which imply  $F(a) - \mu(a)A_g \geq 0$  and  $\tilde{\mu}(a)A_g - \tilde{F}(a) \geq 0$ . As a consequence, the right hand side of (3.6) is non-negative. Thus the claim (3.3) is proved.

By (2.5) and (2.6), we have both  $\Psi$  and  $\tilde{\Psi}$  are monotone increasing on the intervals  $[t_0, t_1] \setminus \cup_{a \in \mathcal{C}(f, g)} (a - \varepsilon, a + \varepsilon)$  for  $\varepsilon > 0$ . Then we get

$$\int_{t_0}^{t_1} \Psi' dt \leq \sum_{a \in \mathcal{C}(f, g)} (\Psi(a^-) - \Psi(a^+)) + F_{\Omega}^2 - A_g f(t_0) |\Omega|^2. \quad (3.7)$$

Similarly, we derive that

$$\int_{t_0}^{t_1} \tilde{\Psi}' dt \leq \sum_{a \in \mathcal{C}(f,g)} (\tilde{\Psi}(a^-) - \tilde{\Psi}(a^+)) + A_g f(t_1) |\Omega|^2 - F_{\Omega}^2. \tag{3.8}$$

By Proposition 2.3, (3.7)–(3.8) and (3.3), we get

$$\begin{aligned} A_g |\Omega|^2 (f(t_1) - f(t_0)) &\geq \int_{t_0}^{t_1} \left( A_g f'(t) \mu^2(t) + 2f(t) I_{\Omega}^2(\mu(t)) \right) dt \\ &\quad + \int_{t_0}^{t_1} \left( A_g f'(t) \tilde{\mu}^2(t) + 2f(t) I_{\Omega}^2(\tilde{\mu}(t)) \right) dt \\ &= \int_{t_0}^{t_1} A_g f'(t) \left( \mu(t)^2 + (|\Omega| - \mu(t))^2 \right) dt \\ &\quad + 4 \int_{t_0}^{t_1} f(t) I_{\Omega}^2(\mu(t)) dt \\ &= A_g |\Omega|^2 \int_{t_0}^{t_1} f'(t) dt + 4 \int_{t_0}^{t_1} f(t) I_{\Omega}^2(\mu(t)) dt \\ &\quad + 2A_g \int_{t_0}^{t_1} f'(t) \mu(t) (\mu(t) - |\Omega|) dt \end{aligned} \tag{3.9}$$

where we used  $\tilde{\mu} = |\Omega| - \mu$  and  $I_{\Omega}(\mu) = I_{\Omega}(|\Omega| - \mu)$ . Since  $f$  is differentiable, inequality (3.9) yields:

$$0 \geq A_g \int_{t_0}^{t_1} f'(t) \mu(t) (\mu(t) - |\Omega|) dt + 2 \int_{t_0}^{t_1} f(t) I_{\Omega}^2(\mu(t)) dt,$$

or equivalently

$$A_g \int_{t_0}^{t_1} f'(t) \mu(t) (|\Omega| - \mu(t)) dt \geq 2 \int_{t_0}^{t_1} f(t) I_{\Omega}^2(\mu(t)) dt,$$

which proves (3.1). □

**Remark 3.1.** Instead of using the inequality  $\mathcal{H}^1(\{u = t\} \cap \Omega) \geq I_{\Omega}(\mu(t))$  from (2.13), we can keep the term  $\mathcal{H}^1(\{u = t\})$  and repeat the arguments of deriving (3.1) to get

$$\frac{A_g}{2} \int_{-\infty}^{\infty} f'(t) \mu(t) (|\Omega| - \mu(t)) dt \geq \int_{-\infty}^{\infty} f(t) [\mathcal{H}^1(\{u = t\} \cap \Omega)]^2 dt. \tag{3.10}$$

In Section 4, we will get the uniqueness result asserted in Theorem 1.1 from the above inequality (3.10).

**Remark 3.2.** When  $\Omega$  is replaced by the manifold  $M$ , we can also derive a counterpart result of Proposition 3.1. Then a similar result of [30, Theorem 1.1] and some related conclusions can be obtained by the same arguments (see [30, section 3] for more details).

### 4. The uniqueness of the Neumann problem (1.3)

In this section, we shall apply the inequalities established in the preceding section to prove the main results of the Neumann problem (1.3).

With

$$f(u) = \lambda \frac{e^u}{\int_{\Omega} e^u dx} \quad \text{and} \quad g(u) = \beta u, \tag{4.1}$$

the problem (2.2) turns to be

$$\begin{cases} \Delta u - \beta u + \lambda \frac{e^u}{\int_{\Omega} e^u dx} = 0 & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ \partial_{\nu} u = 0 & \text{on } \Omega. \end{cases} \tag{4.2}$$

It is easy to check that  $f, g$  verify the condition (1.9). In the next result we shall prove

$$|\mathcal{C}(f, g)| = \left| \mathcal{C} \left( \lambda \frac{e^u}{\int_{\Omega} e^u dx}, \beta u \right) \right| < +\infty.$$

**Lemma 4.1.** *Let  $f, g$  be defined in (4.1). Then*

$$1 \leq |\mathcal{C}(f, g)| \leq 2.$$

*Proof.* We divide our proof into two steps.

Step 1.  $|\mathcal{C}(f, g)| \geq 1$ . Using the boundary condition, we have

$$\int_{\Omega} f(u) dx = \beta \int_{\Omega} u dx. \tag{4.3}$$

Then we claim that  $f(u(x)) = \beta u(x)$  must happen for some  $x \in \Omega$ . Otherwise, we have either  $f(u(x)) > \beta u(x)$  or  $f(u(x)) < \beta u(x)$  for all  $x \in \Omega$ , which leads to  $\int_{\Omega} f(u) dx > \beta \int_{\Omega} u dx$  or  $\int_{\Omega} f(u) dx < \beta \int_{\Omega} u dx$ , it contradicts to (4.3). Thus  $\{u | f(u) = \beta u\} \neq \emptyset$ .

Step 2.  $|\mathcal{C}(f, g)| \leq 2$ . Considering the function  $h(x) = f(x) - \beta x$ , which is convex by noticing that  $f$  is a convex function. As a consequence, the function  $h$  possesses at most two roots. Hence, we finish the proof. □

Next we prove the following theorem, which is equivalent to Theorem 1.1.

**Theorem 4.2.** *Let  $u$  be a non-constant solution of problem (4.2) with  $\text{osc}_{\partial\Omega}(u) = 0$ . Then the following inequality holds:*

$$\beta \int_{\Omega} u dx > 8\pi.$$

*Proof.* To prove Theorem 4.2, we need the following inequality (see Remark 3.1)

$$\frac{A_g}{2} \int_{-\infty}^{\infty} f'(t) \mu(t) (|\Omega| - \mu(t)) dt \geq \int_{-\infty}^{\infty} f(t) [\mathcal{H}^1(\{u = t\} \cap \Omega)]^2 dt. \tag{4.4}$$

By the setting of  $g$ , we have  $A_g = \beta \bar{u}$ . Setting  $u_0 := u|_{\partial\Omega}$ , we consider the regular values of  $u$ :

$$\text{Reg}(u) := \{t \in \mathbb{R} : \nabla u(x) \neq 0, \forall x \in u^{-1}(t)\}.$$

From Sard’s theorem we have  $\mathbb{R} \setminus \text{Reg}(u)$  has zero Lebesgue measure, which along with the implicit function theorem implies that

$$\{u = t\} \Subset \Omega, \{u = t\} \text{ is a } 1\text{-submanifold of class } C^1 \text{ for } t \in \text{Reg}(u) \setminus \{u_0\}.$$

Next we claim that  $\{u < t\}$  or  $\{u > t\}$  must be contained in a domain enclosed by some connected branch of the set  $\{u = t\}$ . Indeed, for any  $t \neq u_0$ , without loss of generality we may assume  $t < u_0$ . Then it is evident to see that  $\{u < t\} \cap \partial\Omega = \emptyset$  and the claim holds. As a consequence

$$\mathcal{H}^1(\{u = t\} \cap \Omega) = \mathcal{H}^1(\{u = t\}).$$

Next, by using the isoperimetric inequality we have

$$[\mathcal{H}^1(\{u = t\})]^2 \geq 4\pi \min\{\mu(t), |\Omega - \mu(t)|\}, \quad \forall t \in \text{Reg}(u) \setminus \{u_0\}. \tag{4.5}$$

From (4.5) we get

$$\frac{f(t)[\mathcal{H}^1(\{u = t\})]^2}{f'(t)\mu(t)(|\Omega| - \mu(t))} \geq \frac{4\pi}{\max\{\mu(t), |\Omega| - \mu(t)\}} > \frac{4\pi}{|\Omega|}, \quad t \in (t_0, t_1). \tag{4.6}$$

Combining with the fact  $\mathbb{R} \setminus \text{Reg}(u)$  has zero Lebesgue measure, (4.4) and (4.6), we have

$$\beta\bar{u} > \frac{8\pi}{|\Omega|},$$

which yields the result. □

Before proving Theorem 1.2, we shall give the following result on the general solutions of (1.3) on the unit disc.

**Proposition 4.3.** *Let  $\Omega$  be the unit disc. If  $u$  is a non-constant solution solving the problem (1.3), then*

$$\lambda > \frac{32}{\pi}.$$

*Proof of Proposition 4.3.* When  $\Omega$  is a disc, it is proved in [42, 18.1.3] that

$$\frac{I_{\Omega}^2(s)}{s(|\Omega| - s)} > \frac{16}{\pi|\Omega|}, \quad \forall s \in \left(0, \frac{|\Omega|}{2}\right).$$

Using the condition  $f \geq f' > 0$ , we have

$$\frac{I_{\Omega}^2(s)f(s)}{s(|\Omega| - s)f'(s)} > \frac{16}{\pi|\Omega|},$$

which along with Proposition 3.1 implies

$$\beta\bar{u} > \frac{32}{\pi|\Omega|}.$$

On the other hand, we have  $\lambda = \beta \int_{\Omega} u$ . Thus, we get  $\lambda > \frac{32}{\pi}$  and it proves the conclusion. □

In the following, we shall consider the case that  $u$  is invariant under a rotation  $R_\theta$ . To study the class of functions which are invariant by a rotation  $R_{\frac{2\pi}{m}}$ , we recall a definition in [28, 30]:

**Definition 4.1.** (isoperimetric profile) Given a group  $G$  of isometries of  $\Omega$ , we consider the class of open subsets:

$$\mathcal{O}_\Omega^G := \{\omega \subset \mathcal{O}_\Omega : g(\omega) = \omega, \forall g \in G\}.$$

The “ $G$ -isoperimetric profile” of  $\Omega$  is defined as

$$I_\Omega^G(s) := \inf\{\mathcal{H}^1(\partial\omega \cap \Omega) : \omega \in \mathcal{O}_\Omega^G, \mathcal{H}^2(\omega) = s\}, \quad \forall s \in (0, |\Omega|].$$

We set  $I_\Omega^G(0) = 0$ .

In the setting of “ $G$ -isoperimetric profile”, we can generalize the Proposition 3.1 to the following result:

**Proposition 4.4.** Let  $\Omega \subseteq \mathbb{R}^2$  be a piecewise  $C^1$ -domain. If  $G$  is a group of isometry of  $\Omega$  and  $u \in C^2(\bar{\Omega}) \cap H^G$  is a non-constant solution of (4.2), then

$$\frac{\beta \bar{u}}{2} \int_{-\infty}^{\infty} f'(t)\mu(t)(|\Omega| - \mu(t))dt \geq \int_{-\infty}^{\infty} f(t)[I_\Omega^G(\mu(t))]^2 dt.$$

*Proof.* We can follow the proof of Proposition 3.1 step by step to prove Proposition 4.4, just by noticing that for almost every  $t \in \mathbb{R}$  the level sets  $\{u > t\}$  and  $\{u < t\}$  belong to  $\mathcal{O}_\Omega^G$ .  $\square$

To continue our discussion, we need the following two lemmas. For the proof we refer the readers to [28, Lemma 4.6–4.7].

**Lemma 4.5.** Let  $B$  be a disc of  $\mathbb{R}^2$  and set  $G = \langle R_{2\pi/m} \rangle$ ,  $m \geq 2$ .

(a) If  $\omega \in \mathcal{O}_B^G$  is such that

$$(i) \mathcal{H}^1(\bar{\omega} \cap \partial B) = 0 \text{ or } (ii) \mathcal{H}^1(\overline{B \setminus \omega} \cap \partial B) = 0,$$

then  $\mathcal{H}^1(\partial\omega \cap B) \geq \min\{(4\pi\omega)^{\frac{1}{2}}, (4\pi[|B| - |\omega|])^{\frac{1}{2}}\}$ .

(b) If  $\omega = \cup_{i=0}^{m-1} R_{\frac{2\pi}{m}}^i(\omega_0)$  (disjoint union) with  $\omega_0$  satisfying

$$\omega_0 \in \mathcal{O}_B, \mathcal{H}^1(\bar{\omega}_0 \cap \partial B) > 0 \text{ and } \mathcal{H}^1(\overline{B \setminus \omega_0} \cap \partial B) > 0, \tag{4.7}$$

then

$$\mathcal{H}^1(\partial\omega \cap B) \geq nI_B\left(\frac{|\omega|}{m}\right). \tag{4.8}$$

If  $B \setminus \omega = \cup_{i=0}^{m-1} R_{\frac{2\pi}{m}}^i(\omega_0)$  with  $\omega_0$  satisfying (4.7), then (4.8) still holds

**Lemma 4.6.** For a disc  $B \subset \mathbb{R}^2$  and  $G = \langle R_{2\pi/m} \rangle$  it holds

$$\frac{[I_B^G(s)]^2}{s(|B| - s)} > \min\left\{\frac{4\pi}{|B|}, \frac{16m}{\pi|B|}\right\}.$$

With the above two lemmas, we can prove Theorem 1.2.



*Proof of Theorem 1.2.* For the solution  $u \in H^G$  with  $G = \langle R_{2\pi/m} \rangle$ , each of the level set  $\Omega_t := \{u > t\}$  is invariant under the action of the group. Then we deduce from Proposition 4.4 that

$$\frac{\beta \int_{\Omega} u dx}{2|\Omega|} \int_{-\infty}^{\infty} f'(t)\mu(t)(|\Omega| - \mu(t)) dt \geq \int_{-\infty}^{\infty} f(t) [I_{\Omega}^G(\mu(t))]^2 dt,$$

with  $f(u) = \lambda \frac{e^u}{\int_{\Omega} e^u dx}$  and  $\Omega = B$ . Using Lemma 4.6, we get

$$\frac{\lambda}{2|\Omega|} > \begin{cases} \frac{32}{\pi|\Omega|}, & \text{if } m = 2, \\ \frac{4\pi}{|\Omega|}, & \text{if } m \geq 3. \end{cases}$$

Then

$$\lambda > \begin{cases} \frac{64}{\pi}, & \text{if } m = 2, \\ 8\pi, & \text{if } m \geq 3, \end{cases}$$

which finishes the whole proof. □

Next we shall apply Theorem 1.2 to derive the optimal inequalities for the functional  $J_{\lambda}(u)$  :

$$J_{\lambda}(u) = \frac{1}{2} \int_{\Omega} |\nabla u|^2 dx + \frac{\beta}{2} \int_{\Omega} |u|^2 dx - \lambda \log \left( \int_{\Omega} e^u dx \right).$$

**Proposition 4.7.** *Let  $J_{\lambda}^m(\cdot)$  be the restriction of  $J_{\lambda}(\cdot)$  to the space  $H^G$  with  $G = \langle R_{2\pi/m} \rangle$ ,  $m \geq 2$ . Then the following holds:*

- a. *The functional  $J_{\lambda}^m(\cdot)$  is bounded from below whenever  $\lambda \leq 8\pi$ .*
- b. *If  $m \geq 3$  and  $\lambda \leq 8\pi$ , the functional  $J_{\lambda}^m(\cdot)$  admits a unique global minimizer given by  $u \equiv \frac{\lambda}{\beta|\Omega|}$ .*
- c. *For  $m = 2$ , the functional  $J_{\lambda}^m(\cdot)$  admits a global minimizer for each  $\lambda \leq 8\pi$ . Furthermore, the global minimizer is unique and given by  $\frac{\lambda}{\beta|\Omega|}$  whenever  $\lambda \leq \frac{64}{|\Omega|}$ .*

*Proof.* Let us first prove that when  $\lambda \leq 8\pi$  we may find a constant  $C > 0$  depending only on  $\lambda$  and  $|\Omega|$  such that

$$J_{\lambda}(u) \geq -C, \quad \forall u \in H^{\langle R_{2\pi/m} \rangle}.$$

We need the following inequality. For a bounded domain  $\Omega$  of  $\mathbb{R}^2$  whose boundary is  $C^2$ -piecewise with finite number of vertexes, denote by  $\theta_{\Omega}$  the minimum interior angle among all the vertexes. Then we have the following Moser–Trudinger inequality (cf. [43, 44]):

$$\int_{\Omega} e^{2\theta_{\Omega} \left( \frac{u - \bar{u}}{\|\nabla u\|_2} \right)^2} dx \leq C_0, \quad \forall u \in H^1(\Omega).$$

As a consequence, we have

$$\begin{aligned} \log \int_{\Omega} e^u dx &= \frac{1}{|\Omega|} \int_{\Omega} u dx + \log \int_{\Omega} e^{u-\bar{u}} dx \\ &\leq \frac{1}{|\Omega|} \int_{\Omega} u dx + \log \int_{\Omega} e^{\frac{1}{8\theta_{\Omega}} \|\nabla u\|_2^2 + 2\theta_{\Omega} \frac{(u-\bar{u})^2}{\|\nabla u\|_2^2}} dx \\ &\leq \frac{\beta}{8\theta_{\Omega}} \int_{\Omega} |u|^2 dx + \frac{1}{8\theta_{\Omega}} \int_{\Omega} |\nabla u|^2 dx + \frac{2\theta_{\Omega}}{\beta|\Omega|} + \log C_0, \end{aligned}$$

and it implies that

$$\frac{1}{2} \int_{\Omega} |\nabla u|^2 dx + \frac{\beta}{2} \int_{\Omega} |u|^2 dx - 4\theta_{\Omega} \log \left( \int_{\Omega} e^u dx \right) \geq -C_1, \quad u \in H^1(\Omega). \tag{4.9}$$

Now fixing  $P \neq 0$ , we consider a fundamental domain

$$\Pi(P, \theta) := \left\{ x \in B \setminus \{0\} : 0 < \arccos \left( \frac{x \cdot OP}{\|x\| \|OP\|} \right) < \frac{\theta}{2} \right\},$$

and split  $\Pi(P, \theta)$  as follows:

$$\begin{aligned} \Pi_+(P, \theta) &:= \left\{ x \in \Sigma : x \cdot R_{\frac{\theta}{2}}(OP) > 0 \right\}, \\ \Pi_-(P, \theta) &:= \left\{ x \in \Sigma : x \cdot R_{\frac{\theta}{2}}(OP) < 0 \right\}. \end{aligned}$$

In  $\Pi$ , for any  $u \in H^{(R_{2\pi/m})}$ , we have

$$\begin{aligned} J_{\lambda}(u) &:= \frac{1}{2} \int_B |\nabla u|^2 dx + \frac{\beta}{2} \int_B |u|^2 dx - \lambda \log \left( \int_B e^u dx \right) \\ &= m \left( \frac{1}{2} \int_{\Pi} |\nabla u|^2 dx + \frac{\beta}{2} \int_{\Pi} |u|^2 dx - \frac{\lambda}{m} \log \left( \int_{\Pi} e^u dx \right) \right). \end{aligned} \tag{4.10}$$

For the domain  $\Pi$ , we apply the inequality (4.9) with  $\theta_{\Pi}$  given by

$$\theta_{\Pi} = \begin{cases} \frac{\pi}{2}, & \text{if } m = 2, 3, \\ \frac{2\pi}{m}, & \text{if } m \geq 4. \end{cases}$$

Then we see that (4.10) is uniformly bounded from below if

$$\frac{\lambda}{m} \leq 4\theta_{\Pi} = \begin{cases} 2\pi, & \text{if } m = 2, 3, \\ \frac{8\pi}{m}, & \text{if } m \geq 4, \end{cases}$$

which is equivalent to

$$\lambda \leq \begin{cases} 2m\pi, & \text{if } m = 2, 3, \\ 8\pi, & \text{if } m \geq 4. \end{cases}$$

This proves that (4.10) is uniformly bounded from below provided  $\lambda \leq 8\pi$  and  $m \geq 4$ .

For  $m = 2, 3$ , using the Steiner symmetrization, we can assume that the function  $u$  is radial. As a consequence, the fundamental domain  $\Pi$  is replaced by  $\Pi_+$  and

$$\begin{aligned}
 J_\lambda(u) &:= \frac{1}{2} \int_B |\nabla u|^2 dx + \frac{\beta}{2} \int_B |u|^2 dx - \lambda \log \left( \int_B e^u dx \right) \\
 &= 2m \left( \frac{1}{2} \int_{\Pi_+} |\nabla u|^2 dx + \frac{\beta}{2} \int_{\Pi_+} |u|^2 dx - \frac{\lambda}{2m} \log \left( \int_{\Pi} e^u dx \right) \right).
 \end{aligned}
 \tag{4.11}$$

Notice that in  $\Pi_+$ , the constant  $\theta_{\Pi_+} = \frac{\pi}{m}$ . Using (4.9), we see (4.11) is bounded from below if

$$\frac{\lambda}{2m} \leq 4 \frac{\pi}{m},$$

which implies  $\lambda \leq 8\pi$ . Thus we have deduced that the functional  $J_\lambda(u)$  restricted to the space  $H^G$  is bounded from below when  $\lambda \leq 8\pi$ . From the standard variational argument we can find  $u_{\min} \in H^{(R_{2\pi/m})}$  such that

$$J_\lambda(u) \geq J_\lambda(u_{\min}), \quad \forall u \in H^{(R_{2\pi/m})}.$$

Then the remaining assertion of Proposition 4.7 is a direct consequence of Theorem 1.2. □

Now we are ready to prove Theorem 1.3.

*Proof of Theorem 1.3.* Applying Proposition 4.7 and the arguments in [5, Theorem 1.1], we can conclude that any solution  $(v, u) \in H^G \times H^\circ G$  of (1.1) is globally defined whenever  $\lambda < \Lambda_m$  and show that

$$\limsup_{t \rightarrow \infty} (\|v(x, t)\|_{L^\infty(\Omega)} + \|u(x, t)\|_{L^\infty(\Omega)}) < \infty.$$

From the result [13, Theorem 1.1], we get that the classical solutions converge in  $C^1(\bar{B})$  as  $t \rightarrow \infty$  to a stationary solution. Particularly, this convergence holds true for a subsequence  $(t_k)_{k \in \mathbb{N}}$  in the sense that

$$v(t_k) \rightarrow \frac{e^{u_\infty}}{\int_\Omega e^{u_\infty} dx} \text{ in } L^2(B) \text{ as } t_k \rightarrow \infty,$$

and

$$u(t_k) \rightarrow u_\infty \text{ in } H^1(B) \text{ as } t_k \rightarrow \infty,$$

where  $u_\infty$  is a solution of (1.3). It is known by Theorem 1.2 that  $u = \frac{\lambda}{\beta\pi}$  is the only solution to the problem (1.3) provided  $\lambda < \Lambda_m$ . Thus, a solution  $(v, u) \in H^G \times \mathring{H}^G$  of system (1.1) must converge to the constant solution  $(\frac{\lambda}{\pi}, \frac{\lambda}{\beta\pi})$  as  $t \rightarrow \infty$  when  $\lambda < \Lambda_m$ . Finally, if  $\lambda < 8\pi$ , the convergence to the constant equilibrium of any radial solution of (1.1) results from Remark 1.1. □

### 5. Uniqueness of the Dirichlet problem (1.12)

In this section, we shall provide a complete proof of Theorem 1.4. Indeed, we may consider a more general problem:

$$\begin{cases} \Delta u - g(u) + \lambda \frac{e^u}{\int_{\Omega} e^u dx} = 0 & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \tag{5.1}$$

where  $g$  satisfies that  $g(x), g'(x) > 0$  for  $x > 0$ .

Concerning the problem (5.1), we obtain the following conclusion,

**Theorem 5.1.** *Let  $\Omega$  be an open bounded and simply connected set. Assuming that  $u_i \in C^2(\Omega) \cap C(\bar{\Omega}) (i = 1, 2)$  solve the Eq. (5.1) and  $u_1 \not\equiv u_2$ , then  $\lambda > 8\pi$ .*

We prove the above result by following the same spirit as treated for the Neumann problem (1.3). Precisely, we focus on the difference between the integration of  $g(u_2)$  and  $g(u_1)$  with respect to the level set of  $u_2 - u_1$ , instead of the pointwise comparison between  $g(u_2)$  and  $g(u_1)$ . To begin with the argument, let us recall the classical Bol’s isoperimetric inequality, see [42, 45–47] for a detailed history of the Bol’s inequality.

**Theorem 5.A.** *Let  $\Omega \subset \mathbb{R}^2$  be a simply-connected and assume  $w \in C^2(\Omega) \cap C(\bar{\Omega})$  satisfies*

$$\Delta w + e^w \geq 0, \quad \int_{\Omega} e^w dx \leq 8\pi.$$

*Then for every  $\omega \Subset \Omega$  of class  $C^1$  the following inequality holds*

$$\left( \int_{\partial\omega} e^{\frac{w}{2}} dx \right)^2 \geq \frac{1}{2} \left( \int_{\omega} e^w dx \right) \left( 8\pi - \int_{\omega} e^w dx \right).$$

*Moreover, the above inequality is strict if  $\Delta w + e^w > 0$  somewhere in  $\omega$  or  $\omega$  is not simply connected.*

For  $\theta > 0$ , the function defined by

$$U_{\theta}(x) = -2 \log \left( 1 + \frac{\theta^2 |x|^2}{8} \right) + 2 \log \theta \tag{5.2}$$

satisfies the following property

$$\Delta U_{\theta} + e^{U_{\theta}} = 0, \text{ and } \left( \int_{\partial B_r} e^{\frac{U_{\theta}}{2}} dx \right)^2 = \frac{1}{2} \left( \int_{B_r} e^{U_{\theta}} dx \right) \left( 8\pi - \int_{B_r} e^{U_{\theta}} dx \right),$$

for all  $r > 0$  and  $\theta > 0$ , where  $B_r$  denotes the ball of radius  $r$  centered at the origin in  $\mathbb{R}^2$ .

Next, we shall recall some facts about the rearrangement with the measures. Such discussions have been detailed in [33, 34, 36, 45–47], we shall sketch the process here only. For any function  $\phi \in C^2(\bar{\Omega})$  which is constant on  $\partial\Omega$  can be equimeasurably rearranged with respect to the measures  $e^u dx$  and  $e^{U_{\theta}} dx$ , where  $u$  is a function satisfying that  $\Delta u + e^u \geq 0$  and  $U_{\theta}$  is defined in (5.2). For any  $t > \min_{x \in \bar{\Omega}} \phi$ , we define  $\Omega_t^*$  be a ball centered at the origin such that

$$\int_{\Omega_t^*} e^{U_{\theta}} dx = \int_{\{\phi > t\}} e^u dx.$$

Then we define  $\phi^* : \Omega^* \rightarrow \mathbb{R}$  by

$$\phi^*(x) := \sup\{t \in \mathbb{R} : x \in \Omega_t^*\},$$

which gives an equimeasurable rearrangement of  $\phi$  with respect to the measures  $e^u dx$  and  $e^{U_0} dx$  :

$$\int_{\{\phi^* > t\}} e^{U_0} dx = \int_{\{\phi > t\}} e^u dx, \quad \forall t > \min_{x \in \Omega} \phi.$$

For functions  $\phi$  and  $\phi^*$ , by using Proposition 5.A, the following conclusions hold.

**Proposition 5.B.** [36] *Let  $u \in C^{0,1}(\bar{\Omega})$  satisfy*

$$\Delta u + e^u \geq 0 \text{ in } \Omega,$$

*and let  $U_\theta$  be given in (5.2). Suppose  $\phi \in C^2(\bar{\Omega})$  and  $\phi \equiv C$  on  $\partial\Omega$ . Let  $\phi^*$  be the equimeasurable symmetric rearrangement of  $\phi$  with respect to the measures  $e^u dx$  and  $e^{U_0} dx$ , then it holds for all  $t > \min_{x \in \bar{\Omega}} \phi(x)$  that*

$$\int_{\{\phi^* = t\}} |\nabla \phi^*| d\sigma \leq \int_{\{\phi = t\}} |\nabla \phi| d\sigma.$$

The following two lemmas are the consequences of the Bols's inequality and reversed Bol's inequality in the radial setting respectively.

**Lemma 5.C.** [36] *Assume that  $\psi \in C^{0,1}(\bar{B}_R)$  is a strictly decreasing, radial, Lipschitz function and satisfies*

$$\int_{\partial B_r} |\nabla \psi| d\sigma \leq \int_{B_r} e^\psi dx \tag{5.3}$$

*a.e.  $r \in (0, R)$  and  $\psi = U_{\theta_1} = U_{\theta_2}$  for some  $\theta_2 > \theta_1$  on  $\partial B_R$ . Then there holds*

$$\text{either } \int_{B_R} e^\psi dx \leq \int_{B_R} e^{U_{\theta_1}} dx \quad \text{or} \quad \int_{B_R} e^\psi dx \geq \int_{B_R} e^{U_{\theta_2}} dx. \tag{5.4}$$

*Moreover if the inequality in (5.3) is strict in a set with positive measure in  $(0, R)$ , then the inequalities in (5.4) are also strict.*

**Lemma 5.D.** [39] *Assume that  $\psi \in C^{0,1}(\mathbb{R}^2 \setminus B_R)$  is a strictly decreasing, radial, Lipschitz function, and satisfies*

$$\int_{\partial B_r} |\nabla \psi| d\sigma \leq 8\pi - \int_{\mathbb{R}^2 \setminus B_r} e^\psi dx \tag{5.5}$$

*a.e.  $r \in (R, +\infty)$  and  $\psi = U_{\theta_1} = U_{\theta_2}$  for some  $\theta_2 > \theta_1$  on  $\partial B_R$ . Then there holds*

$$\int_{\mathbb{R}^2 \setminus B_R} e^{U_{\theta_2}} dx \leq \int_{\mathbb{R}^2 \setminus B_R} e^\psi dx \leq \int_{\mathbb{R}^2 \setminus B_R} e^{U_{\theta_1}} dx. \tag{5.6}$$

*Moreover if the inequality in (5.5) is strict in a set with positive measure in  $(R, +\infty)$ , then the inequalities in (5.6) are also strict.*

We shall also need the following lemma.

**Lemma 5.E.** [39] Assume that  $\psi \in C^{0,1}(\overline{B_R})$  is a strictly decreasing and radial function satisfying (5.3) for a.e.  $r \in (0, R)$ . If

$$\int_{B_R} e^\psi dx = \int_{B_R} e^{U_\theta} dx < 8\pi,$$

then  $U_\theta(R) \leq \psi(R)$ .

Now we are ready to prove Theorem 5.1.

*Proof of Theorem 5.1.* Without loss of generality, we may assume  $\int_\Omega g(u_2) dx \geq \int_\Omega g(u_1) dx$ . Let

$$v_i = u_i + \log \lambda - \log \int_\Omega e^{u_i} dx, \quad i = 1, 2.$$

Then  $v_i$  ( $i = 1, 2$ ) satisfy the following equation

$$\Delta v_i + e^{v_i} = g(u_i), \quad i = 1, 2, \tag{5.7}$$

where  $\int_\Omega e^{v_1} = \int_\Omega e^{v_2} = \lambda$ . It is not difficult to see that  $v_1 \neq v_2$ . Indeed, if  $v_1 \equiv v_2$ , then we get  $g(u_1) = g(u_2)$  by (5.7). Combined with the fact  $g(u)$  is strictly increasing, we get  $u_1 = u_2$  and contradiction arises. Thus,  $v_1 \neq v_2$ .

Set

$$t_i = \inf_\Omega (v_2 - v_1) \quad \text{and} \quad t_s = \sup_\Omega (v_2 - v_1).$$

Then we claim that

$$\int_{\{v_2 - v_1 > t\}} (g(u_2) - g(u_1)) dx \geq 0, \quad \forall t \in [t_i, t_s]. \tag{5.8}$$

We notice that

$$\int_{\{v_2 - v_1 > t\}} (g(u_2) - g(u_1)) dx = \int_{\{u_2 - u_1 > t^*\}} (g(u_2) - g(u_1)) dx,$$

where

$$t^* = t + \log \int_\Omega e^{u_2} dx - \log \int_\Omega e^{u_1} dx.$$

If  $t^* \geq 0$ , then there is nothing to prove, while if  $t^* < 0$ , we have

$$\int_{\{u_2 - u_1 > t^*\}} (g(u_2) - g(u_1)) dx = \int_\Omega (g(u_2) - g(u_1)) dx - \int_{\{u_2 - u_1 \leq t^*\}} (g(u_2) - g(u_1)) dx \geq 0.$$

Thus we have proved the claim.

Next we divide our arguments into two steps.

Step 1. We prove that  $\lambda \geq 8\pi$ . Suppose  $\lambda < 8\pi$ , we choose  $\theta > 0$  and  $R \in (0, \infty)$  such that

$$\int_\Omega e^{v_1} dx = \int_{B_R} e^{U_\theta} dx,$$

and let  $\phi$  be the symmetrization of  $v_2 - v_1$  with respect to the measure  $e^{v_1} dx$  and  $e^{U_\theta} dx$ . Then it follows from Proposition 5.B and Fubini's theorem that

$$\begin{aligned} \int_{\{\phi=t\}} |\nabla \phi| d\sigma &\leq \int_{\{v_2-v_1=t\}} |\nabla(v_2-v_1)| d\sigma = - \int_{\{v_2-v_1>t\}} \Delta(v_2-v_1) dx \\ &= \int_{\{v_2-v_1>t\}} (e^{v_2} - e^{v_1}) dx - \int_{\{v_2-v_1>t\}} (g(u_2) - g(u_1)) dx \\ &\leq \int_{\{v_2-v_1>t\}} (e^{v_2} - e^{v_1}) dx = \int_{\{\phi>t\}} e^{U_\theta+\phi} dx - \int_{\{\phi>t\}} e^{U_\theta} dx \\ &= \int_{\{\phi>t\}} e^{U_\theta+\phi} dx - \int_{\{\phi=t\}} |\nabla U_\theta| d\sigma, \end{aligned}$$

for  $t \in (t_b, t_s]$ , where  $t_b = (v_2 - v_1)|_{\partial\Omega}$  and we used (5.8). If  $t < t_b$ , we find that

$$\partial\{v_2 - v_1 > t\} = \{v_2 - v_1 = t\} \cup \partial\Omega.$$

On the other hand, we notice that  $v_2 - v_1 \leq t$  for  $t < t_b$  implies that  $u_2 - u_1 \leq 0$ , so  $g(u_2) - g(u_1) \leq 0$  in  $\{v_2 - v_1 \leq t\}$  for  $t < t_b$ . Then

$$\begin{aligned} \int_{\{\phi=t\}} |\nabla \phi| d\sigma &\leq \int_{\{v_2-v_1=t\}} |\nabla(v_2-v_1)| d\sigma \\ &= - \int_{\{v_2-v_1>t\}} \Delta(v_2-v_1) dx + \int_{\partial\Omega} \partial_\nu(v_2-v_1) d\sigma \\ &= \int_{\{v_2-v_1>t\}} (e^{v_2} - e^{v_1}) dx + \int_{\{v_2-v_1\leq t\}} (g(u_2) - g(u_1)) dx \\ &\leq \int_{\{v_2-v_1>t\}} (e^{v_2} - e^{v_1}) dx = \int_{\{\phi>t\}} e^{U_\theta+\phi} dx - \int_{\{\phi>t\}} e^{U_\theta} dx \\ &= \int_{\{\phi>t\}} e^{U_\theta+\phi} dx - \int_{\{\phi=t\}} |\nabla U_\theta| d\sigma, \end{aligned}$$

for  $t \in [t_i, t_b)$ , where we used

$$\begin{aligned} \int_{\partial\Omega} \partial_\nu(v_2-v_1) d\sigma &= \int_\Omega \Delta(v_2-v_1) dx = \int_\Omega (g(u_2) - g(u_1)) dx - \int_\Omega (e^{v_2} - e^{v_1}) dx \\ &= \int_\Omega (g(u_2) - g(u_1)) dx. \end{aligned}$$

Hence

$$\int_{\{\phi=t\}} |\nabla(U_\theta + \phi)| d\sigma \leq \int_{\{\phi>t\}} e^{U_\theta+\phi} dx$$

for all a.e.  $t > t_i$ . Since  $\phi$  is decreasing in  $r$ ,  $\psi := U_\theta + \phi$  is strictly decreasing function, and

$$\int_{\partial B_r} |\nabla \psi| d\sigma \leq \int_{B_r} e^\psi dx, \text{ a.e. } r \in (0, R).$$

Since  $v_1 \neq v_2$  and  $\int_\Omega e^{v_1} dx = \int_\Omega e^{v_2} dx$ , then  $v_2 < v_1$  on a subset of  $\Omega$  with positive measure. Hence  $\phi(R) < 0$ , therefore

$$\psi(R) = U_\theta(R) + \phi(R) < U_\theta(R).$$

It is a contradiction by Lemma 5.E., and therefore  $\lambda \geq 8\pi$ .

Step 2. We prove  $\lambda \neq 8\pi$ . Suppose  $\lambda = 8\pi$  and let  $\theta_1 > 0$ . From the above arguments we can find there exists  $\psi = U_{\theta_1} + \phi \in C^{0,1}(\mathbb{R}^2)$  such that

$$\int_{\Omega} e^{v_1} dx = \int_{\mathbb{R}^2} e^{U_{\theta_1}} dx = 8\pi = \int_{\Omega} e^{v_2} dx = \int_{\mathbb{R}^2} e^{\psi} dx,$$

and

$$\int_{\partial B_r} |\nabla \psi| d\sigma \leq \int_{B_r} e^{\psi} dx \text{ for a.e. } r \in (0, \infty).$$

Since  $\int_{\mathbb{R}^2} e^{\psi} dx = \int_{\mathbb{R}^2} e^{U_{\theta_1}} dx$ , there exists  $r_0 \in (0, \infty)$  such that  $\psi(r_0) = U_{\theta_1}(r_0)$ . Let  $\theta_2 = \frac{8}{r_0^2 \theta_1}$ , then it is easy to see that

$$U_{\theta_2}(r_0) = U_{\theta_1}(r_0) = \psi(r_0)$$

by the expression of  $U_\theta$  in (5.2). We notice that  $\psi > U_{\theta_1}$  in  $B_{r_0}$ , it follows from Lemma 5.C that  $\theta_1 < \theta_2$  and

$$\int_{B_{r_0}} e^{\psi} dx \geq \int_{B_{r_0}} e^{U_{\theta_2}} dx.$$

While in  $\mathbb{R}^2 \setminus B_{r_0}$ , we have  $\psi < U_{\theta_1}$  and it follows from Lemma 5.D that

$$\int_{\mathbb{R}^2 \setminus B_{r_0}} e^{\psi} dx \geq \int_{\mathbb{R}^2 \setminus B_{r_0}} e^{U_{\theta_2}} dx.$$

Hence

$$8\pi = \lambda = \int_{\mathbb{R}^2} e^{\psi} dx \geq \int_{\mathbb{R}^2} e^{U_{\theta_2}} dx = 8\pi. \tag{5.9}$$

Since the solution  $u$  of (5.1) is positive, we have  $\Delta v_i + e^{v_i} > 0$ ,  $i = 1, 2$ , as a consequence, the inequality in (5.9) is strict by Lemma 5.C and Lemma 5.D. Thus  $\lambda \neq 8\pi$  and it finishes the proof.  $\square$

Before proving Theorem 1.4, we shall derive the degree counting formula for (1.12) in  $H_0^1(\Omega)^1$ .

For any solution of (1.12) in  $H_0^1(\Omega)$ , an inequality of Brezis and Strauss [48] asserts that (also see [49, Lemma 2.3])

$$\|u\|_{W^{1,q}(\Omega)} \leq C \text{ for } 1 < q < 2 \text{ with a constant } C = C_q > 0.$$

We decompose  $u = w + v$ , where  $w, v$  satisfy

$$\Delta w - \beta u = 0 \text{ in } \Omega, \quad w = 0 \text{ on } \partial\Omega,$$

and

$$\Delta v + \lambda \frac{he^v}{\int_{\Omega} he^v dx} = 0 \text{ in } \Omega, \quad v = 0 \text{ on } \partial\Omega, \tag{5.10}$$

<sup>1</sup> $H_0^1(\Omega) = \{u \in H^1(\Omega) : u = 0 \text{ on } \partial\Omega\}$



respectively, where  $h = e^w$ . Using the standard elliptic estimate and the fact  $u \in W^{1,q}(\Omega)$ , we get  $w \in W^{3,q}(\Omega)$  and  $\|w\|_{C^1(\Omega)} \leq C$ . We denote  $H = \lambda \frac{h}{\int_{\Omega} h e^v dx}$  and it is easy to see that  $\|H\|_{C^1(\Omega)} \leq C$ . If there is a sequence of solutions  $\{u_n\}$  of (1.12) with  $\lambda = \lambda_n$  such that  $\max_{\Omega} u_n \rightarrow +\infty$  as  $n \rightarrow +\infty$ , then for Eq. (5.10) with  $\lambda$  replaced by  $\lambda_n$ , we get  $\max_{\Omega} v_n \rightarrow +\infty$  as  $n \rightarrow +\infty$ . By [50, Theorem 1], we have  $\lambda_n \rightarrow 8m\pi$  for some  $m \in \mathbb{N}$ . As a consequence, if  $\lambda \neq 8m\pi$ , then any solution  $v$  of (5.10) is uniformly bounded, and we can find a constant  $C > 0$  such that  $\|u\|_{L^\infty(\Omega)} \leq C$  for any solution of (1.12). Set  $T(\lambda)$  to be

$$T(\lambda) = \Delta^{-1} \left( \lambda \frac{e^u}{\int_{\Omega} e^u dx} - \beta u \right),$$

which acts in  $H_0^1(\Omega)$ . Then the Leray–Schauder degree

$$d_\lambda := \text{deg}(I + T(\lambda), B_R, 0)$$

is well-defined for  $\lambda \neq 8m\pi$  and

$$B_R = \{u \in H_0^1(\Omega) : \|u\|_{H^1} \leq R\} \text{ with } R \text{ sufficiently large.}$$

Next, we introduce a homotopy deformation for (1.12)

$$\begin{cases} \Delta u - t\beta u + \lambda \frac{e^u}{\int_{\Omega} e^u dx} = 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases} \tag{5.11}$$

When  $t = 1$ , Eq. (5.11) is (1.12), while if  $t = 0$ , Eq. (5.11) is the mean field equation on bounded domain. It is not difficult to see that as  $t$  changes from 1 to 0, we can argue as above to get that the solution  $u_t$  of (5.11) is uniformly bounded provided  $\lambda \neq 8m\pi$ . Therefore the Leray–Schauder degree of the above equation is the same for  $t = 1$  and  $t = 0$ . Together with the degree formula for the mean field equation on bounded domain (see [20, Theorem 1.3]), we get

**Theorem 5.2.** *Let  $\Omega$  be a non-empty bounded open set and  $8m\pi < \lambda < 8(m + 1)\pi$  for some positive integer  $m$ . Then the Leray–Schauder degree of (1.12)  $d_\lambda = \binom{m + g - 1}{m}$ , where  $g$  denotes the number of holes in  $\Omega$ . While if  $0 \leq \lambda < 8\pi$ ,  $d_\lambda = 1$ .*

Next we prove Theorem 1.4 by Theorems 5.1 and 5.2.

*Proof of Theorem 1.4.* First we claim that all the solutions to the Eq. (1.12) are positive. Indeed, if  $u$  is not positive in  $\Omega$ , letting  $p$  be the point in  $\Omega$  where  $u$  obtains its minimal value, then we have  $u(p) \leq 0$  and  $\Delta u(p) \geq 0$ . Therefore

$$\Delta u - \beta u + \lambda \frac{e^u}{\int_{\Omega} e^u dx} > 0 \text{ at } p,$$

which gives a contradiction. Hence, the claim is true. Next, it is easy to see that  $g(u) = u$  satisfies the condition  $g(x), g'(x) > 0$  for  $x > 0$ . Then Theorem 1.4 follows by Theorems 5.1 and 5.2. Thus we finish the proof. □

## Funding

Part of this work was finished when the third author was a postdoctoral fellow in the Department of Applied Mathematics in the Hong Kong Polytechnic University supported by the project G-YBKT. The research of J. Wang is supported by NSFC of China (Grants 11571140, 11671077), Fellowship of Outstanding Young Scholars of Jiangsu Province (BK20160063), The Six big talent peaks project in Jiangsu Province (XYDXX-015) and NSF of Jiangsu Province (BK20150478). The research of Z. Wang is supported by the Hong Kong Polytechnic University internal grant PolyU 153041/15P. The research of W. Yang is supported by CAS Pioneer Hundred Talents Program (Y8Y3011001) and NSFC No. 11801550.

## References

- [1] Keller, E. F., Segel, L. A. (1970). Initiation of slime mold aggregation viewed as an instability. *J. Theor. Biol.* 26(3):399–415.
- [2] Keller, E. F., Segel, L. A. (1971). Traveling bands of chemotactic bacteria: A theoretical analysis. *J. Theor. Biol.* 30(2):235–248.
- [3] Hillen, T., Potapov, A. (2004). The one-dimensional chemotaxis model: Global existence and asymptotic profile. *Math. Methods Appl. Sci.* 27(15):1783–1801. DOI: [10.1002/mma.569](https://doi.org/10.1002/mma.569).
- [4] Osaki, K., Yagi, A. (2001). Finite dimensional attractors for one-dimensional Keller–Segel equations. *Funkcial. Ekvacioj. Ser. Int.* 44:441–469.
- [5] Nagai, P., Senba, T., Yoshida, K. (1997). Application of the Moser-Trudinger inequality to a parabolic system of chemotaxis. *Funkc. Ekvacioj. Ser. Int.* 40:411–433.
- [6] Herrero, M. A., Velázquez, J. J. L. (1996). Chemotactic collapse for the Keller–Segel model. *J. Math. Biol.* 35(2):177–194. DOI: [10.1007/s002850050049](https://doi.org/10.1007/s002850050049).
- [7] Horstmann, D., Wang, G. (2001). Blow-up in a chemotaxis model without symmetric assumptions. *Eur. J. Appl. Math.* 12:159–177.
- [8] Winkler, M. (2013). Finite-time blow-up in the higher-dimensional parabolic–parabolic Keller–Segel system. *J. Math. Pure. Appl.* 100(5):748–767. DOI: [10.1016/j.matpur.2013.01.020](https://doi.org/10.1016/j.matpur.2013.01.020).
- [9] Bellomo, N., Bellouquid, A., Tao, Y. S., Winkler, M. (2015). Towards a mathematical theory of Keller–Segel models of pattern formation in biological tissues. *Math. Models Methods Appl. Sci.* 25(9):1663–1763. DOI: [10.1142/S021820251550044X](https://doi.org/10.1142/S021820251550044X).
- [10] Hillen, T., Painter, K. J. (2009). A user’s guide to PDE models for chemotaxis. *J. Math. Biol.* 58(1–2):183–217. DOI: [10.1007/s00285-008-0201-3](https://doi.org/10.1007/s00285-008-0201-3).
- [11] Horstmann, D. (2003). From 1970 until present: The Keller–Segel model in chemotaxis and its consequences. I. *Jahresber. Deutsch. Math.-Verien.* 105:103–106.
- [12] Perthame, B. (2007). *Transport Equations in Biology: Frontiers in Mathematics*. Basel: Birkhäuser Verlag.
- [13] Feireisl, E., Laurençot, P., Petzeltová, H. (2007). On convergence to equilibria for the Keller–Segel chemotaxis model. *J. Diff. Eq.* 236(2):551–569. DOI: [10.1016/j.jde.2007.02.002](https://doi.org/10.1016/j.jde.2007.02.002).
- [14] Moser, J. (1970/71). A sharp form of an inequality by N. Trudinger. *Indiana Univ Math. J.* 20:1077–1092.
- [15] Caglioti, E., Lions, P. L., Marchioro, C., Pulvirenti, M. (1992). A special class of stationary flows for two-dimensional euler equations: A statistical mechanics description. *Commun. Math. Phys.* 143(3):501–525. DOI: [10.1007/BF02099262](https://doi.org/10.1007/BF02099262).
- [16] Chanillo, S., Kiessling, M. (1994). Rotational symmetry of solutions of some nonlinear problems in statistical mechanics and in geometry. *Commun. Math. Phys.* 160(2):217–238. DOI: [10.1007/BF02103274](https://doi.org/10.1007/BF02103274).

- [17] Kiessling, M. (1993). Statistical mechanics of classical particles with logarithmic interactions. *Commun. Pure Appl. Math.* 46(1):27–56. DOI: [10.1002/cpa.3160460103](https://doi.org/10.1002/cpa.3160460103).
- [18] Tarantello, G. (1996). Multiple condensate solutions for the Chern–Simons–Higgs theory. *J. Math. Phys.* 37:769–796.
- [19] Struwe, M., Tarantello, G. (1998). On multiplevortex solutions in Chern–Simons gauge theory. *Boll. U.M.I.* 8:109–121.
- [20] Chen, C. C., Lin, C. S. (2003). Topological degree for a mean field equation on Riemann surfaces. *Commun. Pure Appl. Math.* 56(12):1667–1727. DOI: [10.1002/cpa.10107](https://doi.org/10.1002/cpa.10107).
- [21] Malchiodi, A. (2008). Morse theory and a scalar field equation on compact surfaces. *Adv. Diff. Eq.* 13(11–12):1109–1129.
- [22] Wang, G. F., Wei, J. C. (2002). Steady state solutions of a reaction-diffusion system modeling chemotaxis. *Math. Nachr.* 233–234(1):221–236. DOI: [10.1002/1522-2616\(200201\)233:1<221::AID-MANA221>3.3.CO;2-D](https://doi.org/10.1002/1522-2616(200201)233:1<221::AID-MANA221>3.3.CO;2-D).
- [23] Horstmann, D. (1999). *Aspekte Positiver Chemotaxis*. Dissertation, Univ. Köln.
- [24] Battaglia, L. (2019). A general existence result for stationary solutions to the Keller–Segel system. *Discr Contin. Dyn. Syst.* 39(2):905–926. DOI: [10.3934/dcds.2019038](https://doi.org/10.3934/dcds.2019038).
- [25] Lin, C. S. (2000). Uniqueness of solutions to the mean field equations for the spherical onsager vortex. *Arch. Ration. Mech. Anal.* 153(2):153–176. DOI: [10.1007/s002050000085](https://doi.org/10.1007/s002050000085).
- [26] Onofri, E. (1982). On the positivity of the effective action in a theory of random surfaces. *Commun. Math. Phys.* 86(3):321–326. DOI: [10.1007/BF01212171](https://doi.org/10.1007/BF01212171).
- [27] Lin, C. S., Lucia, M. (2006). Uniqueness of solutions for a mean field equation on torus. *J. Diff. Eq.* 229(1):172–185. DOI: [10.1016/j.jde.2005.11.003](https://doi.org/10.1016/j.jde.2005.11.003).
- [28] Horstmann, D., Lucia, M. (2011). Uniqueness and symmetry of equilibria in a chemotaxis model. *J. Reine Angew. Math.* 654:83–124.
- [29] Senba, T., Suzuki, T. (2000). Some structures of the solution set for a stationary system of chemotaxis. *Adv. Math. Sci. Appl.* 10:191–224.
- [30] Lucia, M. (2009). Isoperimetric profile and uniqueness for Neumann problems. *Ann. Inst. Henri Poincaré Anal. Non Linéaire* 26(1):81–100. DOI: [10.1016/j.anihpc.2007.07.002](https://doi.org/10.1016/j.anihpc.2007.07.002).
- [31] López-Gómez, J., Nagai, T., Yamada, T. (2014). Non-trivial  $\omega$ -limit sets and oscillating solutions in a chemotaxis model in  $\mathbb{R}^2$  with critical mass. *J. Funct. Anal.* 266:3455–3507. DOI: [10.1016/j.jfa.2014.01.015](https://doi.org/10.1016/j.jfa.2014.01.015).
- [32] López-Gómez, J., Nagai, T., Yamada, T. (2013). The basin of attraction of the steady-states for a chemotaxis model in  $\mathbb{R}^2$  with critical mass. *Arch. Rational Mech. Anal.* 207(1):159–184. DOI: [10.1007/s00205-012-0560-1](https://doi.org/10.1007/s00205-012-0560-1).
- [33] Suzuki, T. (1992). Global analysis for a two dimensional elliptic eigenvalue problem with exponential nonlinearity. *Ann. Inst. Henri Poincaré Anal. Non Linéaire* 9(4):367–398. DOI: [10.1016/S0294-1449\(16\)30232-3](https://doi.org/10.1016/S0294-1449(16)30232-3).
- [34] Chang, S. Y. A., Chen, C. C., Lin, C. S. (2003). *Extremal Functions for a Mean Field Equation in Two Dimension. Lecture on Partial Differential Equations*, Vol. 2: *New Studies on Advanced Mathematics*. Somerville: Int. Press, 6193 p.
- [35] Bartolucci, D., Lin, C. S. (2014). Existence and uniqueness for mean field equations on multiply connected domains at the critical parameter. *Math. Ann.* 359(1–2):1–44. DOI: [10.1007/s00208-013-0990-6](https://doi.org/10.1007/s00208-013-0990-6).
- [36] Gui, C. F., Moradifam, A. (2018). The sphere covering inequality and its applications. *Invent. Math.* 214(3):1169–1204. DOI: [10.1007/s00222-018-0820-2](https://doi.org/10.1007/s00222-018-0820-2).
- [37] Gui, C. F., Jevnikar, A., Moradifam, A. (2018). Symmetry and uniqueness of solutions to some liouville-type problems: Asymmetric sinh–Gordon equation, cosmic string equation and toda system. *Commun. Part. Diff. Eq.* 43(3):428–447. DOI: [10.1080/03605302.2018.1446164](https://doi.org/10.1080/03605302.2018.1446164).
- [38] Gui, C. F., Moradifam, A. (2018). Symmetry of solutions of a mean field equation on flat tori. *Int. Math. Res. Notices.* 2019(3):799–809.
- [39] Gui, C. F., Moradifam, A. Uniqueness of solutions of mean field equations in  $\mathbb{R}^2$ . *Proc. Am. Math. Soc.* 146(3):1231–1124. DOI: [10.1090/proc/13814](https://doi.org/10.1090/proc/13814).

- [40] Shi, Y. G., Sun, J. C., Tian, G., Wei, D. Y. (2019). Uniqueness of the mean field equation and rigidity of Hawking Mass. *Calc. Var.* 58(2). DOI: [10.1007/s00526-019-1496-1](https://doi.org/10.1007/s00526-019-1496-1).
- [41] Giusti, E. (1984). *Minimal Surfaces and Functions of Bounded Variation, Monographs in Mathematics*, Vol. 80. Basel: Birkhäuser Verlag.
- [42] Burago, Y. D., Zalgaller, V. A. (1988). *Geometric Inequalities, Grundlehren Der Mathematischen Wissenschaften*, Vol. 285. Berlin: Springer-Verlag.
- [43] Chang, A., Yang, P. C. (1988). Conformal deformation of metrics on  $S^2$ . *J. Diff. Geom.* 27(2):259–296. DOI: [10.4310/jdg/1214441783](https://doi.org/10.4310/jdg/1214441783).
- [44] Cianchi, A. (2005). Moser-Trudinger inequalities without boundary conditions and isoperimetric problems. *Indiana Univ. Math. J.* 54(3):669–705. DOI: [10.1512/iumj.2005.54.2589](https://doi.org/10.1512/iumj.2005.54.2589).
- [45] Bandle, C. (1980). *Isoperimetric Inequalities and Applications*. London, UK: Pitman
- [46] Bartolucci, D., Lin, C. S. (2009). Uniqueness results for mean field equations with singular data. *Commun. Part. Diff. Eq.* 34(7):676–702. DOI: [10.1080/03605300902910089](https://doi.org/10.1080/03605300902910089).
- [47] Bol, G. (1941). Isoperimetrische ungleichungen für bereiche auf flächen. *Jber. Deutsch. Math. Verein.* 51:219–257.
- [48] Brezis, H., Strauss, W. A. (1973). Semi-linear second order elliptic equations in  $L^1$ . *J. Math. Soc. Jpn.* 25:565–590.
- [49] Ding, W., Jost, J., Li, J., Wang, G. F. (1997). The differential equation  $\Delta u = 8\pi - 8\pi h e^u$  on a compact Riemann surface. *Asian J. Math.* 1:653–666.
- [50] Ma, L., Wei, J. C. (2001). Convergence for a liouville equation. *Comment. Math. Helv.* 76(3):506–514. DOI: [10.1007/PL00013216](https://doi.org/10.1007/PL00013216).

## Appendix

**Lemma 6.1.** *Let  $u$  be a solution to the boundary value problem (1.3) with  $\beta > 0$ . Then*

$$u > 0 \text{ in } \Omega.$$

*Proof.* Let

$$a = \min_{x \in \Omega} u(x).$$

We shall prove  $a \geq 0$  by contradiction. Suppose  $a < 0$ , we first prove the value  $a$  cannot be obtained by  $u$  in  $\Omega$ , indeed if  $u(x_0) = a$  with  $x_0 \in \Omega$ , we get

$$\Delta u(x_0) - \beta u(x_0) + \lambda \frac{e^{u(x_0)}}{\int_{\Omega} e^u dx} > 0,$$

contradiction arises. Hence,  $a$  can only be obtained by  $u$  on  $\partial\Omega$ , say at  $x_0 \in \partial\Omega$ . By the continuity of  $u$ , we can always find  $\Omega' \subset \Omega$  with  $x_0 \in \bar{\Omega}' \cap \bar{\Omega}$  and  $u(x) \leq 0$  in  $\Omega'$ . As a consequence, we can see that

$$\Delta u = \beta u - \lambda \frac{e^u}{\int_{\Omega} e^u dx} \leq 0 \text{ in } \Omega'.$$

By the Hopf boundary lemma, we get  $\frac{\partial u}{\partial \nu}(x_0) < 0$ , which contradicts to the Neumann boundary condition. Thus, the assumption is not true and  $a \geq 0$ . On the other hand, by the above arguments we can also show that  $u(x)$  cannot reach the value  $a$  in  $\Omega$  if  $a = 0$ . Therefore  $u(x) > 0$  in  $\Omega$  and the lemma is proved.  $\square$