

Eigenvalues of Nonnegative Tensors

Liqun Qi and Qingzhi Yang

January 9, 2012

A. Definitions of Eigenvalues of Tensors

- [1] L.-H. Lim, “Singular values and eigenvalues of tensors: a variational approach”, *Proceedings of the IEEE International Workshop on Computational Advances in Multi-Sensor Adaptive Processing (CAMSAP '05)*, 1 (2005) 129-132.
- [2] L. Qi, “Eigenvalues of a real supersymmetric tensor”, *J. Symbolic Computation*, 40 (2005) 1302-1324.

B. Generalization of the Perron-Frobenius Theorem

- [3] K.C. Chang, K. Pearson and T. Zhang, “Perron Frobenius Theorem for nonnegative tensors”, *Comm. Math. Sci.*, 6 (2008) 507-520.
- [4] K.C. Chang, K. Pearson and T. Zhang, “Primitivity, the convergence of the NZQ method, and the largest eigenvalue for nonnegative tensors”, *SIAM Journal on Matrix Analysis and Applications*, 32 (2011) 806-819.
- [5] S. Friedland, S. Gaubert and L. Han, “Perron-Frobenius theorem for nonnegative multilinear forms and extensions”, to appear in: *Linear Algebra and Applications*. <http://www.sciencedirect.com/science/article/pii/S0024379511001765>
- [6] K.J. Pearson, “Essentially positive tensors”, *International Journal of Algebra*, 9 (2010) 421-427.
- [7] Q. Yang and Y. Yang, “Further results for Perron-Frobenius Theorem for nonnegative tensors II”, *SIAM Journal on Matrix Analysis*, 32 (2011) 1236-1250.
- [8] Y. Yang and Q. Yang, “Further results for Perron-Frobenius Theorem for nonnegative tensors”, *SIAM Journal on Matrix Analysis*, 31 (2010) 2517-2530.
- [9] Y. Yang and Q. Yang, “A note on the geometric simplicity of the spectral radius of nonnegative irreducible tensors”, Arxiv preprint arXiv:1101.2479, 2011.

C. Algorithms and Convergence Analysis

- [10] S. Hu, Z.H. Huang and L. Qi, “Finding the spectral radius of a nonnegative tensor”, arXiv: 1111.2138v1 [math.NA] 9 Nov 2011. <http://www.polyu.edu.hk/ama/staff/new/qilq/HHQ.pdf>
- [11] Y. Liu, G. Zhou and N.F. Ibrahim, “An always convergent algorithm for the largest eigenvalue of an irreducible nonnegative tensor”, *Journal of Computational and Applied Mathematics*, 235 (2010) 286-292.
- [12] M. Ng, L. Qi and G. Zhou, “Finding the largest eigenvalue of a non-negative tensor”, *SIAM Journal on Matrix Analysis and Applications*, 31 (2009) 1090-1099.
- [13] Y. Yang, Q. Yang and Y. Li, “An algorithm to find the spectral radius of nonnegative tensors and its convergence analysis”, Arxiv preprint arXiv:1102.2668, 2011.
- [14] L. Zhang and L. Qi, “Linear convergence of an algorithm for computing the largest eigenvalue of a nonnegative tensor”, to appear in: *Numerical Linear Algebra with Applications*. <http://www.polyu.edu.hk/ama/staff/new/qilq/ZQ.pdf>

- [15] L. Zhang, L. Qi and Z. Luo, “The dominant eigenvalue of an essentially nonnegative tensor”, arXiv: 1110.6261v1 [math.NA] 28 Oct 2011.
<http://www.polyu.edu.hk/ama/staff/new/qilq/ZQL.pdf>
- [16] L. Zhang, L. Qi and Y. Xu, “Linear convergence of the LZI algorithm for weakly positive tensors”, *Journal of Computational Mathematics*, 30 (2012) 24-33.

D. Singular Values of Nonnegative Rectangular Tensors

- [17] K.C. Chang, L. Qi and G. Zhou, “Singular values of a real rectangular tensor”, *Journal of Mathematical Analysis & Applications*, 370 (2010) 284-294.
- [18] G. Zhou, L. Caccetta and L. Qi, “Convergence of an algorithm for the largest singular value of a nonnegative rectangular tensor”, to appear in: *Linear Algebra and Applications*. <http://www.polyu.edu.hk/ama/staff/new/qilq/ZCQ.pdf>
- [19] Y. Yang and Q. Yang, “Singular values of nonnegative rectangular tensors”, *Frontiers of Mathematics in China*, 6 (2011) 363-378.

E. Eigenvalues of Positive Operators

- [20] K.C. Chang, “A nonlinear Krein Rutman theorem”, *Journal of Systems Science and Complexity*, 22 (2009) 542-554.

F. Spectral Hypergraph Theory

- [21] S.R. Bulò and M. Pelillo, “A generalization of the Motzkin-Straus theorem to hypergraphs”, *Optim. Lett.*, 3 (2009) 187-295.
- [22] S.R. Bulò and M. Pelillo, “New bounds on the clique number of graphs based on spectral hypergraph theory”, in: T. Stützle ed., *Learning and Intelligent Optimization*, Springer Verlag, Berlin, (2009) pp. 45-48.
- [23] J Cooper and A Dutle, “Spectra of Hypergraphs”, Arxiv preprint arXiv:1106.4856, 2011.
- [24] S. Hu and L. Qi, “Algebraic connectivity of an even uniform hypergraph”, to appear in: *Journal of Combinatorial Optimization*. <http://www.polyu.edu.hk/ama/staff/new/qilq/HQ.pdf>

G. Higher-Order Markov Chains

- [25] W. Li and M. Ng, Existence and uniqueness of stationary probability vector of a transition probability tensor, Department of Mathematics, Hong Kong Baptist University, March 2011. <http://www.math.hkbu.edu.hk/mng/tensor-research/tensor-li-ng.pdf>
- [26] W. Li and M. Ng, On linear convergence of power method for computing stationary probability vector of a transition probability tensor, Department of Mathematics, Hong Kong Baptist University, April 2011. <http://www.math.hkbu.edu.hk/mng/tensor-research/tensor-li-ng-conv.pdf>